



**HAL**  
open science

## Uncertainties in the adaptation of alpine pastures to climate change based on remote sensing products and modelling

L. Brilli, Raphaël Martin, G. Argenti, M. Bassignana, M. Bindi, R. Bonet, P. Choler, E. Cremonese, M. Della Vedova, C. Dibari, et al.

### ► To cite this version:

L. Brilli, Raphaël Martin, G. Argenti, M. Bassignana, M. Bindi, et al.. Uncertainties in the adaptation of alpine pastures to climate change based on remote sensing products and modelling. *Journal of Environmental Management*, 2023, 336, pp.117575. 10.1016/j.jenvman.2023.117575 . hal-04068837

HAL Id: hal-04068837

<https://hal.inrae.fr/hal-04068837v1>

Submitted on 14 Apr 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

# Uncertainties in the adaptation of alpine pastures to climate change based on remote sensing products and modelling

Brilli, L.<sup>1,2\*</sup>, Martin, R.<sup>3</sup>, Argenti, G.<sup>2</sup>, Bassignana, M.<sup>4</sup>, Bindi, M.<sup>2</sup>, Bonet, R.<sup>5</sup>, Choler, P.<sup>6</sup>, Cremonese, E.<sup>7</sup>, Della Vedova, M.<sup>5</sup>, Dibari, C.<sup>2</sup>, Filippa, G.<sup>7</sup>, Galvagno, M.<sup>7</sup>, Leolini, L.<sup>2</sup>, Moriondo, M.<sup>1,2</sup>, Piccot, A.<sup>4</sup>, Stendardi, L.<sup>2</sup>, Targetti, S.<sup>8</sup>, Bellocchi, G.<sup>3</sup>

<sup>1</sup> National Research Council – Institute of BioEconomy (IBE-CNR) 50145 Sesto Fiorentino, Italy

<sup>2</sup> University of Florence, DAGRI, 50144 Florence, Italy

<sup>3</sup> Université Clermont Auvergne, INRAE, VetAgro Sup, UREP, 63000 Clermont-Ferrand, France

<sup>4</sup> Institut Agricole Régional, 11100 Aosta, Italy

<sup>5</sup> Parc National des Ecrins, Domaine de Charance, 05000 Gap, France

<sup>6</sup> Université Grenoble Alpes, Université Savoie Mont Blanc, CNRS, LECA, 38000 Grenoble, France

<sup>7</sup> Climate Change Unit, Environmental Protection Agency of Aosta Valley, Saint-Christophe, Italy

<sup>8</sup> University of Bologna, Department of Agricultural and Food Sciences, Viale Fanin, 50 - 40127 Bologna

\*Corresponding author. Tel.: +39 0553033711

E-mail address: [lorenzo.brilli@ibe.cnr.it](mailto:lorenzo.brilli@ibe.cnr.it) (L. Brilli)

## Highlights

- Alpine pastures are vulnerable to climate change.
- Remote sensing and modelling support vulnerability analysis in the western Alps.
- Earlier grazing, not changes in cattle density, copes with increased vulnerability.
- Summer water stresses and warming may lower C sequestration by pastures.

27 **Abstract**

28 Over the last century, the management of pastoral systems has undergone major changes to meet  
29 the livelihood needs of alpine communities. Faced with the changes induced by recent global  
30 warming, the ecological status of many pastoral systems has seriously deteriorated in the western  
31 alpine region. We assessed changes in pasture dynamics by integrating information from remote-  
32 sensing products and two process-based models, i.e. the grassland-specific, biogeochemical growth  
33 model PaSim and the generic crop-growth model DayCent. Meteorological observations and  
34 satellite-derived Normalised Difference Vegetation Index (NDVI) trajectories of three pasture  
35 macro-types (high, medium and low productivity classes) in two study areas - *Parc National des*  
36 *Écrins* (PNE) in France and *Parco Nazionale Gran Paradiso* (PNGP) in Italy - were used as a basis  
37 for the model calibration work. The performance of the models was satisfactory in reproducing  
38 pasture production dynamics ( $R^2=0.52$  to  $0.83$ ). Projected changes in alpine pastures due to  
39 climate-change impacts and adaptation strategies indicate that: i) the length of the growing season  
40 is expected to increase between 15 and 40 days, resulting in changes in the timing and amount of  
41 biomass production, ii) summer water stress could limit pasture productivity; iii) earlier onset of  
42 grazing could enhance pasture productivity; iv) higher livestock densities could increase the rate  
43 of biomass regrowth, but major uncertainties in modelling processes need to be considered; and v)  
44 the carbon sequestration potential of pastures could decrease under limited water availability and  
45 warming.

46

47 *Keywords:* alpine pastures, climate-change adaptation, modelling, remote sensing

48

## 49 **1. Introduction**

50 Mountain pastures are important livelihood systems in the European Alps, with a  
51 multifunctional form of land use encompassing agriculture, outdoor recreation, and tourism as well  
52 as conservation needs (Wanner et al., 2021). Rich in terms of biodiversity (Kurtogullari et al., 2020)  
53 and cultural heritage (Jourdain-Annequin and Duclos, 2006), alpine pastures fulfil economic, social  
54 and environmental functions at the same time (Bengtsson et al., 2019). They provide low-cost  
55 fodder for grazing livestock during the summer period and - where traditional transhumance  
56 systems are present – represent a complementary resource for Mediterranean pastoral systems  
57 (Caballero et al., 2009). Shaped by pastoral activities, alpine pastures have undergone multiple  
58 transformations over the centuries, mainly driven by the fragile balance between maximising  
59 agricultural productivity and the limits imposed by the temporal and spatial dynamics of the climate  
60 and forests-grasslands interactions (Kurz, 2013). However, alpine pastoralism manifests its  
61 fragility in the face of the changes induced by recent global warming. Climate changes and their  
62 impacts are visible in the alpine region, which has experienced a temperature increase of almost 2  
63 °C over the last century, along with an important reduction of precipitation in the summer season  
64 (Gobiet et al., 2014). Specifically, droughts have been one of the main manifestations of climate  
65 variability. Corresponding to periods of abnormally low precipitation, they alter grassland  
66 productivity and quality (Nettier et al., 2010; Dibari et al., 2016) by offsetting the positive effect  
67 of summer heatwaves on canopy greenness (Corona-Lozada et al., 2019), as seen in the European  
68 Alps following a series of droughts (Calanca, 2007). The response of European mountain plant  
69 assemblages to increasing temperatures (thermophilisation) also suggests a progressive decline of  
70 cold-tolerant high-altitude grassland communities (Gottfried et al., 2012) and landscape  
71 modifications with warming-induced upward range shifts (Engler et al., 2011). This may lead to  
72 both a decrease in areas suitable for pasture and a reduction in pasture diversity driven by low-

73 quality vegetation types in the Alpine chain (Dibari et al., 2020), together with changes in grazing  
74 practices (Dibari et al., 2021). This is critical because most impacts on grassland ecosystems can  
75 be related to overgrazing and changes in the timing of livestock transhumance, with high stocking  
76 densities in particular causing a range of negative impacts on plant and animal communities, as  
77 observed in central France (Dumont et al., 2009) and in the Italian Maritime Alps (Negro et al.,  
78 2011).

79 In this context, appropriate management can preserve grassland biodiversity, maintain socio-  
80 ecological systems (Altaweel et al., 2015; Alessa et al., 2018) and counteract climate-change  
81 impacts (Nori and Gemini, 2011; Felber et al., 2016). Specifically for the western Alps, global  
82 warming and the increased frequency of extreme climate events such as heatwaves and droughts  
83 have raised awareness of the need to adapt, due to the combined effects of climate and changes in  
84 pastoral practices (Bonet et al., 2016). However, in many alpine zones, specific measures to  
85 manage pastures in the face of climate change are still not implemented, despite the implementation  
86 of agri-environmental and climate measures in the Common Agricultural Policy (EC, 2013). Since  
87 proper management is needed to ensure the environmental, social and economic sustainability of  
88 mountain permanent grasslands, a multi-disciplinary approach is a fundamental starting point,  
89 involving the co-responsibility of livestock farmers and local officers, as well as cooperation based  
90 on observation, modelling and intervention (Della-Vedova and Legeard, 2012).

91 This posture forms the basis of the design and implementation of this study started in 2017 in  
92 two representative areas of the western alpine territory: the *Écrins* (France) and *Gran Paradiso*  
93 (Italy) national parks (PNE and PNGP, respectively). In the pasturelands of the two parks, ground-  
94 based and remotely sensed observation systems, as well as model-based simulations were used to  
95 identify efficient management strategies able to support pastoral management and the sustainability  
96 of pastoral systems. Modelling adaptation strategies was supported by a participatory-based

97 process bringing together different local stakeholders in the two case study areas. The target of the  
98 modelling concerned the performance of pastoral systems and in particular the definition of  
99 production while minimising environmental impacts. Remote sensing supports such modelling by  
100 offering information on the spatial and temporal variation of important canopy state variables  
101 which would be difficult to obtain otherwise. The involvement of local pastoralists was the basis  
102 for the design and assessment of the analytical framework concerning the climate-change  
103 adaptation strategies.

104 In the context of these alpine pastures, the objectives of this study were: (1) to inform modelling  
105 via calibration with remotely sensed data; (2) to use the calibrated models to project climate-change  
106 impacts, and (3) to assess a set of adaptation options for pastoral management identified by  
107 stakeholders.

108

## 109 **2. Materials and Methods**

### 110 *2.1. Study areas*

111 In its wide-ranging perspective, the study considered three macro-types of pastoral vegetation  
112 (high, medium and low productivity) located at different altitudes (low, medium and high) in two  
113 national parks of the western Alps, on either side of the French-Italian border (Stendardi et al.,  
114 2022, Filippa et al., 2022) (Fig. 1).




115 Established in 1973, the *Parc National des Écrins* (PNE) covers an area of ~91800 ha  
116 (approximately in the range 44° 03'-45° 05' N and 06° 05'-06° 35' E) in the two French  
117 departments of Hautes-Alpes (region Provence-Alpes-Côte d'Azur) and Isère (region Auvergne-  
118 Rhône-Alpes). It includes *c.* 70,000 ha of summer pastureland (~30% of the park area), which is  
119 grazed by about 115,000 sheep (75% of the total stocking rate), 5,800 cows and >1,000 of goats  
120 and horses. Transhumance (which is declining across Europe) is still relevant in the study area,

121 with ~1/3 of the total sheep stocking rate in summer pasture being involved in transhumance (Brien,  
122 2018).

123 The *Parco Nazionale Gran Paradiso* (PNGP) is Italy's oldest national park (founded in 1922),  
124 established in the core of the former Piedmontese royal hunting reserve of the alpine ibex, a species  
125 of wild goat (*Capra ibex*) that lives in the mountains of the European Alps. It covers over 71,000  
126 ha approximately in the range 45° 25' -45° 45' N and 07° 00' -07° 30' E in the two Italian regions  
127 of Piedmont and Aosta Valley. Most of the territory (c. 60%) is used for non-agricultural purposes,  
128 a small part (c.11.5%) is covered by forests, while the areas of stable grasslands are constantly  
129 decreasing.

130 The surface of both parks is represented by mountainous environments, located from low valleys  
131 to very high mountains, with the highest peaks of 4,102 m a.s.l. (Barre des Écrins) and 4,061 m  
132 a.s.l. (Gran Paradiso Mountain) for the PNE and PNGP, respectively. The territories of the two  
133 protected areas are characterised by forests, from broadleaf in the lower parts to coniferous in the  
134 higher parts, and by mountain and alpine grasslands and pastures. The climate is generally alpine,  
135 but with different microclimatic conditions due to high variability in topographical features  
136 (elevation, aspect and slope). In addition, there are different lithological formations. All these  
137 complex and variable conditions produce a large typology of different plant communities  
138 characterised by a great richness of vegetation.

139





141 Figure 1. Location and details of the study areas: a) *Parc National des Écrins* (left) and *Parco*  
 142 *Nazionale Gran Paradiso* (right) with localization of the high, medium and low productivity  
 143 macro-types (i.e., green, yellow and red areas); b) Ikonos Panel Sharp (IPS) image from Google  
 144 Earth showing Italy and the position of the two Nation parks in the alpine chain; c) example of  
 145 grazing areas in the *Parc National des Écrins* and; d) in the *Parco Nazionale Gran Paradiso*.

146  
 147 The territories of the two parks lie within the areas of three vegetation macro-typologies (Table  
 148 1), which group the main plant communities that can be found in the subalpine and alpine pastures  
 149 of the French Southern Alps (Jouglet, 1999), the Vanoise and Aosta Valley (Bornard et al., 2007)  
 150 and Piedmont in Italy (Cavallero et al., 2007). These typologies have been harmonised in 13  
 151 categories of common pastures that were further grouped in three productivity macro-types  
 152 (Stendardi et al., 2022).

153  
 154 Table 1. Description of three pastoral macro-types (HP: high productivity; MP: medium  
 155 productivity; LP: low productivity) in the two study areas (PNE: *Parc National des Écrins*; PNGP:  
 156 *Parco Nazionale Gran Paradiso*).

Study area	Description	Unit	Pastoral macro-types		
			HP	MP	LP
PNE	Latitude	degree N	45.04	45.06	45.06
	Longitude	degree E	06.40	06.38	06.37
	Slope	rad	0.14	0.31	0.15
	Aspect	rad	3.06	1.95	2.32
	Elevation	m a.s.l.	2044	2539	2634
	Soil depth	m	0.70	0.65	0.55
	Clay	%	30.3	34.9	27.5
	Silt	%	37.6	40.3	61.1
	Sand	%	32.1	24.8	11.4
	Soil organic carbon	g 100 g <sup>-1</sup>	4.50	14.00	10.50
	Soil pH	-	5.70	5.05	4.75
	Bulk density	g cm <sup>-3</sup>	0.800	0.735	0.960
	Saturated soil water content	m <sup>3</sup> m <sup>-3</sup>	0.490	0.511	0.507
	Field capacity	m <sup>3</sup> m <sup>-3</sup>	0.312	0.345	0.330
	Wilting point	m <sup>3</sup> m <sup>-3</sup>	0.170	0.194	0.153

	Reference pasture type <sup>1</sup>	-	S6	S1	A9	
PNGP	Latitude	degree N	45.56	45.57	45.58	
	Longitude	degree E	07.12	07.19	07.29	
	Slope	rad	0.31	0.33	0.16	
	Aspect	rad	5.76	1.97	1.80	
	Elevation	m a.s.l.	2133	2336	2806	
	Soil depth	m	0.70	0.65	0.55	
	Clay	%	6.8	6.5	6.1	
	Silt	%	20.0	20.0	14.0	
	Sand	%	73.2	73.5	79.9	
	Soil organic carbon	g 100 g <sup>-1</sup>	1.88	2.24	1.90	
	Soil pH	-	5.5	4.9	5.3	
	Bulk density	g cm <sup>-3</sup>	1.48	1.48	1.51	
	Saturated soil water content	m <sup>3</sup> m <sup>-3</sup>	0.39	0.38	0.37	
	Field capacity	m <sup>3</sup> m <sup>-3</sup>	0.130	0.120	0.098	
	Wilting point	m <sup>3</sup> m <sup>-3</sup>	0.053	0.052	0.041	
		Reference pasture type <sup>1</sup>	-	S-II	SA-II	A-I

157 <sup>1</sup> A-I - Alpine intermediate: sparse vegetation on medium to moderate slopes, windy ridges and bumps in the alpine  
158 level (main species: *Carex curvula*, *Trifolium alpinum*, *Avenula versicolor*); SA-II - *Nardus* swards: on lowlands and  
159 slopes in the subalpine or alpine level, vegetation of medium height (0.2-0.3 m), not very dense, dominated by *Nardus*  
160 *stricta* (main species: *Nardus stricta*, *Carex sempervirens*, *Trifolium alpinum*, *Festuca rubra*); A-II – nival: sparse  
161 vegetation in snow combs and moderate slopes in alpine and nival environment (main species: *Alchemilla*  
162 *pentaphyllea*, *Salix herbacea*, *Carex foetida*, *Plantago alpina*); S-II - subalpine intermediate: vegetation in flatlands  
163 and low slopes of the subalpine level with medium-rich soil, 0.3 to 0.5 m high, dense grassy patches dominated by fine  
164 to medium-leaved *Gramineae* (main species: *Festuca rubra*, *Agrostis capillaris*, *Phleum alpinum*, *Alchemilla*  
165 *xanthochlora*); S-III - *Patzkea paniculata* swards: on medium sunny slopes in the subalpine level, vegetation very tall  
166 (over 0.5 m), very dense, dominated by *Gramineae* with long, thick leaves, especially *Patzkea paniculata* (main  
167 species: *Patzkea paniculata*, *Festuca rubra*, *Carex sempervirens*).  
168

## 169 2.2. Data collection

170 The Normalised Difference Vegetation Index (NDVI) is a standard way to determine vegetation  
171 cover and productivity. High NDVI values (approximately >0.7-0.8) indicate healthy vegetation,  
172 dense and productive canopies, while low NDVI values indicate land with little or no vegetation  
173 or stressed canopies. Satellite-derived NDVI data for the period 2018-2020 were retrieved for the  
174 two study areas by processing the Sentinel-2 imagery. The images (10-m spatial resolution,  
175 level2A) were atmospherically and topographically corrected with the Sen2Cor processor  
176 (<https://step.esa.int/main/snap-supported-plugins/sen2cor>). The images were filtered on a per-pixel  
177 basis with the scene classification (SCL) map, retaining only top quality, and cloud- and shadow-  
178 free pixels. The downloading and processing of the data were performed on Google Earth Engine  
179 (<https://earthengine.google.com>) with a dedicated Python (<https://www.python.org>) script

180 (Hufkens, 2017). Seasonal NDVI trajectories were used to retrieve growing season start and end  
181 based on a fixed threshold method (20% seasonal amplitude) similar to what proposed by Shen et  
182 al., (2015a). These dates were then compared to those retrieved from the snow cover seasonal  
183 pattern (table 1) and found to be in good agreement. Aboveground biomass (AGB) and leaf area  
184 index (LAI) were measured in both areas following standardised protocols (Filippa et al., 2015).  
185 An empirical model was fitted between AGB/LAI observations and the corresponding S2-derived  
186 NDVI, and the resulting equations (Supplementary material, section 1) were then used to convert  
187 S2-NDVI data in AGB and LAI data for the three productivity macro-types of each study area.

188

### 189 2.3. *Climate-change scenarios*

190 Simulated pastoral outputs were obtained by forcing impact models (section 2.5) with daily  
191 downscaled weather data, which were selected to map a broad range of climate outcomes for impact  
192 modelling (Wilcke and Barring, 2016). Supplementary material (section 2) describes the methods  
193 used in processing and post-processing the climate output used in the generation of climate  
194 scenarios.

195 Climate data from three Regional Climate Models (RCMs) from Med-CORDEX (Ruti et al.,  
196 2016) - CNRM-ALADIN ( $0.11^\circ \times 0.11^\circ$ ), ICTP-RGCM4 ( $0.44^\circ \times 0.44^\circ$ ), and CMCC-CCLM4  
197 ( $0.44^\circ \times 0.44^\circ$ ) for the reference period 1981-2010 (near past) and for two future time-slices 2011-  
198 2040 (near future) and 2041-2070 (mid future). For near past period ambient CO<sub>2</sub> concentration  
199 was fixed to 400 ppm. For future periods, Representative Concentration Pathways 4.5 and 8.5  
200 (RCP4.5, RCP4.5) were selected, with ambient CO<sub>2</sub> concentration at 450 (RCP4.5) and 470 ppm  
201 (RCP8.5) for near future and 540 and 670 ppm for mid future.

202 The delta-change approach was applied as a downscaling procedure, where the observed daily  
203 weather data available for each given site were modified using as forcing factors the outcomes

204 obtained from the RCM simulations. These were calculated as the mean absolute monthly  
205 differences between the RCM baseline (1981-2010) and the future RCM periods selected for  
206 simulations (2041-2070, 2071-2100) for minimum and maximum air temperatures and the  
207 percentage variation in monthly cumulated rainfall, wind speed and solar radiation. These  
208 differences were then added, month by month, to the observed daily meteorological data from PNE  
209 and PNGP to derive future weather data that were used to feed model simulations for future periods.  
210 The three daily datasets deriving from RCMs downscaling were finally merged into a single dataset  
211 reproducing the mean change in climate conditions for each study area in RCP4.5 and 8.5 for 2031-  
212 2040, 2041-2070 and 2071-2100 time-slices.

213

#### 214 2.4. *Participatory approach*

215 To understand the impact of climatic events and changes in grazing practices, and to preserve  
216 (or restore) the sustainable management of these areas, the "Sentinel Alpine Pastures" programme  
217 focuses on how to adapt to different phenomena as part of a long-term approach to the complex  
218 dynamics of climate change, to anticipate adaptive strategies (Dobremez et al., 2014). These  
219 sources of information thus represent a unique opportunity to environmentally characterise these  
220 pastoral areas by using advanced techniques such as remote sensing and process-based simulation  
221 modelling. As a basis for the design and assessment of the analytical framework, a participatory  
222 process was conducted since 2018 with groups of *c.* 100 local stakeholders in each park including  
223 farmers, technicians, representatives of the two parks and officials from local institutions. The  
224 participatory process involved meetings, interviews and informal discussions that took place in  
225 parallel with data collection and territorial analysis (Targetti et al., 2019). Participation addressed  
226 three main topics: i) current pastoral practices, related barriers and incentives, and key drivers of  
227 socio-economic change; ii) effective adaptation measures already implemented in the western

228 Alps; and iii) which measures should be prioritised (Piccot et al., 2022). In this study, we assessed  
229 the effect of prioritised adaptation options from a modelling perspective as it emerged from the  
230 participatory approach, recognising the limited set of modelling assumptions contained in the  
231 adaptation requests, which represent a fraction of plausible adaptations and a step towards  
232 transformative changes (Holman et al., 2019).

233

### 234 2.5. *Grassland modelling*

235 Process-based models are important tools in agricultural and environmental research to  
236 extrapolate local observations over time and space, and to assess the impact of climate and  
237 agricultural practices on the soil-plant-atmosphere continuum through plant-soil feedback effects.  
238 These widely tested models are also recognised as effective tools for studying the magnitude and  
239 spatial-temporal patterns of C-N (carbon-nitrogen) fluxes, playing a prominent role in testing the  
240 effect of specific changes in management, plant properties or environmental factors, and in  
241 designing policies specific to the soil, climate and agricultural conditions of a location or region.  
242 However, the results from different models often differ, presenting a range of possible impacts and  
243 adaptation responses (Brilli et al., 2017), which are influenced by the models' users' knowledge  
244 and expertise, and their understanding of the variables determined in the target agroecosystems  
245 (Albanito et al., 2022).

246 Here, the soil-vegetation generic model DayCent (Parton et al., 1994, 1998) and the grassland-  
247 specific model PaSim (Riedo et al., 1998) were chosen to simulate alpine pastures. Both provide a  
248 mechanistic view of the multiple processes and interactions occurring in grassland systems and are  
249 able to simulate grassland productivity and C and N fluxes under alternative management options.  
250 DayCent is the daily time-step adaptation of the biogeochemical model CENTURY (Parton et al.,  
251 1994), which simulates plant growth, soil C dynamics, N leaching, gaseous emissions (e.g. nitrous

252 oxide) and C fluxes (e.g. net ecosystem exchange) in a variety of managed ecosystems. PaSim is a  
253 grassland-specific ecosystem model consisting of detailed sub-models for vegetation, animals,  
254 microclimate, soil biology, soil physics and management to simulate grassland productivity and C-  
255 N fluxes.

256

## 257 2.6. *Simulation design*

258 The modelling work was carried out in three suites of simulations: suite 1 with observational  
259 data (model calibration), suite 2 with projected scenarios of climate change (impact projections),  
260 and suite 3 with altered management under projected scenarios of climate change (adaptation  
261 assessment).

262 Model calibration (suite 1) was carried out over the years 2018 to 2020 in the two parks, setting  
263 management practices (grazing intensity and periods) as defined in Table 2 (one or two short  
264 periods with short-term, intensive management), on a set of parameters (Table S1 and Table S2) to  
265 which model sensitivity was determined in previous studies for both DayCent (e.g. Fitton et al.,  
266 2014; Nécipalová et al., 2015) and PaSim (e.g. Ben Touhami et al., 2013; Ma et al., 2015; Pulina et  
267 al., 2018; Sándor et al., 2018). The agreement between simulated and observed dry matter (DM)  
268 was assessed by inspection of time-series graphs (fluctuations of output variables over time), and  
269 numerically, through two commonly used performance metrics of model evaluation (Richter et al.,  
270 2012): root mean square error (best,  $0 \leq \text{RMSE} < +\infty$  g DM m<sup>-2</sup>, worst) and coefficient of  
271 determination (worst,  $0 \leq R^2 \leq 1$ , best).

272 Table 2. Management of three pastoral macro-types (HP: high productivity; MP: medium productivity; LP: low productivity) in the two  
 273 study areas (PNE: *Parc National des Écrins*; PNGP: *Parco Nazionale Gran Paradiso*). Grazing 1 and Grazing 2 refer to the first and  
 274 second (if present) grazing periods expressed as days of the years, respectively, over the investigated macro-types. Livestock Standard  
 275 Unit (LSU) refers to a dairy cow producing 3000 kg of milk per year, without additional concentrated feed (EC, 2008).

Site	Pasture macro-type	Grazing 1 <sup>st</sup>						Grazing 2 <sup>th</sup>					
		Period (days of year)			Stocking density (LSU ha <sup>-1</sup> d <sup>-1</sup> )			Period (days of year)			Stocking density (LSU ha <sup>-1</sup> d <sup>-1</sup> )		
		2018	2019	2020	2018	2019	2020	2018	2019	2020	2018	2019	2020
PNE	HP	196-197	197-198	191	120	113	126	287-288	272-273	262	43	37	76
	MP	213-214	213-214	214-215	51	49	62	-	-	-	-	-	-
	LP	217	220	217	12	10	9	-	-	-	-	-	-
PNGP	HP	194-195	198	196-197	104	98	84	261	264	264	102	106	118
	MP	229-230	230-231	229-230	79	75	57	-	-	-	-	-	-
	LP	217-218	202	222-223	30	14	20	-	-	-	-	-	-

276

277 With suite 2, we assessed the projected response of DayCent and PaSim to climate-change forcing  
278 options described in section 2.3. Impacts of climate change were calculated on the changes in a set  
279 of climate and ecosystem variables related to biomass production and C-N fluxes (Table 3).



280 Table 3. Climate-change impact metrics.

Type	Output	Acronym	Unit	Description
Date	Snow cover start	SCs	day of year (doy)	First of 10 consecutive days of the year with snow cover $\geq 5$ cm
	Snow cover end	SCe		First of 10 consecutive days of the year with snow cover $\leq 5$ cm
	Growing seasons start	GSs		First day of the year with aboveground biomass (SCe + 1 day)
	Growing seasons end	GSe		Last day of the year with aboveground biomass (SCs - 1 day)
	Biomass peak date (period 1)	BP1a		Day of the year with the highest value of aboveground biomass before the first grazing period
	Biomass peak date (period 2, HP)	BP2a		Day of the year with the highest value of aboveground biomass after the first grazing period and before the second grazing period
Count	Snow cover length	SC	days	Number of days between SCs and SCe
	Growing season length	GS		Number of days between the GSs and GSe
Amount	Biomass peak (period 1)	BP1b	kg DM m <sup>-2</sup>	Aboveground biomass value at the first peak date
	Biomass peak (period 2, HP)	BP2b		Aboveground biomass value at the second peak date
	Above ground biomass	AGB	kg DM m <sup>-2</sup> yr <sup>-1</sup>	Annual mean aboveground biomass
	Net ecosystem exchange	NEE	kg C m <sup>-2</sup> yr <sup>-1</sup>	C-N fluxes  (they include emissions from ecosystem respiration (RECO = plant + soil + animal respiration), as well as estimates of the plant production of organic compounds from atmospheric CO <sub>2</sub> (GPP) and other system variables: NEE = RECO - GPP, NPP = GPP - plant respiration, enteric emissions of CH <sub>4</sub> from grazing animals and N <sub>2</sub> O emissions from the N cycle)
	Net primary production	NPP		
	Ecosystem respiration	RECO		
	Gross primary production	GPP		
	Methane	CH <sub>4</sub>		
	Nitrogen dioxide	N <sub>2</sub> O	kg N m <sup>-2</sup> yr <sup>-1</sup>	
Soil water content	SWC	m <sup>3</sup> m <sup>-3</sup>	Soil water content	

281

282 With the adaptation assessment (suite 3), we show the simulated outputs using the two grassland  
283 models fed with the following adaptation practices, defined during the participatory process  
284 (section 2.4) combined with climate-change forcing: the stocking rate in the pasture was increased  
285 or decreased by 20% (LD-20% and LD+20%, respectively), and the grazing period was advanced  
286 by 14 days (GDadv).

287 Simulation results are presented separately per study area, comparing DayCent and PaSim  
288 outputs with satellite-derived AGB data (suite 1). Time-series graphs are presented to illustrate the  
289 dynamics of selected variables (AGB, SWC, C fluxes and CH<sub>4</sub> and N<sub>2</sub>O emissions) for suites 2 and  
290 3, as well as two-dimensional colour data visualisations (heatmap graphs).

291

### 292 **3. Results**


293 For greater clarity in the presentation of results and discussion, we present in detail only the  
294 results obtained in the high productivity macro-type for which a full modelling analysis is available.  
295 We also briefly present the results obtained in the medium and low productivity macro-types,  
296 which are fully provided in the Supplementary material.

297

#### 298 *3.1. Climate analysis*

299 The monthly distribution of air temperatures in the two study areas (Fig. 2), averaged from the  
300 outputs of the ICTP-REGCM4, CMCC-CCLM4 and CNRM-ALADIN climate models, showed an  
301 overall increase in temperature towards the far future, similar for both parks, with a distinct  
302 seasonal trend, with the highest increases in summer (+4 °C at PNE and +3.7 °C at PNGP under  
303 the warmest scenario) and the lowest in autumn-winter (+2.5 °C at PNE and +2.3 °C at PNGP  
304 under the warmest scenario). Analysis of simulated monthly rainfall data (Fig. 2) showed increases  
305 in autumn-winter (November-February) relative to the baseline in both scenarios and sites (PNE:

306 +3.3% and +9.9%; PNGP: +5.4 % and +9.5 %, for RCP4.5 and RCP8.5, respectively), while  
 307 spring-summer exhibited a strong decrease in rainfall, more pronounced in the PNE (-11.7% and -  
 308 18.8%, for RCP4.5 and RCP8.5, respectively) than in the PNGP (-10 % in both scenarios). In both  
 309 parks, no clear trend was observed, nor was a clear pattern evident when analysing the differences  
 310 between time-slices, as there was no trend of increasing/decreasing monthly precipitation in the  
 311 progression from the near to the far future.



312  
 313 Figure 2. Absolute change (°C) in monthly mean air temperature (top graphs) and relative change  
 314 (%) of monthly cumulated rainfall (bottom graphs) generated in the two study areas with the RCM  
 315 ensemble (ICTP-REGCM4, CMCC-CCLM4 and CNRM-ALADIN) for two climate scenarios  
 316 (RCP4.5, RCP8.5) and two future periods - 2011-2040 and 2041-2070 - over the baseline period  
 317 1981-2010.

318

319 3.2. *Suite 1 of simulations: model evaluation against observed data*

320 The resulting sets of parameter values allowed the outputs of the two impact models to be  
321 compared for each study area. Plant parameters served to accommodate changes in the sward  
322 structures driven by local environmental conditions and management. Although no formal  
323 sensitivity analysis was conducted for the model parameters, the calibration applied separately to  
324 each study area allowed us to explore the variability of parameter values between the two parks.  
325 An indication from the calibration work is that, for both models, the parameter values can be  
326 considerably different across alternative conditions (Table S1 and Table S2). For instance, the  
327 different vegetation patterns in the two parks are reflected in the PaSim parameter “maximum  
328 specific leaf area”, whose lower values tend to be associated with the PNGP (e.g. for the high-  
329 productivity pastoral vegetation macro-type, the value decreased from  $\sim 37 \text{ m}^2 \text{ kg}^{-1}$  in the PNE to  
330  $\sim 22 \text{ m}^2 \text{ kg}^{-1}$  in the PNGP). Photosynthetic rates estimated with PaSim (Table S1) were lower in the  
331 PNGP during the reproductive stage ( $pmco2rep \sim 25 \mu\text{mol C m}^{-2} \text{ s}^{-1}$  against  $\sim 32 \mu\text{mol C m}^{-2} \text{ s}^{-1}$  in  
332 the PNE) and higher during the vegetative stage ( $pmco2veg \sim 16 \mu\text{mol C m}^{-2} \text{ s}^{-1}$  against  $\sim 13 \mu\text{mol}$   
333  $\text{C m}^{-2} \text{ s}^{-1}$  in the PNE). With DayCent, air temperature thresholds (optimal and maximum), the  
334 number of soil layers influencing water and nutrient availability, and the allocation of C to different  
335 plant organs influenced plant growth and C fluxes. Specifically, for the high-productivity pastoral  
336 vegetation macro-type, the coefficient for calculating potential monthly aboveground biomass  
337 production as a function of solar radiation outside the atmosphere lowers from 4.1 in the PNGP to  
338  $\sim 1.0 \text{ m}^2 \text{ kg}^{-1}$  in the PNE, while the thresholds for optimal air temperatures were slightly higher in  
339 the PNGP than in the PNE (Table S2).

340 Standing biomass simulations (Fig. S2, Table S3) indicate that estimates substantially reflect  
341 patterns of vegetation dynamics ( $R^2 > 0.50$ ) although some departures from observed data are noted.  
342 RMSE values ( $> 70 \text{ g DM m}^{-2}$ ) are comparable with results from previous modelling studies (e.g.

343 Sándor et al., 2018), with simulations for grasslands being generally less accurate compared to  
344 arable crops (e.g. Kollas et al., 2015). We also note that the great deal of fundamental research  
345 incorporated into the most mechanistic PaSim model has not always improved the results.

346

### 347 3.3. *Suite 2 and 3 of simulations: impacts of future scenarios and adaptation strategies*


348 For both parks, we assessed the sensitivity of the two grassland models to (suite 2) climate  
349 change (RCP4.5 and RCP8.5 for the near and far future) with business-as-usual (BaU) management  
350 and to (suite 3) management scenarios (GDadv and LD±20%). Multi-year mean responses for  
351 selected production (AGB), biophysical (SWC) and biogeochemical (C-N fluxes) outputs are  
352 presented below.

353

### 354 3.4. *Growing season*

355 Under the climate-change scenarios, with both grassland models, the estimated length of the  
356 snow season decreases in both areas due to earlier spring snowmelt and later autumn/winter  
357 snowpack accumulation. This condition leads to an earlier onset and later end of the growing season  
358 (GS) in both parks, especially in the far future (i.e. 2041-2070) (Fig. 3). Specifically, using  
359 DayCent, the start of the growing season (GSs) was on average 11 and 28 days earlier in the PNE,  
360 and 12 and 39 days earlier in the PNGP, for the 2011-2040 and 2041-2070. The end of the growing  
361 season (GSe) was delayed on average 8 and 17 days in the PNE, and 17 days in the PNGP for 2041-  
362 2070. In contrast, no changes in GSe were observed in PNGP for the period 2011-2040.

363 Using PaSim, GSs was advanced by 14 and 31 days on average in the PNE, and by 7 and 19  
364 days in the PNGP for the periods 2011-2040 and 2041-2070. GSe was delayed by 5 and 23 days  
365 on average for the periods 2011-2040 and 2041-2070 in the PNE, and by 36 days in the PNGP for  
366 both time-slices.



368

369 Figure 3. Estimated durations (20-year mean values) of snow-cover periods (SC, grey bars) and  
 370 vegetation growing seasons (green bars) with two grassland models for baseline and climate-  
 371 change scenarios under business-as-usual management in both parks for the high productivity (HP)  
 372 macro-type. The annual pattern was reported at daily time-step (DOY: day of the year).

373


374 The MP and LP macro-types showed similar growing season patterns to those observed in the  
 375 HP macro-type, with GSs advanced and GSe delayed towards the end of the century, with the  
 376 largest impacts using RCP8.5. For all three macro-types, DayCent reported a mean GS extension  
 377 ranging between 15 and 40 days in the PNE, and between 12 to 45 days in the PNGP for the periods  
 378 2011-2040 and 2041-2070, respectively. Using PaSim, the increase in GS ranged between 17 and  
 379 44 days in the PNE, and between 23 and 35 days in the PNGP for the periods 2011-2040 and 2041-  
 380 2070, respectively (Fig. S3 and Fig. S4). Overall, both models suggested a longer growing season  
 381 of 2 to 5 weeks when approaching the warmest scenarios.

382

383 3.5. Soil water content (0.30 m topsoil)

384 Under the climate-change scenarios, both models indicated an earlier decline in SWC, near or  
385 below the permanent wilting point (Table 1), especially during the warm season in both parks (Fig.  
386 4). PaSim showed less pronounced oscillations in SWC ( $\sim 0.30\text{-}0.40\text{ m}^3\text{ m}^{-3}$  in the PNE and  $\sim 0.15\text{-}$   
387  $0.25\text{ m}^3\text{ m}^{-3}$  in the PNGP), while DayCent interpreted the increased water supply projected by  
388 climate modelling in winter (Fig. 2) to amplify seasonal differences (i.e. an excess SWC in winter  
389 followed by a deficit in summer), with  $\sim 0.15\text{-}0.60\text{ m}^3\text{ m}^{-3}$  in the PNE and  $\sim 0.05\text{-}0.35\text{ m}^3\text{ m}^{-3}$  in the  
390 PNGP (i.e. even below the permanent wilting point). Despite the differences between the two  
391 models, for both parks the simulated patterns suggest that with drier summer conditions, grassland  
392 growth may be limited by water in summer (Fig. 4).

393



394

395 Figure 4. Simulated annual pattern (20-year mean values) of 0.30-m soil water content (SWC) with  
396 two grassland models (DayCent, PaSim), for baseline and climate-change scenarios under

397 business-as-usual management in both parks for the high productivity (HP) macro-type. The annual  
398 pattern was reported at daily time-step (DOY: day of the year).

399  
400 The MP and LP macro-types showed SWC patterns similar to those observed for the HP macro-  
401 type, with a reduction in SWC when approaching warmer scenarios and less pronounced SWC  
402 oscillations in PaSim compared to DayCent (Fig. S5 and Fig. S6). In the MP macro-type, the SWC  
403 simulated by DayCent ranged over  $\sim 0.20\text{-}0.65 \text{ m}^3 \text{ m}^{-3}$  in the PNE and  $\sim 0.05\text{-}0.40 \text{ m}^3 \text{ m}^{-3}$  in the  
404 PNGP, whereas with PaSim, the SWC was in the range  $\sim 0.30\text{-}0.45 \text{ m}^3 \text{ m}^{-3}$  in the PNE and  $\sim 0.12\text{-}$   
405  $0.24 \text{ m}^3 \text{ m}^{-3}$  in the PNGP (Fig. S5). In the LP macro-type, the SWC simulated by DayCent ranged  
406 from  $\sim 0.20\text{-}0.65 \text{ m}^3 \text{ m}^{-3}$  in the PNE and  $\sim 0.05\text{-}0.40 \text{ m}^3 \text{ m}^{-3}$  in the PNGP, while with PaSim, the  
407 SWC was in the range  $\sim 0.32\text{-}0.48 \text{ m}^3 \text{ m}^{-3}$  in the PNE and  $\sim 0.12\text{-}0.22 \text{ m}^3 \text{ m}^{-3}$  in the PNGP (Fig. S6).


408

### 409 3.6. *Aboveground biomass*

410 Figure 5 shows the yearly average AGB production patterns under baseline management in both  
411 parks for the HP macro-type as obtained with the two grassland models, while the yearly average  
412 AGB patterns obtained with all alternative management options can be found in the Supplementary  
413 material (Figs. S7-S10). The main differences in AGB patterns among alternative management and  
414 climate scenarios were assessed based on changes in peak biomass dates (BP1a and BP2a) and  
415 corresponding AGB values (BP1b and BP2b), which strongly influence stakeholders' and farmers'  
416 decisions in choosing the most suitable periods for grazing.

417





418  
 419 Figure 5. Simulated annual pattern (20-year mean values) of aboveground biomass (AGB) with  
 420 two grassland models, for baseline and climate-change scenarios under business-as-usual  
 421 management in both parks for the high productivity (HP) macro-type. The annual pattern was  
 422 reported at daily time-step (DOY: day of the year).

423  
 424 Under the baseline climate scenarios, DayCent reported the first biomass peak (BP1a) on day  
 425 189 ( $\pm 9$  standard deviation) and 190 ( $\pm 8$  standard deviation) for the PNE and PNGP, respectively.  
 426 Under future climate scenarios, the model indicated an advance of BP1a of 7-10 days for the PNE  
 427 and 3-7 days for the PNGP (Table S4). In contrast, the biomass peak simulated by PaSim was  
 428 mainly driven by the effect of grazing, showing only a slight advance under the future scenarios  
 429 (i.e. 2-3 days) for both PNE ( $194 \pm 4$ ) and PNGP ( $196 \pm 5$ , Table S5).

430 For the second biomass peak (BP2a), DayCent indicated that biomass peaks were at day 267  
 431 ( $\pm 14$  standard deviation) in the PNE and day 244 ( $\pm 13$  standard deviations) in the PNGP under the  
 432 baseline scenarios, while future scenarios suggested advanced biomass peaks of 3 to 15 days in the

433 PNE and contrasting patterns (from -3 to +2 days) in the PNGP (Table S4). PaSim indicated that  
434 BP2a was on day 262 ( $\pm 7$  standard deviation) in the PNE and on day 260 ( $\pm 2$  standard deviation)  
435 in the PNGP under baseline scenarios, while the future scenarios indicated no or slight delay (1-5  
436 days) in the PNGP and PNE, respectively (Table S5).

437 In the baseline scenarios, the biomass production of the first peak (BP1b) is similar with both  
438 models in the PNE ( $\sim 0.52 \pm 0.06$  kg DM m<sup>-2</sup>), while in the PNGP it is  $\sim 38\%$  lower with PaSim  
439 compared to DayCent ( $\sim 0.61 \pm 0.17$  kg DM m<sup>-2</sup>). For the second peak (BP2b), the biomass value  
440 provided by DayCent ( $0.44 \pm 0.06$  kg DM m<sup>-2</sup>) was close to that provided by PaSim ( $0.43 \pm 0.08$  kg  
441 DM m<sup>-2</sup>) in the PNE, while at the PNGP the biomass simulated by DayCent ( $0.52 \pm 0.14$  kg DM m<sup>-2</sup>)  
442 was higher compared to that provided by PaSim ( $0.41 \pm 0.06$  kg DM m<sup>-2</sup>). Future patterns for  
443 BP2b partly mirror those of BP1b, with PaSim providing an increase in biomass production of  
444  $\sim 18\%$  in the PNE and  $\sim 41\%$  in the PNGP as the warmer scenarios are approached, while DayCent  
445 reported a decrease in biomass production of  $\sim 20\%$  in both study areas (Table S4 and Table S5).  
446 These results mainly reflect calibration against observational patterns (Fig. S2), with the PaSim  
447 production profile indicating faster plant growth in spring, with a distinct peak biomass, and rapid  
448 regrowth in summer. This behaviour is much more evident in the climate-change scenarios,  
449 resulting in differences in AGB that are about 38-45% higher at the peak with PaSim than with  
450 DayCent (Fig. 5), likely due to the absence of sensible water deficits simulated by PaSim (Fig. 4).

451 For the MP and LP macro-types (Tables S6-S9), the biomass peaks (BP1b and BP2b) partly  
452 reflect the trends found in the HP macro-type. Specifically, while PaSim reported an increase in  
453 peak biomass value of 50-100% with warmer scenarios in all macro-types for both parks, DayCent  
454 indicated a decrease of 3-20% with the sole exception of the LP macro-type in the PNE, where  
455 biomass production increased of  $\sim 25\%$ . For the impact of adaptation strategies, the value of peak  
456 biomass obtained with alternative management practices (i.e. BaU + adaptation management


457 options) was compared with the peak biomass of business-as-usual (BaU) management under  
458 projected scenarios. To simplify the reading, only the first biomass peak of the HP macro-type in  
459 both parks is reported here (Fig. 6), while the dynamics of the second peak (Fig. S11) and those of  
460 the MP and LP macro-types are reported in the Supplementary material (Tables S6, S7, S8 and  
461 S9).

462 Using DayCent, in the PNE under RCP4.5 (blue), on average, the highest AGB values at the  
463 first biomass peak compared to BaU ( $0.52 \pm 0.06$  kg DM m<sup>-2</sup>) was obtained with LD+20% at both  
464 current (+18.3%) and advanced (+13.5%) dates (Fig. 6). Only a slight increase was observed with  
465 the other strategies (+1 to +7.7%). Under RCP8.5 (orange), BP1b shows a similar pattern to that  
466 observed under RCP4.5, with higher values occurring with the adoption of the LD+20% strategy  
467 at both current (+16.3%) and advanced (+13.4%) dates, and a slight mean increase using other  
468 strategies (+3.8 to +6.7%). In the PNGP, under RCP4.5, a decrease in BP1b values compared to  
469 current BaU ( $0.61 \pm 0.17$  kg DM m<sup>-2</sup>) was observed with all alternative strategies, with the smallest  
470 decrease when adopting LD+20% (-5.4%) and the highest when using GDadv\_LD-20% (-18%).  
471 Under RCP8.5 (Fig. 6b), BP1b showed a similar pattern and magnitude to those observed under  
472 RCP4.5, with the largest decrease when adopting Gdadv\_LD-20% (-17.2%) and the lowest when  
473 using LD+20% (-4.9%).

474 Using PaSim, all management options showed an increase in peak biomass under all climate  
475 scenarios and time-slices. Specifically, in the PNE under RCP4.5, higher BP1b values compared  
476 to BaU ( $0.50 \pm 0.17$  kg DM m<sup>-2</sup>) were observed, on average, when the same grazing dates were  
477 maintained with all management options (+43%) while a smaller increase was observed when  
478 grazing dates were advanced (+11.7%). Under RCP8.5, BP1b shows a similar pattern to that  
479 observed under RCP4.5, with higher values occurring when both current (+46%) and advanced  
480 (+16.7%) grazing dates are adopted. In the PNGP, under RCP4.5, BP1b values compared to BaU

481 (0.37±0.11 kg DM m<sup>-2</sup>) were observed, on average, both maintaining the same grazing dates with  
482 all management options (+47.3%) and advancing grazing dates (+23.9%). Under RCP8.5, BP1b  
483 showed the same pattern as under RCP4.5, with higher values at both current (+53.6%) and  
484 advanced (+32.4%) grazing dates. Overall, DayCent showed less variability in peak biomass  
485 production in the PNE than in the PNGP, with increasing variability as we approach the far future  
486 (2041-2070) with the warmest scenario (i.e. RCP8.5) in both parks. In contrast, PaSim indicated  
487 greater variability in peak biomass production in the PNE than in the PNGP, with decreasing  
488 variability towards the far future with the warmest scenario in the PNE and contrasting patterns in  
489 the PNGP.

490



491  
 492 Figure 6. Changes in the first (BP1b) peak aboveground biomass (kg DM m<sup>-2</sup>) between business-  
 493 as-usual management (BaU) under baseline climate (black histogram) and all alternative  
 494 management options under RCP4.5 (cyan and blue histograms) and RCP8.5 (clear and dark orange  
 495 histograms) for high productivity pasture (HP) in both parks as provided by DayCent and PaSim.  
 496 Vertical bars are standard deviations.

497

498 For the MP and LP macro-types, PaSim suggested a generalised increase in biomass production  
 499 that was particularly large (>50%) in the PNE and smaller in the PNGP across all macro-types. In  
 500 contrast, DayCent reported no decline or a decrease (-6%) in production for the MP macro-type in  
 501 both parks, regardless of advanced grazing management, while for the LP macro-type it showed  
 502 contrasting patterns. Specifically, a slight decrease in productivity (-4%) was observed in the PNGP  
 503 when approaching the warmest scenario, irrespective of management, while a 10-20% increase in  
 504 productivity was found in the PNE when approaching the warmest scenario at the current grazing  
 505 date and with different livestock densities (i.e. BaU, LD-20% and LD+20%).

506

### 507 3.7. Carbon-nitrogen fluxes

508 Under current climate and management conditions, PaSim shows limited non-CO<sub>2</sub> emissions in  
 509 both parks, i.e. 1.9 and 1.6 g C m<sup>-2</sup> yr<sup>-1</sup> for CH<sub>4</sub> and 1 and 3 g N m<sup>-2</sup> yr<sup>-1</sup> for N<sub>2</sub>O emissions, while  
 510 the C exchanges (NEE) vary from a limited sink in the PNE (-41 g C m<sup>-2</sup> yr<sup>-1</sup>) to a limited source  
 511 in the PNGP (+96 g C m<sup>-2</sup> yr<sup>-1</sup>). DayCent represents a higher sinking pattern (-350 and -308 g C m<sup>-2</sup>  
 512 yr<sup>-1</sup>) and lower CH<sub>4</sub> emissions (2.5E-04 and 1.2E-04 g C m<sup>-2</sup> yr<sup>-1</sup>) in both parks, while N<sub>2</sub>O  
 513 emissions (0.5 and 3.8 g N m<sup>-2</sup> yr<sup>-1</sup>) are in agreement with PaSim (Table 4).

514

515 Table 4. C-N emissions (NEE: net ecosystem CO<sub>2</sub> exchange; CH<sub>4</sub>: methane; N<sub>2</sub>O: nitrous oxide)  
 516 from the two study areas (baseline climate), estimated (20-year mean ± standard deviation) using  
 517 two grassland models. The estimated components of the C budget (GPP: gross primary production;  
 518 NPP: net primary production; RECO: ecosystem respiration) can be found in Supplementary  
 519 material (Table S10).

Site	Model	NEE	CH <sub>4</sub>	N <sub>2</sub> O
		g C m <sup>-2</sup> yr <sup>-1</sup>		g N m <sup>-2</sup> yr <sup>-1</sup>

PNE	DayCent	-350±14	2.5E-04±~0.0	0.5±0.1
	PaSim	-41±12	1.9±0.9	1.0±0.7
PNGP	DayCent	-308±19	1.2E-04±~0.0	3.8±1.3
	PaSim	96±11	1.6±1.0	3.0±0.9

520

521 The absolute values of C-N fluxes (Fig. S12) indicate that both models agree in representing the  
522 magnitude of these fluxes, and the differences are explained by the inherent features of the two  
523 model structures (i.e. animal respiration, enteric fermentation). Heatmaps of the % differences  
524 between current conditions (i.e. baseline climate and BaU management) and combinations of  
525 alternative climate and management scenarios allow the impact of altered climate and management  
526 changes on gas emissions in the two parks to be assessed (Fig. 7).

527 For NEE, in particular, the PaSim heatmaps show overall trends towards C uptake (more  
528 negative NEE values) in both parks (red colour) by moving towards extreme climate conditions  
529 (i.e. RCP8.5 and time-frame 2041-2070), reducing livestock density and advancing grazing dates,  
530 thus reflecting the baseline AGB pattern (Fig. 5) and the inclusion in the model of an animal  
531 component explicitly representing animal respiration and enteric fermentation (Graux et al., 2011).

532 In contrast, DayCent reports an increase in C sourcing (more positive NEE values) of up to 30%  
533 in both parks (green colour) when extreme climate conditions are approached, which is higher  
534 when livestock density is reduced. An increase in C uptake of up to 30% was observed at both  
535 current grazing date and advanced grazing date when the livestock density is increased.

536 As for CH<sub>4</sub> emissions, the PaSim heatmap indicates that emissions are higher (~>20%) as  
537 livestock density increases. While this pattern is clearly observed in the PNE, the results in the  
538 PNGP are more contrasted, as the earlier grazing date also leads to increased CH<sub>4</sub> emissions, even  
539 when livestock density is reduced. Projected climate conditions do not appear to influence the  
540 pattern of emissions, which are mainly driven by management. In contrast, the CH<sub>4</sub> emissions

541 estimated by DayCent are conditional on climatic conditions, with the highest emission values (up  
542 to ~30%) occurring towards the end of the century (i.e. in the period 2041-2070).


543 Finally, the N<sub>2</sub>O emissions estimated by PaSim were mainly driven by the type of management  
544 adopted in the two parks, where an increase in livestock density leads to higher emissions (up to  
545 ~40%), while a decrease in livestock density reduces emissions to ~50%. Unlike PaSim, DayCent  
546 shows contrasting patterns between the two parks. Specifically, N<sub>2</sub>O emissions in the PNE are  
547 mainly driven by management, where increasing livestock density leads to increased emissions (up  
548 to ~30%), while in the PNGP, N<sub>2</sub>O emissions are mainly driven by the climate scenarios, with the  
549 highest emissions (up to ~40%) for the period 2041-2070 under both RCPs (4.5 and 8.5).

550 Under the baseline scenario, PaSim-simulated NEE for the LP macro-type showed contrasting  
551 patterns. Simulated NEE in the PNE ( $195\pm 193$  g C m<sup>-2</sup> yr<sup>-1</sup>) decreased as the warmest scenarios  
552 were approached ( $107\pm 181$  g C m<sup>-2</sup> yr<sup>-1</sup>), while simulated NEE in the PNGP ( $151\pm 72$  g C m<sup>-2</sup> yr<sup>-1</sup>)  
553 increased as the warmest scenarios were approached ( $163\pm 97$  g C m<sup>-2</sup> yr<sup>-1</sup>), making both parks  
554 sources of C (Fig. S14 and Fig. S15). For the MP macro-types, NEE decreased in both parks as  
555 warming scenarios approached, with the PNE still being a source of C ( $448\pm 388$  g C m<sup>-2</sup> yr<sup>-1</sup>) while  
556 the PNGP turned into a sink of C ( $-91\pm 81$  g C m<sup>-2</sup> yr<sup>-1</sup>) (Fig. S14 and Fig. S15). In contrast, under  
557 the baseline climate scenario, DayCent- simulated NEE in both MP ( $-126\pm 36$  and  $-163\pm 135$  g C  
558 m<sup>-2</sup> yr<sup>-1</sup>) and LP ( $-9\pm 19$  and  $-66\pm 41$  g C m<sup>-2</sup> yr<sup>-1</sup>) macro-types showed negative values in both parks.  
559 Under the warmest scenarios, NEE tended to decrease for all macro-types in both parks, with the  
560 sole exception of the LP macro-type in the PNE, where it showed a significant increase (+90%) in  
561 C uptake (Fig. S13 and Fig. S14).

562 The patterns of simulated CH<sub>4</sub> and N<sub>2</sub>O emissions for the LP and MP macro-types matched  
563 those reported for the HP macro-type, where the estimates provided by DayCent were mainly



564 driven by climatic conditions whilst those of PaSim were mainly related to the different  
565 management types (Fig. S13 and Fig. S14).



566

567 Figure 7. Heatmap visualisation of the relative differences (%) between the three main greenhouse gas emissions (NEE: net ecosystem  
 568 exchange; CH<sub>4</sub>: methane; N<sub>2</sub>O: nitrous oxide), estimated using two grassland models (DayCent, PaSim), for alternative management  
 569 and climate-change scenarios compared to the current climate and management in the *Parc National des Écrins* (PNE) and *Parco*  
 570 *Nazionale Gran Paradiso* (PNGP). Absolute values are given in the supplementary material (Fig. S10).

## 571 **4. Discussion**

### 572 *4.1. Uncertainty in climate-change impact assessments*

573 The issue of uncertainty in model outputs remains a real challenge for the implementation of a  
574 modelling framework in decision-making or information processes. Uncertainty can arise from  
575 several sources (model structure, parameterisation, input data, initialisation), and the way it  
576 manifests itself in model estimates can be difficult to determine and requires interpretation by  
577 stakeholders. In this study on climate-change and impact projections in alpine pasturelands, there  
578 are different levels of uncertainty on multiple elements (e.g. biophysical, socio-economic). The  
579 lack of certainty about the future is primarily related to prospective views at various scales: global  
580 (socio-economic scenarios), regional (land and soil use) and field (pastoral systems and  
581 management). In addition, the chaotic character of the climate system, with its interannual  
582 variability, limits the reliability of climate projections. For instance, the CNRM-ALADIN RCM  
583 used to derive forcing factors for downscaling in the alpine context (Rousselot et al., 2012) is  
584 considered relevant for high temporal frequency climate studies in the Euro-Mediterranean region  
585 (Nabat et al., 2020), but substantial improvement in simulating spatial patterns and annual cycles  
586 can be achieved with an ensemble of RCMs (Fantini et al., 2018). In our study, the use of ensemble-  
587 mean results to estimate mean trends for alternative RCMs does not take into account the variability  
588 associated with different RCMs, which respond differently to the same emission scenarios  
589 (Supplementary material, section 3), and represents a simplification in the modelling design.

590 Also in the case of vegetation response to CO<sub>2</sub> enrichment, there are large uncertainties in the  
591 (complex) climate model projections and underlying scenarios. The extent of our imperfect  
592 knowledge of the processes (and their interactions) embedded in models (epistemic uncertainties)  
593 is reflected in climate modelling, where it mainly concerns atmospheric and biosphere physics,  
594 ocean-atmosphere coupling, embedded empirical relationships, parameterisation and spatial

595 resolution. At smaller scales, impact models (such as the grassland models used here) may suffer  
596 from the omission or lack of consideration of long-term climate-related processes (e.g. plant  
597 acclimation; Sándor et al., 2018) and their interactions (model structure), as well as changes in  
598 parameter values due to new climate conditions (which may require re-parameterisation of the  
599 model, e.g. Ben Touhami et al., 2015). Estimates of future conditions should be presented for  
600 review by stakeholders, who can then assess the different levels of uncertainty in the multiple  
601 elements and estimate their own relevant projections based on their own area of expertise. In  
602 climate-change impact studies, a range of emission scenarios are used to feed climate models and,  
603 in turn, impact models. Uncertainty accumulates and propagates throughout the process of climate-  
604 change projection and impact assessment, which is carried out by developing fine-scale climate  
605 data from coarse-scale climate models and feeding the resulting local-scale scenarios into impact  
606 models to determine impacts and assess possible adaptations (Bellocchi et al., 2015). Emission  
607 scenarios are neither forecasts nor policy recommendations (Moss et al., 2010) but are selected to  
608 map a broad range of climate outcomes for further research and assessment, including impact-  
609 modelling studies. In particular, policies contrary to those established to discuss extreme situations  
610 (Cao et al., 2022) should not lead to a misuse of RCP8.5 as a no-climate-policy baseline (Pilke and  
611 Ritchie, 2021).

612 Our work demonstrates the value gained by conducting a process to assess the ability of a  
613 modelling framework and remotely sensed products to represent past (observational) and future  
614 (projected) conditions (two-time frames), based on the outputs of two climate scenarios (radiative  
615 forcing), three climate models (climate forcing) and two grassland models (impacts and  
616 adaptations). For the latter, the study identified grassland models sufficiently contrasted in their  
617 ability to represent processes controlling the dynamics of energy, water and C-N cycles. This  
618 choice was made in order to better assess changes in model estimates as a result of changing

619 assumptions and to identify the most significant ones that reflect reality. In addition, this approach  
620 helps to identify the range of variability over which model outputs may vary. In the selection phase,  
621 two models were identified in which processes are represented at different levels of detail. While  
622 PaSim is a complex (grassland-specific) model, simulating C-N and water cycles in detail (with a  
623 module dedicated to livestock impact and livestock-atmosphere feedbacks), DayCent is a more  
624 empirical (generic) model, with relatively simple relationships between driving variables and  
625 fluxes. The two models differ in the representation of soil properties, vegetation type, agricultural  
626 practices and environmental forcing, as well as in the initialisation of C pools. The common feature  
627 of both models is that they were both designed to be applied considering the grassland community  
628 as a crop with parameters describing the morphological and physiological characteristics of the  
629 vegetation set at values that should represent the mean traits of the community. Consequently, their  
630 ability to characterise interactions in multi-species grasslands is limited, beyond simple mixtures  
631 of legumes and grasses (Van Oijen et al., 2020).

632

#### 633 4.2. *Analysis of climate-change impacts and adaptation strategies*

634 The two impact models adopted agreed in the representation of impacts such as the timing and  
635 extent of the growing season and C-N fluxes, whilst divergences were observed for other outputs  
636 (e.g. biomass production and peak production). The longer growing season simulated by both  
637 models was mainly driven by the extension of the potential growing season in spring, whilst it was  
638 limited during autumn-winter, reflecting long-term observations in large alpine regions  
639 (Barichivich et al., 2013; Ernakovich et. al., 2014; Chen and Yang, 2020). The agreement in the  
640 GS outputs suggests the ability of the models to reproduce the changing seasonality of  
641 photosynthesis in vegetation, including the beginning and end of the growing season. Although  
642 both models were widely applied in various contexts (e.g. Calanca et al., 2007; Abdalla et al., 2010;

643 Vital et al., 2013; Ma et al., 2015; Ben Touhami and Bellocchi, 2015; Pulina et al., 2018; Fitton et  
644 al., 2019; Fucks et al., 2019, 2020; Melo-Damian et al., 2021), this is one of the few studies  
645 reproducing the dynamics in alpine pastoral environments, whose multifaceted structure of territory  
646 and vegetation coupled with extreme weather conditions is difficult to parameterise due to limited  
647 ground-based data for initialisation, calibration and assessment, the complex response of the  
648 vegetation growth and information on critical thresholds (e.g. air temperatures, water needs,  
649 radiation use efficiency) for mixed plant communities. As evidence of this, the mean plant growth  
650 trend simulated with both models (20-year means, Fig. 5) does not seem to reflect the observed  
651 pattern of very slow or no growth during the snow season, followed by a rapid increase in biomass  
652 accumulation as the snow melts. Generally, after the onset of growth in May or early June, plants  
653 grow rapidly, and daily dry matter accumulation can reach a maximum within a few weeks.  
654 Grassland models in their current state are not designed to properly represent such a progression  
655 of biomass accumulation with the rapid attainment of the first biomass peak. If this behaviour was  
656 roughly captured with the calibration work (over three years of AGB data; Fig. S2), and  
657 discrepancies appeared less evident with PaSim. Also, the lack of overlap between snow-cover  
658 periods and vegetation growing seasons was due to limitations inherent to the modelling of  
659 complex ecosystems. Specifically, biogeochemical models are often unable to discriminate the  
660 presence of snow and biomass at the same time on a given surface unit, but instead work separately  
661 on it. Thus, to avoid speculation on the residual amount of snow cover and, therefore, period  
662 overlap, the two components were showed separately. Considering that biophysical data show a  
663 large degree of variability, and that discrepancies between model estimates and actual data are  
664 common despite calibration efforts, research avenues can be opened from this assessment to  
665 improve simulation models for such harsh environments as high altitude mountains, where snow  
666 beds melt late and plants grow rapidly, but the rapid growth period is short during the middle of

667 the growing season, and this pattern evolves with changing climate (e.g. Wang et al., 2020). The  
668 extension of data series is indeed critical, as while calibrated models provide an adequate  
669 description of the available data, they may not capture known trends over long time horizons  
670 (Bellocchi et al., 2010).

671 Regarding C-N fluxes, the two grassland models agreed to some extent across study areas and  
672 pasture macro-types, with differences in magnitude and patterns likely associated with the inherent  
673 structure of each specific C and N sub-model (e.g. Cavalli et al., 2019). The higher C uptake  
674 estimated by DayCent compared to PaSim (Figs. S12, S13 and S14) reflects the fact that PaSim  
675 estimates the animal contribution to ecosystem respiration, which is not accounted for in DayCent's  
676 C budget. Similarly, DayCent estimated limited CH<sub>4</sub> emissions since (unlike PaSim) it does not  
677 account for fermentative digestion in its C-N sub-model. The magnitude of N<sub>2</sub>O emissions was  
678 instead similar between the two models for each study area and macro-type. Overall, the analysis  
679 also revealed that GHG dynamics were mainly driven by weather variables in DayCent and by  
680 livestock management in PaSim. In this way, PaSim estimates of CH<sub>4</sub>, and CO<sub>2</sub> emissions may  
681 better reflect observational studies that underpin the impact of management on the annual C cycle  
682 of grassland systems (as well as cropping systems), in addition to the variability of local  
683 environmental drivers (Pinares-Patiño et al., 2007; Ceschia et al., 2010; Zeeman et al., 2010, 2019).  
684 In particular, the importance of quantifying direct CO<sub>2</sub> emissions from grazing animals is  
685 emphasised (e.g. Pinares-Patiño et al., 2007). In the absence of observational data to compare with  
686 the simulation results, we refer to the literature on the C sequestration capacity of grasslands, which  
687 reports contrasting results similar to those of Table 4, which do not exclude that mountain  
688 grasslands may oscillate between being sinks and moderate sources (e.g. Zeeman et al., 2010).

689 Although the two grassland models generally agree in their impact projections, they often differ  
690 in essential details, for instance with regard to future peak pasture production. Among the reasons

691 for these different results is the uncertainty inherent in the structure of the models, as well as  
692 uncertainty in their parameterisation (and the generalisation of the resulting sets of parameter  
693 values to broad regional studies), which in turn makes the projections themselves uncertain. When  
694 considering the influence of these uncertainties on the interpretation and understanding of the  
695 projections, but also on the direction of the research, it is of great value to know the factors behind  
696 these uncertainties (Dietze et al., 2018). DayCent and PaSim showed different responsiveness to  
697 water-related factors. The water-limited growth of DayCent rather hinders biomass growth, which,  
698 according to climate scenarios, peaks before the start of grazing. With DayCent, the projected  
699 scenarios indicate that the water deficit could be the limiting factor for summer growth, which  
700 could be lower than in the near-past climate baseline. Consequently, DayCent projections towards  
701 a greater C sourcing in the PNE are logically associated with water stress and water-limited biomass  
702 production estimated by this model under future scenarios (Fig. 7), which limits photosynthetically  
703 assimilated C (gross primary production). This condition requires further study of the uncertainty  
704 associated with the model processes. Thus, the mean responses of alpine pasture production (and  
705 related outputs) to climate change, obtained with two impact models, should be considered as two  
706 extreme situations with respect to plausible future realisations: without water stress (liberal PaSim  
707 approach) and with water stress (conservative DayCent approach).

708 However, there may also be biases in the simulation of SWC, with grassland models that may  
709 not be accurate enough to estimate these dynamics. This is often associated with an unrealistically  
710 low amplitude of the annual cycle (fluctuation damping, after Wu et al., 2002) of the soil water  
711 content curve compared to field measurements (Sándor et al., 2017). It is known that the quality of  
712 soil water content simulations can seriously affect model outputs. In case of poor estimation of soil  
713 water content, model calibration may result in biased parameter values. Several factors, such as  
714 permanent wilting point, root distribution and maximum transpiration rate, are in fact related to the



715 rate of water infiltration into the soil during precipitation events and snowmelt periods (Philip,  
716 1993), which would require detailed datasets for an accurate description. Because of the known  
717 role of soil water content in determining evapotranspiration rates, stomatal conductance and other  
718 processes, this issue has obvious implications for sites and seasons where water shortage is a typical  
719 feature. The response of the models to water-limited conditions is thus questionable, which means  
720 that the applicability of the models in semi-arid or arid pastoral systems may not always be  
721 supported. This is to some extent related to the ability of roots to extract water from the soil (Volaire  
722 and Lelièvre, 2001). The usefulness of soil water content estimation is not as straightforward as for  
723 other variables and it is clear the development of improved models is fundamentally necessary for  
724 soil hydrology, to rectify structural errors in models and to avoid systematic errors associated with  
725 some of the model parameters. These may involve further study of runoff, diffusion and percolation  
726 processes, while accounting for features such as ponding water formation, and snowmelt and  
727 groundwater movement (Hidy et al., 2016).

728

#### 729 4.3. *Effect of adaptation measures*

730 The climate-change scenarios exhibited an increase in the air temperature together with higher  
731 winter precipitation and prolonged drier conditions during the spring-summer period. This  
732 condition logically translates into an accentuation of seasonality, with faster snowmelt and a longer  
733 growing season (i.e. +15 to +40 days), providing higher estimated yields and evapotranspiration in  
734 both study areas. In this perspective, earlier grazing dates and changes in livestock density, which  
735 were *a priori* hypothesised as adaptation options, proved to be coherent to cope with these projected  
736 changes.

737 The earlier grazing date better matched the future biomass peaks simulated by DayCent, also  
738 agreeing with observations in alpine regions (i.e. Xu et al., 2016) and thus resulting in a more

739 efficient use of pastures. Although this pattern paved the way for increasing the number of grazing  
740 events during the growing season, reduced biomass regrowth due to dry summer conditions  
741 inhibited the possibility of further grazing. Reduced soil water availability and increased number  
742 of days of heat stress can lead to stomatal closure and inhibit biomass production in summer, which  
743 must be taken into account in addition to the increase in plant photosynthetic rates with increasing  
744 CO<sub>2</sub> concentration in the atmosphere, not to mention the possible degradation of grasslands due to  
745 severe summer drought episodes (Moreau and Lorgeou, 2007). For that, any excess water in winter  
746 can be used to reduce the water deficit when the soil is drier than the field capacity. While this  
747 would hint at the possibility of using irrigation even at high altitudes to extend the number of  
748 grazing events, the balance between costs (e.g. of energy and irrigation systems) and benefits (e.g.  
749 increased end product) would need to be thoroughly investigated. In contrast, in the scenario  
750 depicted by PaSim, which would be too liberal in the sense that it simulates optimal growing  
751 conditions in midsummer (i.e. without an expectation of summer water stress), this adaptation  
752 strategy would not be necessary.

753 The model results indicate higher biomass production when LD is increased (although changes  
754 in livestock density do not particularly modify production levels). This result is likely due to a  
755 higher N availability to plants provided by a higher amount of N excreted in faeces and urine which,  
756 together with a simulated non-linear effect of grazing on production, led to a faster and higher  
757 biomass regrowth rate. While both models consider N from excretion and the effect of grazing on  
758 production, detrimental effects on biomass production due to the impact of soil compaction when  
759 animal density increases were not considered. However, higher animal numbers may increase soil  
760 compaction, resulting in poor water retention and altered (slower) mineralisation processes that  
761 may reduce biomass regrowth and forage quality (Li et al., 2017). In this perspective, the response  
762 of biomass growth to changes in LD could be partly overestimated or affected by a certain level of

763 uncertainty in both models. Despite these limitations, we can conclude that the alpine region is set  
764 to become warmer and wetter, and that yields in these areas are highly dependent on both water  
765 availability and the type of management adopted. The projected climate scenarios and adaptation  
766 options considered are not expected to substantially worsen the GHG balance, although a caveat is  
767 that C sequestration by pasturelands may be reduced in a warmer climate. However, there is a need  
768 to further develop and evaluate grassland models for key processes and outputs, such as CO<sub>2</sub> and  
769 non-CO<sub>2</sub> emissions, as well as to systematically and more accurately characterise the extent and  
770 timing of human intervention in a range of grazing areas covering broader climatic gradients.

771

## 772 **5. Concluding remarks**

773 Research on mountain pastures in two western alpine parks shows that variations in climate-  
774 change impacts and adaptations of these systems are linked to natural and anthropogenic factors to  
775 different degrees depending on the pastoral macro-type class studied (defined by an altitudinal  
776 productivity gradient). While the use of modelling approaches and remote-sensing products in  
777 vulnerability studies is not new *per se*, the integration of these tools within alpine pastoral  
778 communities has a point of originality, as the analysis carried out can help to solve  
779 multidisciplinary challenges such as which areas are vulnerable and how they compare under harsh  
780 climatic conditions. The findings of this study indicate an increase in the length of the growing  
781 season by 15 to 40 days, leading to expected changes in the timing and amount of biomass  
782 production and a likely decrease in biomass regrowth during the summer season due to prolonged  
783 drought conditions. The greatest uncertainties were found in the GHG balance and mitigation  
784 capacity of alpine pastures, where contrasting patterns were observed between the impact models  
785 used (ranging from -350 to +100 g C m<sup>-2</sup> yr<sup>-1</sup> for NEE), mainly due to the different flux simulation  
786 approaches. Similarly, earlier grazing dates appeared to be the most suitable adaptation strategy,

787 especially when combined with increasing livestock density, while decreasing livestock density  
788 did not show any significant change.

789 The elaboration of adaptation measures, carried out in this study with local herding and farming  
790 communities, provides a basis for appropriate agricultural policy and land management measures  
791 adapted to ongoing climate change. However, although different modelling approaches are able to  
792 capture distinct aspects of the adaptive process, they tend to be applied in relative isolation, without  
793 producing unified representations. The corollary of this is that the usefulness of future projections  
794 of climate-change impacts by grassland models, such as those represented here, is strongly  
795 influenced by the quality of the climate model data used to run them and the field data used to  
796 calibrate them. Social impact assessment studies are now needed to examine how  
797 production/biophysical/biogeochemical impacts, i.e. the effects of climatic anomalies on alpine  
798 pasture performances, propagate through the socio-economic and political system. Such an  
799 integrated approach, which would include the potential for adaptation and adjustment to climate  
800 pressure, would reflect the reality of pastoral communities much better than the modelling used  
801 and raises fruitful research questions on the vulnerability of alpine territories and their adaptive  
802 capacity.

803

#### 804 **Acknowledgement**

805 The present work was produced under the co-finance of the EC LIFE programme for the  
806 Environment and Climate Action (2014-2020) in the framework of the Project LIFE PASTORALP  
807 ‘Pastures vulnerability and adaptation strategies to climate change impacts in the Alps’  
808 (LIFE16/CCA/IT/000060). It falls within the thematic area of the French government IDEX-ISITE  
809 initiative (reference: 16-IDEX-0001; project CAP 20-25). The authors wish to acknowledge the  
810 commitment of Dehia Hadjsaadi (Université Clermont Auvergne, INRAE, VetAgro Sup, UREP,

811 Clermont-Ferrand, France), who supported PaSim simulations in 2020-2021. The research contract  
812 of the author Luisa Leolini was co-funded by the European Union - PON Research and Innovation  
813 2014-2020 in accordance with Article 24, paragraph 3a), of Law No. 240 of December 30, 2010,  
814 as amended, and Ministerial Decree No. 1062 of August 10, 2021.

815

816

817 **References**

- 818 Abdalla, M., Jones, M., Yeluripati, J., Smith, P., Burke, J., Williams, M., 2010. Testing DayCent  
819 and DNDC model simulations of N<sub>2</sub>O fluxes and assessing the impacts of climate change on the  
820 gas flux and biomass production from a humid pasture. *Atmos. Environ.* 44, 2961–2970.  
821 <https://doi.org/10.1016/j.atmosenv.2010.05.018>
- 822 Albanito, F., McBey, D., Harrison, M., Smith, P., Ehrhardt, F., Bhatia, A., Bellocchi, G., Brillì, L.,  
823 Carozzi, M., Christie, K., Doltra, J., Dorich, C., Doro, L., Grace, P., Grant, B., Léonard, J.,  
824 Liebig, M., Ludemann, C., Martin, R., Meier, E., Meyer, R., De Antoni Migliorati, M.,  
825 Myrgiotis, V., Recous, S., Sándor, R., Snow, V., Soussana, J.F., Smith, W.N., Fitton, F., 2022.  
826 How Modelers Model: the Overlooked Social and Human Dimensions in Model  
827 Intercomparison Studies. *Environmental Science & Technology* 2022 56 (18), 13485-13498.  
828 <https://doi.org/10.1021/acs.est.2c02023>
- 829 Alessa, L., Kliskey, A., Gosz, J., Griffith, D., & Ziegler, A., 2018. MtnSEON and social–ecological  
830 systems science in complex mountain landscapes. *Frontiers in Ecology and the Environment*,  
831 16(S1), S4-S10. <https://doi.org/10.1002/fee.1753>
- 832 Altaweel, M., Virapongse, A., Griffith, D., Alessa, L., & Kliskey, A., 2015. A typology for  
833 complex social-ecological systems in mountain communities. *Sustainability: Science, Practice  
834 and Policy*, 11(2), 1-13. <https://doi.org/10.1080/15487733.2015.11908142>
- 835 Ayar, P.V., Vrac, M., Bastin, S., Carreau, J., Déqué, M., Gallardo, C., 2016. Intercomparison of  
836 statistical and dynamical downscaling models under the EURO-and MED-CORDEX initiative  
837 framework: present climate evaluations. *Climate Dynamics* 46, 1301-1329.  
838 <https://doi.org/10.1007/s00382-015-2647-5>
- 839 Barichivich, J., Briffa, K.R., Myneni, R.B., Osborn, T.J., Melvin, T.M., Ciais, P., Piao, S., Tucker,  
840 C., 2013. Large-scale variations in the vegetation growing season and annual cycle of

841 atmospheric CO<sub>2</sub> at high northern latitudes from 1950 to 2011. *Global Change Biology* 19,  
842 3167-3183. <https://doi.org/10.1111/gcb.12283>

843 Bellocchi, G., Rivington, M., Donatelli, M., Matthews, K., 2010. Validation of biophysical models:  
844 issues and methodologies. A review. *Agronomy for Sustainable Development* 30, 109-130.

845 Bellocchi, G., Rivington, M., Matthews, K., Acutis, M., 2015. Deliberative processes for  
846 comprehensive evaluation of agroecological models. A review. *Agronomy for Sustainable*  
847 *Development* 35, 589-605. <https://doi.org/10.1051/agro/2009001>

848 Ben Touhami, H., Bellocchi, G., 2015. Bayesian calibration of the Pasture Simulation model  
849 (PaSim) to simulate European grasslands under water stress. *Ecological Informatics* 30, 356-  
850 364. <https://doi.org/10.1016/j.ecoinf.2015.09.009>

851 Ben Touhami, H., Lardy, R., Barra, V., Bellocchi, G., 2013. Screening parameters in the Pasture  
852 Simulation model using the Morris method. *Ecological Modelling* 266, 42-57.

853 Bengtsson, J., Bullock, J.M., Egoh, B., Everson, C., Everson, T., O'Connor, T., O'Farrell, P.J.,  
854 Smith, H.G., Lindborg, R., 2019. Grasslands - more important for ecosystem services than you  
855 might think. *Ecosphere* 10, e02582. <https://doi.org/10.1002/ecs2.2582>

856 Bonet, R., Arnaud, F., Bodin, X., Bouche, M., Boulangeat, I., Bourdeau, P., Bouvier, M., Cavalli,  
857 L., Choler, P., Delestrade, A., Dentant, C., Dumas, D., Fouinat, L., Gardent, M., Lavergne, S.,  
858 Lavorel, S., Naffrechoux, E., Nellier, Y., Perga, M.-E., Sagot, C., Senn, O., Thibert, E., Thuiller,  
859 W., 2016. Indicators of climate: Ecrins National Park participates in long-term monitoring to  
860 help determine the effects of climate change. *Journal on Protected Mountain Areas Research*  
861 *and Management* S. 44-52. <https://doi.org/10.1553/eco.mont-8-1s3>

862 Bornard A., Bassignana M., Bernard-Brunet C., Labonne S., Cozic Ph., 2007. Les végétations  
863 d'alpage de la Vanoise. Description agro-écologique et gestion pastorale. Éd. Quae, Versailles.  
864 (in French)

865 Brien, P., 2018. Analyse de l'évolution de l'utilisation des alpages sur le territoire du Parc National  
866 des Écrins (1996-2012), perceptions des tendances en cours et des enjeux à venir. L'exemple de  
867 la prédation et du changement climatique en alpage. Rapport de stage en entreprise, Université  
868 Savoie-Mont Blanc. (in French)

869 Brillì, L., Bechini, L., Bindi, M., Carozzi, M., Cavalli, D., Conant, R., Dorich, C.D., Doro, L.,  
870 Ehrhardt, F., Farina, R., Ferrise, R., Fitton, N., Francaviglia, R., Grace, P., Iocola, I., Klumpp,  
871 K., Léonard, J., Martin, R., Massad, R.S., Recous, S., Seddaiu, G., Sharp, J., Smith, P., Smith,  
872 W.N., Soussana, J.-F., Bellocchi, G., 2017. Review and analysis of strengths and weaknesses of  
873 agro-ecosystem models for simulating C and N fluxes. *Sci. Total Environ.* 598, 445-470.  
874 <https://doi.org/10.1016/j.scitotenv.2017.03.208>

875 Caballero, R., Fernández-González, F., Pérez Badia, R., Molle, G., Roggero, P.P., Bagella, S.,  
876 D'Ottavio, P., Papanastasis, V.P., Fotiadis, G., Sidiropoulou, A., Ispikoudis, I., 2009. Grazing  
877 systems and biodiversity in Mediterranean areas: Spain, Italy and Greece. *Pastos* 39, 9-152.

878 Calanca, P., 2007. Climate change and drought occurrence in the Alpine region: How severe are  
879 becoming the extremes? *Global and Planetary Change* 57, 151-160.  
880 <https://doi.org/10.1016/j.gloplacha.2006.11.001>

881 Calanca, P., Vuichard, N., Campbell, C., Viovy, N., Cozic, A., Fuhrer, J., Soussana, J.F., 2007.  
882 Simulating the fluxes of CO<sub>2</sub> and N<sub>2</sub>O in European grasslands with the Pasture Simulation  
883 Model (PaSim). *Agriculture, Ecosystems & Environment* 121, 164-174.  
884 <https://doi.org/10.1016/j.agee.2006.12.010>

885 Cao, W., Duan, C., Yang, T., Wang, S., 2022. Higher Heat Stress Increases the Negative Impact  
886 on Rice Production in South China: A New Perspective on Agricultural Weather Index  
887 Insurance. *Atmosphere*, 13, 1768. <https://doi.org/10.3390/atmos13111768>



888 Cavallero A., Aceto P., Gorlier A., Lombardi G., Lonati M., Martinasso B., Tagliatori C., 2007. I  
889 tipi pastorali delle Alpi piemontesi. A. Perdisa Editore, Bologna. pp XII-467 p., ill. ,  
890 EAN:9788883723216

891 Cavalli, D., Bellocchi, G., Corti, M., Marino Gallina, P., Bechini, L., 2019. Sensitivity analysis of  
892 C and N modules in biogeochemical crop and grassland models following manure addition to  
893 soil. *Eur. J. Soil Sci.* 70, 833-846. <https://doi.org/10.1111/ejss.12793>

894 Ceschia, E., Béziat, P., Dejoux, J.F., Aubinet, M., Bernhofer, Ch., Bodson, B., Buchmann, N.,  
895 Carrara, A., Cellier, P., Di Tommasi, P., Elbers, J.A., Eugster, W., Grünwald, T., Jacobs, C.M.J.,  
896 Jans, W.W.P., Jones, M., Kutsch, W., Lanigan, G., Magliulo, E., Marloie, O., Moors, E.J.,  
897 Moureaux, C., Olioso, A., Osborne, B., Sanz, M.J., Saunders, M., 2010. Management effects on  
898 net ecosystem carbon and GHG budgets at European crop sites. *Agriculture, Ecosystems and*  
899 *Environment* 139, 363–383. <https://doi.org/10.1016/j.agee.2010.09.020>

900 Chen, X., Yang, Y., 2020. Observed earlier start of the growing season from middle to high  
901 latitudes across the Northern Hemisphere snow-covered landmass for the period 2001–2014.  
902 *Environ. Res. Lett.* 15: 034042. <https://doi.org/10.1088/1748-9326/ab6d39>

903 Chow, L., Xing, Z., Rees, H.W., Meng, F., Monteith, J., Stevens, L., 2009. Field performance of  
904 nine soil water content sensors on a sandy loam soil in New Brunswick, Maritime Region,  
905 Canada. *Sensors* 9, 9398-9413. <https://doi.org/10.3390/s9110939>

906 Corona-Lozada, M.C., Morin, S., Choler, P., 2019. Drought offsets the positive effect of summer  
907 heat waves on the canopy greenness of mountain grasslands. *Agricultural and Forest*  
908 *Meteorology.* <https://doi.org/10.1016/j.agrformet.2019.107617>

909 Della-Vedova M., Legéard J.P., 2012. Alpages sentinelles dans le Parc national des Écrins : un  
910 espace de dialogue pour anticiper l’impact des aléas climatiques. Assemblée Générale du  
911 Cerpam, La Chapelle-en-Valgaudemar (05), 18 septembre 2012. (in French)

912 Dibari, C., Bindi, M., Moriondo, M., Staglianò, N., Targetti, S., Argenti, G., 2016. Spatial data  
913 integration for the environmental characterization of pasture macrotypes in the Italian Alps.  
914 Grass and Forage Science 71, 219-234. <https://doi.org/10.1111/gfs.12168>

915 Dibari, C., Costafreda-Aumedes, S., Argenti, G., Bindi, M., Carotenuto, F., Moriondo, M.,  
916 Padovan, G., Pardini, A., Staglianò, N., Vagnoli, C., Brilli, L., 2020. Expected changes to Alpine  
917 pastures in extent and composition under future climate conditions. Agronomy 10: 926.  
918 <https://doi.org/10.3390/agronomy10070926>

919 Dibari, C., Pulina, A., Argenti, G., Aglietti, C., Bindi, M., Moriondo, M., Mula, L., Pasqui, M.,  
920 Seddaiu, G., Roggero, P.P., 2021. Climate change impacts on the Alpine, Continental and  
921 Mediterranean grassland systems of Italy: A review. Italian Journal of Agronomy, 16:1843.  
922 <https://doi.org/10.4081/ija.2021.1843>

923 Dietze, M.C., Fox, A., Beck-Johnson, L.M., Betancourt, J.L., Hooten, M.B., Jarnevich, C.S., Keitt,  
924 T.H., Kenney, M.A., Laney, C.M., Larsen, L.G., Loescher, H.W., Lunch, C.K., Pijanowski,  
925 B.C., Randerson, J.T., Read, E.K., Tredennick, A.T., Vargas, R., Weathers, K.C., White, E.P.,  
926 2018. Iterative near-term ecological forecasting: Needs, opportunities, and challenges.  
927 Proceedings of the National Academy of Sciences of the United States of America 115, 1424-  
928 1432. <https://doi.org/10.1073/pnas.1710231115>

929 Dumont, B., Farruggia, A., Garel, J.-P., Bachelard, P., Boitier, E., Frain, M., 2009. How does grazing  
930 intensity influence the diversity of plants and insects in a species-rich upland grassland on basalt  
931 soils? Grass and Forage Science 64, 92–105. <https://doi.org/10.1111/j.1365-2494.2008.00674.x>

932 EC, 2008. Regulation (EC) No 1166/2008 of the European Parliament and of the Council of 19  
933 November 2008 on farm structure surveys and the survey on agricultural production methods  
934 and repealing Council Regulation (EEC). Official Journal of the European Union, Brussels.

935 EC, 2013. Regulation (EC) No 1306/2013 of the European Parliament and of the Council of 17  
936 December 2013 on support for rural development by the European Agricultural Fund for Rural  
937 Development (EAFRD) and repealing Council Regulation (EC) No 1698/2005. Official Journal  
938 of the European Union, Brussels.

939 Engler R, Randin C, Thuiller W, Dullinger S, Zimmermann NE, Araújo MB, Pearman PB, Le Lay  
940 G, Piedallu C, Albert CH, Choler P, Coldea G, De Lamo X, Dirnböck T, Gégout J-C, Gómez-  
941 García D, Grytnes J-A, Heegaard E, Høistad F, Nogués-Bravo D, Normand S, Puşcaş M,  
942 Sebastià M-T, Stanisci A, Theurillat J-P, Trivedi MR, Vittoz P, Guisan A., 2011. 21st century  
943 climate change threatens mountain flora unequally across Europe. *Global Change Biology* 17,  
944 2330-2341. <https://doi.org/10.1111/j.1365-2486.2010.02393.x>

945 Ernakovich, J.G., Hopping, K.A., Berdanier, A.B., Simpson, R.T., Kachergis, E.J., Steltzer, H.,  
946 Wallenstein, M.D., 2014. Predicted responses of arctic and alpine ecosystems to altered  
947 seasonality under climate change. *Glob Chang Biol* 20, 3256-3269.  
948 <https://doi.org/10.1111/gcb.12568>

949 Fantini, A., Raffaele, F., Torma, C., Bacer, S., Coppola, E., Giorgi, F., Ahrens, B., Dubois, C.,  
950 Sanchez, E., Verdecchia, M., 2018. Assessment of multiple daily precipitation statistics in ERA-  
951 Interim driven Med-CORDEX and EURO-CORDEX experiments against high resolution  
952 observations. *Climate Dynamics* 51, 877-900. <https://doi.org/10.1007/s00382-016-3453-4>

953 Felber R., Bretscher D., Münger A., Neftel A., Ammann C., 2016. Determination of the carbon  
954 budget of a pasture: effect of system boundaries and flux uncertainties. *Biogeosciences* 13,  
955 2959-2969. <https://doi.org/10.5194/bg-13-2959-2016>

956 Filippa, G., Cremonese, E., Migliavacca, M., Galvagno, M., Sonnentag, O., Humphreys, E.,  
957 Hufkens, K., Ryu, Y., Verfaillie, J., di Cella, U.M., Richardson, A.D., 2015. Five years of  
958 phenological monitoring in a mountain grassland: inter-annual patterns and evaluation of the

959 sampling protocol. *Int J Biometeorol* 59, 1927–1937. [https://doi.org/10.1007/s00484-015-0999-](https://doi.org/10.1007/s00484-015-0999-5)  
960 [5](https://doi.org/10.1007/s00484-015-0999-5)

961 Filippa, G., Cremonese, E., Galvagno, M., Bayle, A., Choler, P., Bassignana, M., Piccot, A.,  
962 Poggio, L., Oddi, L., Gascoin, S., Costafreda-Aumedes, S., Argenti, A., Dibari, C., 2022. On  
963 the distribution and productivity of mountain grasslands in the Gran Paradiso National Park,  
964 NW Italy: A remote sensing approach. *International Journal of Applied Earth Observations and*  
965 *Geoinformation*. <https://doi.org/10.1016/j.jag.2022.102718>

966 Fitton, N., Bindi, M., Brilli, L., Cichota, R., Dibari, C., Fuchs, K., Huguenin-Elie, O., Klumpp, K.,  
967 Lieffering, M., Lüscher, A., Martin, R., McAuliffe, R., Merbold, L., Newton, P., Rees, R.M.,  
968 Smith, P., Topp, C.F.E., Snow, V., 2019. Modelling biological N fixation and grass-legume  
969 dynamics with process-based biogeochemical models of varying complexity. *European Journal*  
970 *of Agronomy* 106, 58-66. <https://doi.org/10.1016/j.eja.2019.03.008>

971 Fitton, N., Datta, A., Smith, K., Williams, J.R., Hastings, A., Kuhnert, M., Topp, C.F.E., Smith, P.,  
972 2014. Assessing the sensitivity of modelled estimates of N<sub>2</sub>O emissions and yield to input  
973 uncertainty at a UK cropland experimental site using the DailyDayCent model. *Nutrient Cycling*  
974 *in Agroecosystems* 99, 119-133. <https://doi.org/10.1007/s10705-014-9622-0>

975 Fuchs, K., Merbold, L., Buchmann, N., Bellocchi, G., Bindi, M., Brilli, L., Conant, R.T., Dorich,  
976 C.D., Ehrhardt, F., Fitton, N., Grace, P., Klumpp, K., Liebig, M., Lieffering, M., Martin, R.,  
977 McAuliff, R., Newton, P.C.D., Rees, R.M., Recous, S., Smith, P., Soussana, J.F., Topp, C.F.E.  
978 and Snow, V., 2020. Evaluating the potential of legumes to mitigate N<sub>2</sub>O emissions from  
979 permanent grassland using process-based models. *Global Biogeochemical Cycles* 34:  
980 e2020GB006561. <https://doi.org/10.1029/2020GB006561>

981 Fuchs, K., Merbold, L., Buchmann, N., Bretscher, D., Brilli, L., Fitton, N., Topp, C.F.E., Klumpp,  
982 K., Lieffering, M., Martin, R., Newton, P.C.D., Rees, R.M., Rolinski, S., Smith, P., Snow, V.,

983 2019. Multi-model evaluation of nitrous oxide emissions from an intensively managed  
984 grassland. *Journal of Geophysical Research: Biogeosciences* 125: e2019JG005261.  
985 <https://doi.org/10.1029/2019JG005261>

986 Gobiet A., Kotlarski S., Beniston M., Heinrich G., Rajczak J., Stoffel M., 2014. 21st century  
987 climate change in the European Alps - A review. *Science of the Total Environment* 493, 1138-  
988 1151. <https://doi.org/10.1016/j.scitotenv.2013.07.050>

989 Gottfried M., Pauli H., Futschik A., Akhalkatsi M., Baranok P., Alonso J.L.B., Coldea G., Jan D.,  
990 Erschbamer B., Calzado M.R.F., Kazakis G., Kraji J., Larsson P., Mallaun M., Michelsen O.,  
991 Moiseev D., Moiseev P., Molau U., Merzouki A., Nagy L., Nakhutsrishvili G., Pedersen B.,  
992 Pelino G., Puscas M., Rossi G., Stanisci A., Theurillat J.P., Tomaselli M., 2012. Continent-wide  
993 response of mountain vegetation to climate change. *Nature Climate Change* 2, 111-115.  
994 <https://doi.org/10.1038/nclimate1329>

995 Graux, A.-I., Gaurut, M., Agabriel, J., Baumont, R., Delagarde, R., Delaby, L., Soussana, J.-F.,  
996 2011. Development of the Pasture Simulation Model for assessing livestock production under  
997 climate change. *Agriculture, Ecosystems & Environment* 144, 69-91.  
998 <https://doi.org/10.1016/j.agee.2011.07.001>

999 Hidy, D., Barcza, Z., Marjanovič, H., Ostrogovič Sever, M.Z., Dobor, L., Gelybó, Gy., Fodor, N.,  
1000 Pintér, K., Churkina, G., Running, S.W. Thornton, P.E., Bellocchi, G., Haszpra, L., Horváth, F.,  
1001 Suyker, A., Nagy, Z., 2016. Terrestrial ecosystem process model Biome-BGCMuSo: summary  
1002 of improvements and new modeling possibilities. *Geoscientific Model Development* 9, 4405-  
1003 4437. <https://doi.org/10.5194/gmd-9-4405-2016>

1004 Holman, I.P., Brown, C., Carter, T.R., Harrison, P.A., Rounsevell, M., 2019. Improving the  
1005 representation of adaptation in climate change impact models. *Regional Environmental Change*  
1006 19, 711-721. <https://doi.org/10.1007/s10113-018-1328-4>

1007 Jouglet, J-P., 1999. Les végétations des alpages des Alpes françaises du Sud. Guide technique pour  
1008 la reconnaissance et la gestion des milieux pâturés d'altitude. Cemagref Éditions and ATEN  
1009 (Atelier Technique des Espaces Naturels). (in French)

1010 Jourdain-Annequin, C., Duclos, J.-C., 2006. Aux origines de la transhumance : les Alpes et la vie  
1011 pastorale d'hier à aujourd'hui. Paris: Picard. (in French)

1012 Kollas, C., Kersebaum, K.C., Nendel, C., Manevski, K., Müller, C., Palosuo, T., Armas-Herrera,  
1013 C.M., Beaudoin, N., Bindi, M., Charfeddine, M., Conradt, T., Constantin, J., Eitzinger, J., Ewert,  
1014 F., Ferrise, R., Gaiser, T., deCortazar-Atauri, I.G., Giglio, L., Hlavinka, P., Hoffmann, H.,  
1015 Hoffmann, M.P., Launay, M., Manderscheid, M., Mary, B., Mirschel, W., Moriondo, M.,  
1016 Olesen, J.E., Öztürk, I, Pacholski, A., Ripoche-Wachter, D., Roggero, P.P., Roncossek, S.,  
1017 Rötter, R.P., Ruget, F., Sharif, B., Trnka, M., Ventrella, D., Waha, K., Wegehenkel, M., Weigel,  
1018 H.-J., Wu, L., 2015. Crop rotation modelling – a European model intercomparison. European  
1019 Journal of Agronomy 70, 98-111. <https://doi.org/10.1016/j.eja.2015.06.007>

1020 Kurtogullari, Y., Rieder, N. S., Arlettaz, R., Humbert, J. Y., 2020. Conservation and restoration of  
1021 Nardus grasslands in the Swiss northern Alps. Applied vegetation science, 23(1), 26-38.  
1022 <https://doi.org/10.1111/avsc.12462>

1023 Kurz, P., 2013. Management strategies and landscape diversity in commonly governed mountain  
1024 pastures: a case study from Austrian Alps. European Countryside, 5(3), 212-231.  
1025 <https://doi.org/10.2478/euco-2013-0014>

1026 Li, W., Cao, W., Wang, J., Li, X., Xu, C., Shi, S., 2017. Effects of grazing regime on vegetation  
1027 structure, productivity, soil quality, carbon and nitrogen storage of alpine meadow on the  
1028 Qinghai-Tibetan Plateau. Ecological Engineering 98, 123–133.  
1029 <https://doi.org/10.1016/j.ecoleng.2016.10.026>

1030 Ma, S., Lardy, R., Graux, A-I., Ben Touhami, H., Klumpp, K., Martin, R., Bellocchi, G., 2015.  
1031 Regional-scale analysis of carbon and water cycles on managed grassland systems.  
1032 Environmental Modelling & Software 72, 356-371.  
1033 <https://doi.org/10.1016/j.envsoft.2015.03.007>

1034 Mathieu, J., 2000. Storia delle Alpi 1500-1900: Ambiente, sviluppo e società (volume 4).  
1035 Bellinzona: Casagrande. (in Italian)

1036 Melo Damian, J., da Silva Matos, E., Carneiro e Pedreira, B., de Faccio Carvalho, P.C., Premazzi,  
1037 L.M., Williams, S., Paustian, K., Pellegrino Cerri, C.E., 2021. Predicting soil C changes after  
1038 pasture intensification and diversification in Brazil. Catena 202: 105238.  
1039 <https://doi.org/10.1016/j.catena.2021.105238>

1040 Moreau, J.C., Lorgeou, J., 2007. First elements for a prospective study of the effects of climatic  
1041 changes on pastures, maize and the forage systems. Fourrages Journée de l'A.F.P.F. (Association  
1042 Française pour la Production Fourragère), 191, pp 285-295, 11 p ; ref : 1/4 p, ISSN 0429-2766

1043 Moss, R.H., Edmonds, J.A., Hibbard, K.A., Manning, M.R., Rose, S.K., van Vuuren, D.P., Carter,  
1044 T.R., Emori, S., Kainuma, M., Kram, T., Meehl, G.A., Mitchell, J.F, Nakicenovic, N., Riahi, K.,  
1045 Smith, S.J., Stouffer, R.J., Thomson, A.M., Weyant, J.P., Wilbanks, T.J., 2010. The next  
1046 generation of scenarios for climate change research and assessment. Nature 463, 747-756.  
1047 <https://doi.org/10.1038/nature08823>

1048 Nabat, P., Somot, S., Cassou, C., Mallet, M., Michou, M., Bouniol, D., Decharme, B., Drugé, T.,  
1049 Roehrig, R., Saint-Martin, D., 2020. Modulation of radiative aerosols effects by atmospheric  
1050 circulation over the Euro-Mediterranean region. Atmospheric Chemistry and Physics 20, 8315-  
1051 8349. <https://doi.org/10.5194/acp-20-8315-2020>

1052 Necpálová, M., Anex, R.P., Fienen, M.N., del Grosso, S.J., Castellano, M.J., Sawyer, J.E., Iqbal,  
1053 J., Pantoja, J.L., Barker, D.W., 2015. Understanding the DayCent model: calibration, sensitivity,

1054 and identifiability through inverse modeling. *Environmental Modelling & Software* 66, 110-  
1055 130. <https://doi.org/10.1016/j.envsoft.2014.12.011>

1056 Negro, M., Rolando, A., Palestini, C., 2011. The impact of overgrazing on dung beetle diversity  
1057 in the Italian Maritime Alps. *Environmental Entomology* 40, 1081–1092.  
1058 <https://doi.org/10.1603/EN11105>

1059 Nettier, B., Dobremez, L., Coussy, J.L., Romagny, T., 2010. Attitudes of livestock farmers and  
1060 sensitivity of livestock farming systems to drought conditions in the French Alps. *Journal of*  
1061 *Alpine Research* 98-4. <http://rga.revues.org/index1294.html>

1062 Nori S., Gemini M., 2011. The Common Agricultural Policy vis-à-vis European pastoralists:  
1063 principles and practices. *Pastoralism: Research, Policy and Practice* 1: 27.  
1064 <https://doi.org/10.1186/2041-7136-1-27>

1065 Parton, W.J., Hartman, M., Ojima, D., Schimel, D., 1998. DAYCENT and its land surface  
1066 submodel: description and testing. *Global and Planetary Change* 19, 35-48.  
1067 [https://doi.org/10.1016/S0921-8181\(98\)00040-X](https://doi.org/10.1016/S0921-8181(98)00040-X)

1068 Parton, W.J., Schimel, D.S., Ojima, D., Cole, C.V., 1994. A general model for soil organic matter  
1069 dynamics: sensitivity to litter chemistry, texture and management. In: Bryant, R.B., Arnoldm,  
1070 R.W. (Eds.), *Quantitative modeling of soil forming processes*. Soil Science Society of America,  
1071 Madison, pp 147–167. <https://doi.org/10.2136/sssaspecpub39.c9>

1072 Philip, J.R., 1993. Variable-head ponded infiltration under constant or variable rainfall. *Water*  
1073 *Resources Research* 29, 2155–2165. <https://doi.org/10.1029/93WR00748>

1074 Piccot, A., Argenti, G., Bellocchi, G., Brien, P., Cremonese, E., Della-Vedova, M., Dibari, C,  
1075 Galvagno, M., Ghidotti, S., Napoléone, C., Stendardi, L., Targetti, S., Trombi, G., Varese, P.,  
1076 Bassignana, M., 2022. Adaptation policies and measures to cope with climate change in Alpine



1077 mountain farming. Proceedings. ISCRAES (International Symposium on climate-resilient agri-  
1078 environmental systems) Dublin, 28-31 August 2022

1079 Pielke, R., Ritchie, J. 2021. Distorting the view of our climate future: The misuse and abuse of  
1080 climate pathways and scenarios. *Energy Research & Social Science* 72, 101890.  
1081 <https://doi.org/10.1016/j.erss.2020.101890>

1082 Pinares-Patiño, C.S., D'Hour, P., Jouany, J.-P., Martin, C., 2007. Effects of stocking rate on  
1083 methane and carbon dioxide emissions from grazing cattle. *Agriculture, Ecosystems and*  
1084 *Environment* 121, 30–46. <https://doi.org/10.1016/j.agee.2006.03.024>

1085 Pulina, A., Lai, R., Salis, L., Seddaiu, G., Roggero, P.P., Bellocchi, G., 2018. Modelling pasture  
1086 production and soil temperature, water and carbon fluxes in Mediterranean grassland systems  
1087 with the Pasture Simulation model. *Grass and Forage Science* 73, 272-283.  
1088 <https://doi.org/10.1111/gfs.12310>

1089 Richter, K., Atzberger, C., Hank, T.B., Mauser, W., 2012. Derivation of biophysical variables from  
1090 Earth observation data: validation and statistical measures. *Journal of Applied Remote Sensing*  
1091 6: 063557. <https://doi.org/10.1117/1.JRS.6.063557>

1092 Riedo, M., Grub, A., Rosset, M., Fuhrer, J., 1998. A Pasture Simulation model for dry matter  
1093 production, and fluxes of carbon, nitrogen, water and energy. *Ecological Modelling* 105, 141-  
1094 183. [https://doi.org/10.1016/S0304-3800\(97\)00110-5](https://doi.org/10.1016/S0304-3800(97)00110-5)

1095 Rousselot, M., Durand, Y., Giraud, G., Mérindol, L., Dombrowski-Etchevers, I., Déqué, M., &  
1096 Castebrunet, H., 2012. Statistical adaptation of ALADIN RCM outputs over the French Alps -  
1097 application to future climate and snow cover. *The Cryosphere* 6, 785-805.  
1098 <https://doi.org/10.5194/tc-6-785-2012>

1099 Ruti, P.M., Somot, S., Giorgi, F., Dubois, C., Flaounas, E., Obermann, A., Dell'Aquila, A.,  
1100 Pisacane, G., Harzallah, A., Lombardi, E., Ahrens, B., Akhtar, N., Alias, A., Arsouze, T., Aznar,

1101 R., Bastin, S., Bartholy, J., Béranger, K., Beuvier, J., Bouffies-Cloch , S., Brauch, J., Cabos,  
1102 W., Calmanti, S., Calvet, J.-C., Carillo, A., Conte, D., Coppola, E., Djurdjevic, V., Drobinski,  
1103 P., Elizalde-Arellano, A., Gaertner, M., Gal n, P., Gallardo, C., Gualdi, S., Goncalves, M.,  
1104 Jorba, O., Jord , G., L’Heveder, B., Lebeaupin-Brossier, C., Li, L., Liguori, G., Lionello, P.,  
1105 Maci s, D., Nabat, P.,  nol, B., Raikovic, B., Ramage, K., Sevault, F., Sannino, G., Struglia,  
1106 M.V., Sanna, A., Torma, C., Vervatis, V., 2016. Med-CORDEX initiative for Mediterranean  
1107 climate studies. *Bulletin of the American Meteorological Society* 97, 1187-1208.  
1108 <https://doi.org/10.1175/BAMS-D-14-00176.1>

1109 S ndor, R., Barcza, Z., Acutis, M., Doro, L., Hidy, D., K chy, M., Minet, J., Lellei-Kov cs, E.,  
1110 Ma, S., Perego, A., Rolinski, S., Ruget, F., Sanna, M., Seddaiu, G., Wu, L., Bellocchi, G., 2017.  
1111 Multi-model simulation of soil temperature, soil water content and biomass in Euro-  
1112 Mediterranean grasslands: Uncertainties and ensemble performance. *European Journal of*  
1113 *Agronomy* 88, 22-40. <https://doi.org/10.1016/j.eja.2016.06.006>

1114 S ndor, R., Picon-Cochard, C., Martin, R., Louault, F., Klumpp, K., Borr s, D., Bellocchi, G.,  
1115 2018. Plant acclimation to temperature: Developments in the Pasture Simulation model. *Field*  
1116 *Crops Research* 222, 238-255. <https://doi.org/10.1016/j.fcr.2017.05.030>

1117 Shen, M., Piao, S., Cong, N., Zhang, G., Janssens, I.A., 2015a. Precipitation impacts on vegetation  
1118 spring phenology on the Tibetan Plateau. *Glob. Change Biol.* 21, 3647-3656.  
1119 <https://doi.org/10.1111/gcb.12961>

1120 Soudani, K., Hmimina, G., Delpierre, N., Pontailleur, J.Y., Aubinet, M., Bonal, D., Caquet, B., de  
1121 Grandcourt, A., Burban, B., Flechard, C., Guyon, D., Granier, A., Gross, P., Heinesh, B.,  
1122 Longdoz, B., Loustau, D., Moureaux, C., Ourcival, J.M., Rambal, S., Saint Andr , L., Dufr ne,  
1123 E., 2012. Ground-based network of NDVI measurements for tracking temporal dynamics of

1124 canopy structure and vegetation phenology in different biomes. *Remote Sensing of Environment*  
1125 123, 234-245. <https://doi.org/10.1016/j.rse.2012.03.012>

1126 Stendardi, L., Dibari, C., Bassignana, M., Bindi, M., Brilli, L., Choler, P., Cremonese, E., Filippa,  
1127 G., Piccot, A., Argenti, G., 2022. Pasture areas in the Gran Paradiso National Park. *Journal of*  
1128 *Maps*. <https://doi.org/10.1080/17445647.2022.2120835>

1129 Targetti, S., Mouléry, M., Napoléone, C., 2019. Agricultural policy, climate change adaptation and  
1130 sustainability: Assessing drivers and mechanisms in an alpine case study area. Contributed paper  
1131 at the 8<sup>th</sup> AIEAA (Italian Association of Agricultural and Applied Economics) Congress, 13-  
1132 14 June, Pistoia, Italy. <https://www.aieaa.org/aieaaconference2019>

1133 Van Oijen, M., Barcza Z., Confalonieri R., Korhonen P., Kröel-Dulay G., Lellei-Kovács E., Louarn  
1134 G., Louault F., Martin R., Moulin T., Moredi E., Picon-Cochard C., Rolinski S., Viovy N.,  
1135 Wirth S.B., Bellocchi, G., 2020. Incorporating biodiversity into biogeochemistry models to  
1136 improve prediction of ecosystem services in temperate grasslands: review and roadmap.  
1137 *Agronomy* 10: 259. <https://doi.org/10.3390/agronomy10020259>

1138 Van Vuuren, D.P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G.C.,  
1139 Kram, T., Krey, V., Lamarque, J.-F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S.J.,  
1140 Rose, S.K., 2011. The representative concentration pathways: an overview. *Climatic Change*  
1141 109, 5-31. <https://doi.org/10.1007/s10584-011-0148-z>

1142 Vital, J.A., Gaurut, M., Lardy, R., Viovy, N., Soussana, J.F., Bellocchi, G., Martin, R., 2013. High-  
1143 performance computing for climate change impact studies with the Pasture Simulation model.  
1144 *Computers and Electronics in Agriculture* 98, 131-135.  
1145 <https://doi.org/10.1016/j.compag.2013.08.004>

1146 Volaire, F., Lelièvre, F., 2001. Drought survival in *Dactylis glomerata* and *Festuca arundinacea*  
1147 under similar rooting conditions. *Plant and Soil* 229, 225-234.  
1148 <https://doi.org/10.1023/A:1004835116453>

1149 Wanner, A., Pröbstl-Haider, U., Feilhammera, M., 2021. The future of Alpine pastures -  
1150 Agricultural or tourism development? Experiences from the German Alps. *Journal of Outdoor*  
1151 *Recreation and Tourism* 35: 100405. <https://doi.org/10.1016/j.jort.2021.100405>

1152 Weitz, A.M., Grauel, W.T., Keller, M., Veldkamp, E., 1997. Calibration of time domain  
1153 reflectometry technique using undisturbed soil samples from humid tropical soils of volcanic  
1154 origin. *Water Resources Research* 33, 1241-1249. <https://doi.org/10.1029/96WR03956>

1155 Wilcke, R.A., Barring L., 2016. Selecting regional climate scenarios for impact modelling studies.  
1156 *Environmental Modelling & Software* 78, 191-201.  
1157 <https://doi.org/10.1016/j.envsoft.2016.01.002>

1158 Wu, W., Geller, M.A., Dickinson, R.E., 2002. A case study for land model evaluation: Simulation  
1159 of soil moisture amplitude damping and phase shift. *Journal of Geophysical Research* 107, ACL  
1160 20-1-ACL 20-13. <https://doi.org/10.1029/2001JD001405>

1161 Xu, C., Liu, H., Williams, A.P., Yin, Y., Wu, X., 2016. Trends toward an earlier peak of the  
1162 growing season in Northern Hemisphere mid-latitudes. *Glob. Chang. Biol.* 22, 2852–2860.  
1163 <https://doi.org/10.1111/gcb.13224>

1164 Zeeman, M.J., Hiller, R., Gilgen, A.K., Michna, P., Plüss, P., Buchmann, N., Eugster, W., 2010.  
1165 Management and climate impacts on net CO<sub>2</sub> fluxes and carbon budgets of three grasslands  
1166 along an elevational gradient in Switzerland. *Agricultural and Forest Meteorology* 150, 519–  
1167 530. <https://doi.org/10.1016/j.agrformet.2010.01.011>

1168 Zeeman, M.J., Shupe, H., Baessler, C., Ruehr, N.K., 2019. Productivity and vegetation structure of  
1169 three differently managed temperate grasslands. *Agriculture, Ecosystems and Environment*  
1170 270–271, 129–148. <https://doi.org/10.1016/j.agee.2018.10.003>