

**Supplementary info 1:** Reference list of the article analyzed for the literature review on compositional analysis of plant protein ingredients

- Alamanou, S. and Doxastakis, G. (1997) ‘Effect of wet extraction methods on the emulsifying and foaming properties of lupin seed protein isolates (*Lupinus albus* ssp. *Graecus*)’, *Food Hydrocolloids*, 11(4), pp. 409–413. doi: 10.1016/S0268-005X(97)80038-0.
- Bader, S., Bez, J. and Eisner, P. (2011) ‘Can protein functionalities be enhanced by high-pressure homogenization? – A study on functional properties of lupin proteins’, *Procedia Food Science*, 1, pp. 1359–1366. doi: 10.1016/j.profoo.2011.09.201.
- Barac, M. *et al.* (2010) ‘Profile and functional properties of seed proteins from six pea (*Pisum sativum*) genotypes’, *International Journal of Molecular Sciences*, 11(12), pp. 4973–4990. doi: 10.3390/ijms11124973.
- Benjamin, O. *et al.* (2014) ‘Emulsifying properties of legume proteins compared to  $\beta$ -lactoglobulin and tween 20 and the volatile release from oil-in-water emulsions’, *Journal of Food Science*, 79(10), pp. E2014–E2022. doi: 10.1111/1750-3841.12593.
- Burger, T. G. *et al.* (2021) ‘Comparison of physicochemical and emulsifying properties of commercial pea protein powders’, *Journal of the Science of Food and Agriculture*, (October). doi: 10.1002/jsfa.11592.
- Burgos-Díaz, C. *et al.* (2019) ‘Food-grade Pickering stabilizers obtained from a protein-rich lupin cultivar (AluProt-CGNA®): Chemical characterization and emulsifying properties’, *Food Hydrocolloids*, 87(September 2018), pp. 847–857. doi: 10.1016/j.foodhyd.2018.09.018.
- Chang, C. *et al.* (2015) ‘Effect of pH on the inter-relationships between the physicochemical, interfacial and emulsifying properties for pea, soy, lentil and canola protein isolates’, *Food Research International*, 77, pp. 360–367. doi: 10.1016/j.foodres.2015.08.012.
- Chen, M. *et al.* (2019) ‘Study on the emulsifying stability and interfacial adsorption of pea proteins’, *Food Hydrocolloids*, 88(September 2018), pp. 247–255. doi: 10.1016/j.foodhyd.2018.09.003.
- Cui, L. *et al.* (2021) ‘Improving the functionality of pea protein isolate through co-spray drying with emulsifying salt or disaccharide’, *Food Hydrocolloids*, 113(November 2020), p. 106534. doi: 10.1016/j.foodhyd.2020.106534.
- Donsì, F. *et al.* (2010) ‘Development of novel pea protein-based nanoemulsions for delivery of nutraceuticals’, *Journal of Agricultural and Food Chemistry*, 58(19), pp. 10653–10660. doi: 10.1021/jf101804g.
- Dumoulin, L. *et al.* (2021) ‘Dry and wet fractionation of plant proteins: How a hybrid process increases yield and impacts nutritional value of faba beans proteins’, *Innovative Food Science and Emerging Technologies*, 72(February), p. 102747. doi: 10.1016/j.ifset.2021.102747.
- Fuhrmeister, H. and Meuser, F. (2003) ‘Impact of processing on functional properties of protein products from wrinkled peas’, *Journal of Food Engineering*, 56(2–3), pp. 119–129. doi: 10.1016/S0260-8774(02)00241-8.

Funke, M., Boom, R. and Weiss, J. (2022) ‘Dry fractionation of lentils by air classification - Composition, interfacial properties and behavior in concentrated O/W emulsions’, *Lwt*, 154, p. 112718. doi: 10.1016/j.lwt.2021.112718.

Geerts, M. E. J. *et al.* (2017) ‘Protein nativity explains emulsifying properties of aqueous extracted protein components from yellow pea’, *Food Structure*, 14(September), pp. 104–111. doi: 10.1016/j.foostr.2017.09.001.

Gharsallaoui, A. *et al.* (2009) ‘Interfacial and emulsifying characteristics of acid-treated pea protein’, *Food Biophysics*, 4(4), pp. 273–280. doi: 10.1007/s11483-009-9125-8.

Gumus, C. E., Decker, E. A. and McClements, D. J. (2017) ‘Formation and Stability of  $\omega$ -3 Oil Emulsion-Based Delivery Systems Using Plant Proteins as Emulsifiers: Lentil, Pea, and Faba Bean Proteins’, *Food Biophysics*, 12(2), pp. 186–197. doi: 10.1007/s11483-017-9475-6.

Hinderink, E. B. A. *et al.* (2021) ‘Physical and oxidative stability of food emulsions prepared with pea protein fractions’, *Lwt*, 146(November 2020). doi: 10.1016/j.lwt.2021.111424.

Karaca, A. C., Low, N. and Nickerson, M. (2011) ‘Emulsifying properties of chickpea, faba bean, lentil and pea proteins produced by isoelectric precipitation and salt extraction’, *Food Research International*, 44(9), pp. 2742–2750. doi: 10.1016/j.foodres.2011.06.012.

Kutzli, I. *et al.* (2020) ‘Improvement of emulsifying behavior of pea proteins as plant-based emulsifiers: Via Maillard-induced glycation in electrospun pea protein-maltodextrin fibers’, *Food and Function*, 11(5), pp. 4049–4056. doi: 10.1039/d0fo00292e.

Lqari, H. *et al.* (2002) ‘Lupinus angustifolius protein isolates: Chemical composition, functional properties and protein characterization’, *Food Chemistry*, 76(3), pp. 349–356. doi: 10.1016/S0308-8146(01)00285-0.

Ma, K. K. *et al.* (2022) ‘Functional Performance of Plant Proteins’, *Foods*, 11(4), pp. 1–23. doi: 10.3390/foods11040594.

Möller, A. C., van der Padt, A. and van der Goot, A. J. (2021) ‘From raw material to mildly refined ingredient – Linking structure to composition to understand fractionation processes’, *Journal of Food Engineering*, 291. doi: 10.1016/j.jfoodeng.2020.110321.

Ntone, E. *et al.* (2021) ‘Adsorption of rapeseed proteins at oil/water interfaces. Janus-like napins dominate the interface’, *Journal of Colloid and Interface Science*, 583, pp. 459–469. doi: 10.1016/j.jcis.2020.09.039.

Pelgrom, P. J. M., Boom, R. M. and Schutyser, M. A. I. (2015) ‘Functional analysis of mildly refined fractions from yellow pea’, *Food Hydrocolloids*, 44, pp. 12–22. doi: 10.1016/j.foodhyd.2014.09.001.

Reinkensmeier, A. *et al.* (2015) ‘Characterization of individual proteins in pea protein isolates and air classified samples’, *Food Research International*, 76(P1), pp. 160–167. doi: 10.1016/j.foodres.2015.05.009.

Rempel, C., Geng, X. and Zhang, Y. (2019) ‘Industrial scale preparation of pea flour fractions with enhanced nutritive composition by dry fractionation’, *Food Chemistry*, 276(October 2018), pp. 119–128. doi: 10.1016/j.foodchem.2018.10.003.

Solaesa, Á. G. *et al.* (2020) ‘Protein and lipid enrichment of quinoa (cv.Titicaca) by dry fractionation. Techno-functional, thermal and rheological properties of milling fractions’, *Food Hydrocolloids*, 105(February). doi: 10.1016/j.foodhyd.2020.105770.

Sridharan, S. *et al.* (2020) ‘On the Emulsifying Properties of Self-Assembled Pea Protein Particles’, *Langmuir*, 36(41), pp. 12221–12229. doi: 10.1021/acs.langmuir.0c01955.

Tabatabaei, S. *et al.* (2019) ‘Functional properties of navy bean (*Phaseolus vulgaris*) protein concentrates obtained by pneumatic tribo-electrostatic separation’, *Food Chemistry*, 283(October 2018), pp. 101–110. doi: 10.1016/j.foodchem.2019.01.031.

Tsoukala, A. *et al.* (2006) ‘Adsorption at the air-water interface and emulsification properties of grain legume protein derivatives from pea and broad bean’, *Colloids and Surfaces B: Biointerfaces*, 53(2), pp. 203–208. doi: 10.1016/j.colsurfb.2006.08.019.

Vogelsang-O’Dwyer, M., Zannini, E. and Arendt, E. K. (2021) ‘Production of pulse protein ingredients and their application in plant-based milk alternatives’, *Trends in Food Science and Technology*, 110(January), pp. 364–374. doi: 10.1016/j.tifs.2021.01.090.

Wang, J. *et al.* (2016) ‘Lupine protein enrichment by milling and electrostatic separation’, *Innovative Food Science and Emerging Technologies*, 33, pp. 596–602. doi: 10.1016/j.ifset.2015.12.020.

Yang, J. *et al.* (2021) ‘Foams and air-water interfaces stabilised by mildly purified rapeseed proteins after defatting’, *Food Hydrocolloids*, 112(June 2020), p. 106270. doi: 10.1016/j.foodhyd.2020.106270.