



HAL
open science

Fate of polychlorobiphenyls in *Tenebrio molitor* larvae: consequences for further use as food and feed

J. Ratel, F. Mercier, J. Rivas, H. Wang, M. Angénieux, B. Calmont, S.
Crépieux, Christelle Planche, E. Engel

► **To cite this version:**

J. Ratel, F. Mercier, J. Rivas, H. Wang, M. Angénieux, et al.. Fate of polychlorobiphenyls in *Tenebrio molitor* larvae: consequences for further use as food and feed. *Journal of Insects as Food and Feed*, 2023, 9 (6), pp.781-788. 10.3920/JIFF2022.0104 . hal-04092954

HAL Id: hal-04092954

<https://hal.inrae.fr/hal-04092954v1>

Submitted on 9 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Fate of Polychlorobiphenyls in *Tenebrio molitor* larvae**Fate of polychlorobiphenyls in *Tenebrio molitor* larvae: consequences for further use as food and feed**

J. Ratel¹, F. Mercier¹, J. Rivas¹, H. Wang¹, M. Angénioux¹, B. Calmont², S. Crépieux³, C. Planche^{1*}, E. Engel¹

¹ MASS group, UR QuaPA, INRAE, F-63122 Saint-Genès-Champanelle, France

² SHNAO, Société d'histoire naturelle Alcide-d'Orbigny, F-63170 Aubière, France

³ Invers, F-63720 Saint-Ignat, France

* Corresponding Author:

Christelle Planche: Email: christelle.planche@inrae.fr Tel: +33 4 73624514

<https://orcid.org/0000-0002-7751-4603>

RESEARCH ARTICLE**Abstract**

This study explored the ability of *Tenebrio molitor*, one of the most widely bred and traded insect species in Europe, to bioaccumulate Polychlorobiphenyls (PCBs) from their feeding substrate. *T. molitor* larvae were reared for 20 days in a temperature and humidity-controlled incubator and fed with wheat bran artificially contaminated with PCBs at a concentration of 0.67, 4 or 24.4 ppb. The larvae PCB content was then measured based on an analysis by GC-MS and GC- μ ECD. Whatever the level, a bran contamination by PCBs did not affect the body weight of larvae indicating a high tolerance to PCBs. The bioaccumulation factors (BAF = concentration of PCBs in larvae / concentration of PCBs in wheat bran) obtained with fresh larvae ranged between 0.4 and 1.0 and were significantly higher with the highest contamination level for PCBs 153 and 180. Although there is no bioaccumulation of PCBs in *T. molitor* larvae during rearing, PCBs can be transferred from the diet to the larvae. This study highlights the significant impact of the drying process which induces an increase of the PCB concentration in larvae by a factor of almost 3. This demonstrates the importance of considering the quality of the substrates used for farming insects as food and feed in terms of content in chemical contaminants including persistent organic pollutants like PCBs.

Keywords

PCB, yellow mealworm, bioaccumulation, contaminants, safety

Conflict of interest

The authors declare no conflict of interest.

Funding statement

This study has been implemented in the R&D Booster program funded by the Auvergne Rhône Alpes Region (INSECT2FEED project).

1. Introduction

The United Nations estimate that the global human population is expected to reach 9.7 billion in 2050 increasing the consumption of animal products and the growing demand for feed ingredients (United Nations, 2019). Available resources are currently limited and may raise environmental, societal, and economic questions (Tran *et al.*, 2015). The quest for alternative sustainable animal protein sources is therefore expected to become a considerable issue in the feed market. Edible insects could be an interesting solution with a high nutritional value (Van Huis, 2021). Processed animal protein derived from insects are currently authorised for aquafeeds (Commission Regulation (EU) 2017/893) (EU, 2017), poultry and swine feeds in the EU (Commission Regulation (EU) 2021/1372) (EU, 2021). Among the broad variety of insect species, the yellow mealworm (*Tenebrio molitor*) is one of the most widely bred and traded insect species in Europe (Bordiean *et al.*, 2020).

To develop this new sector in future years, it is essential to assess and control the risks associated with the use of insects. Some studies are already available about the microbiological hazards (Vandeweyer *et al.*, 2021). However, there is a lack of knowledge concerning the chemical safety while some chemical contaminants may be found in insect farming environment or in their feeding substrates, can be produced during processing methods or can migrate from packaging material to insect products (Meyer *et al.*, 2021). While some authors have reported that insects may degrade some mycotoxins (Niermans *et al.*, 2019), some antimicrobials (Cai *et al.*, 2018) or even some pesticides (De Almeida *et al.*, 2017), other chemical contaminants like heavy metals can have a negative impact on insect development inducing, for example, an increase in larval mortality (Bulak *et al.*, 2018). Moreover, previous studies have shown that insects like *T. molitor* can bioaccumulate toxic trace elements during rearing, with very variable bioaccumulation factors up to 1.7 for cadmium, 6.2 for mercury and even 34 for lead (Truzzi *et al.*, 2019).

Among chemical contaminants, persistent organic pollutants (POPs) can be transferred from environment or feed to animal products and therefore are of serious concern in animal-derived food products (Engel *et al.*, 2015). In the black soldier fly (*Hermetia illucens*), Van der Fels-Klerx *et al.* (2020) have determined that the bioaccumulation factors of POPs from their substrates (different whole meals and different types of snack products, both with paperboard carton and with plastic packaging materials) ranged between 0.3 and 1.2 for polycyclic aromatic hydrocarbon (PAHs) and between 1.0 and 2.0 for the sum of dioxins and dioxin-like polychlorinated biphenyls (PCBs). However, to our knowledge, there are no studies regarding the bioaccumulation of POPs in *T. molitor*.

Using *T. molitor* larvae as a model and setting up a pilot experimental farming under controlled and safe conditions allowing the implementation of an exposure study with chemical contaminants, the present work explores the ability of insects to bioaccumulate PCBs from their feeding substrate during the larvae growth. For this purpose, the first part of this paper is focused on the set up of an analytical method to monitor PCBs in *T. molitor* larvae and the assessment of the performance of this method in terms of linearity (coefficients of determination, R^2) and sensitivity (limits of detection, LOD). In a second part, the impact of a feed contamination by a mix of 6 non-dioxin like PCBs (nDL-PCBs) on the growth of larvae was studied. The bioaccumulation factors of PCBs in larvae were determined at the end of the rearing and will be discussed in the context of a further use of larvae as food and feed.

2. Materials and Methods

Chemicals

91 Hexane, dichloromethane, acetone and toluene were organic trace analysis grade solvents
92 (Sigma-Aldrich, Saint-Quentin Fallavier, France). For PCB extraction, Florisil® and activated
93 aluminium oxide (acidic, Brockmann I) were from Sigma-Aldrich. Diatomaceous earth was
94 obtained from Thermo Fisher Scientific (Waltham, MA, USA). The certified reference material
95 AE-00059-H-2X of AccuStandard Europe (Niederbipp, Switzerland) was used for the
96 development of PCB analysis. This mix is composed of the 6 nDL-PCBs (IUPAC numbers 28,
97 52, 101, 138, 153, and 180) with individual concentrations of 20 ppm. Fluorinated internal
98 standards 3-F-PCB-52 and 5'-F-PCB-126 (Chiron, Trondheim, Norway) were used for the
99 accurate quantification of target compounds.

100 Larvae

101 Feeding

102 Larval feed was dry wheat bran produced by local flour mills in Puy de Dome (France). Larvae
103 were fed with non-contaminated feeds (control group) or with PCB contaminated feeds
104 (exposed groups). For exposed groups, the bran was spiked according to a protocol adapted
105 from Planche *et al.* (2015), based on a contaminant addition to the feed via a volatile solvent.
106 For each PCB exposure condition, the spiking was carried out on the total quantity of wheat
107 bran necessary for all the individual rearing (45g), allowing to ensure the quality of replicates.
108 Briefly, 45 g of dry wheat bran were set in a glass jar (9.5cm high×7cm wide) then immersed
109 in acetone (125 mL) containing the 6 nDL-PCBs. The mixture was evaporated down under a
110 hood and roughly homogenised. Three spiking concentrations, 0.67, 4.0 and 24.4 ppb (ng
111 PCB/g wheat bran), were tested for each PCB congener. This corresponds to levels of $\Sigma(6$ nDL-
112 PCBs) that are 0.4, 2.4 and 14.6 times the maximum level (ML) set at 10 ppb by the Regulation
113 (EU) No 277/2012 in feed materials of plant origin. The spiking concentrations are consistent
114 both with the regulation issues and with the scarce literature data dealing with PCB
115 contamination in food of plant origin, like wheat bran, with for example some concentrations
116 of PCBs in bran that can reach 1.84 ppb (Roszko *et al.*, 2014). For the control groups, the bran
117 was either unspiked (bran without solvent or PCB) or spiked with neat solvent (bran with
118 solvent but without PCB).

119 Rearing

120 Larvae of *T. molitor* were produced by the startup INVERS (Saint-Ignat, France). Using a
121 balance (Precisa 410 AM-FR, Precisa Instruments Ltd., Switzerland; d: 0.0001), 10g of larvae
122 at L3 growth stage (approximately 200 individuals) were set on a glass jar containing 45 g of
123 control or contaminated wheat bran. Three glass jars for each modality were prepared and put
124 in an individual incubator (18 L Heratherm™ Compact Microbiological Incubator, Thermo
125 Scientific) customised for sufficient ventilation (300 L/h) (Figure 1). In total, 15 rearing glass
126 jars were distributed in the 5 incubators. These replicates were not randomly distributed over
127 the incubators to avoid a cross contamination between the different exposure conditions. The
128 humidity was maintained at 60% and the temperature regulated at 26°C with a monitoring
129 throughout the rearing (probe: VAISALA HMP110; data acquisition system: AOIP SA32)
130 (Supplementary Figure S1). A water spray was added to each jar containing larvae two times
131 per day. The rearing of larvae was carried out for 20 days in incubators. From the 6th day of
132 the test, the evaluation of larval weight was performed for each trial twice per week. Larvae
133 were separated by using a 1.5 mm mesh sieve. After sieving, the weight of 20 larvae was
134 determined. After each weighting, the sieved content and the larvae were put back in the jar.

135 Figure 1: Pilot experimental farming of *Tenebrio molitor* larvae.

136

137 Analysis of PCBs**138 Sample preparation**

139 At the end of the rearing period, all insects were sacrificed by freezing at -20°C. Larvae were
140 then microwave-dried according to the usual conditions carried out by INVERS company: the
141 power of the microwave was fixed (600 W) and the duration of the drying adapted so that the
142 final relative humidity of the larvae was inferior to 10% (between 5 and 7 min). Prior to
143 extraction, all samples were powdered with a liquid nitrogen grinder. Extraction process was
144 carried out according to Planche *et al.* (2017) with slight modifications. Briefly, 1 g of larvae
145 powder was extracted by accelerated solvent extraction (ASE) using a Dionex ASE 350
146 extractor (Sunnyvale, CA). Stainless-steel extraction cells (22 mL volume) were used, with 5 g
147 of acidic alumina and 5 g Florisil® placed at the bottom of the cells. Paper filters were placed
148 at the bottom and top of the alumina layer. The cells were then filled with 1 g of ground larvae
149 dispersed in diatomaceous earth and hexane was used as extraction solvent at a temperature of
150 100 °C and pressure of 1500 psi. ASE extraction included heating (5 min), static time (5 min)
151 and purging (90 s) with two extraction cycles per sample. After filtration through a 1.2 µm glass
152 fiber prefilter and a 0.45 µm nylon filter (Phenomenex, Torrance, CA), the extract was
153 evaporated (Rocket; Genevac Ltd, Ipswich, UK) using toluene as a keeper to minimise analyte
154 losses during the evaporation step, then 4 mL of dichloromethane were added. To clean up
155 extracts, gel permeation chromatography (GPC) (Gilson, Middleton, WI) was carried out on an
156 S-X3 Bio-Beads column (Bio-Rad, Philadelphia, PA) using dichloromethane as eluting solvent
157 at a flow rate of 5 mL/min. The fraction collected between 15 and 37 min was evaporated to
158 dryness (Rocket, Genevac Ltd), and a mix of 100 µL of hexane and internal standard was added
159 (Figure 2).

160 Figure 2: Simplified diagram of the workflow used to analyse PCBs in *Tenebrio molitor*
161 larvae**162 PCB detection by GC-ToF/MS**

163 For PCB identification in insect matrix, larvae extracts (obtained according to the conditions
164 detailed in the previous section) were spiked with nDL-PCBs then analyzed with a time-of-
165 flight mass spectrometer (Pegasus 4D, Leco) coupled to a gas chromatograph (6890, Agilent
166 Technologies). 1 µL of extract was injected at 280°C in splitless mode into a DB-5MS capillary
167 column (60 m × 0.32 mm × 1 µm; Agilent J&W) with Helium (purity of 99.99995%) as carrier
168 gas at 1 mL/min. Oven temperature was held at 120°C for 1 min, then ramped up to 240°C at a
169 gradient of 20°C/min and to 300°C at a gradient of 2°C/min, and held at 300°C for 15 min. The
170 MS-temperatures were set at 230 °C, 150 °C and 180 °C in the transfer line, the source and the
171 quadrupole, respectively. Electron impact energy was set at 70 eV, and data was collected in
172 full scan in the range of 45 to 600 m/z at a scan range of 10 scans per second.

173 PCB quantification by GC-µECD

174 The PCB levels in the larvae experimentally reared were determined by µECD (Agilent)
175 coupled to a gas chromatograph (6890, Agilent Technologies). The GC parameters were the
176 same as for the GC-MS coupling. The micro-ECD system was operated at 300 °C using data
177

178 acquisition rate of 50 Hz, with nitrogen as make-up gas at a flowrate of 40 ml/min. Calibration
 179 curves were preliminary built from dried larvae extracts spiked with the 6 nDL-PCBs (see Table
 180 1) at the following concentrations: 0.2, 0.5, 1.3, 3.0, 8.0, 20.0, 50.0 ng/g. Larvae extracts
 181 without PCB added were run in triplicate to check the absence of targeted analytes. Each
 182 concentration level was analysed 4 times by GC- μ ECD. Peak areas were determined,
 183 normalized by internal standard, then used for calculations of the standard deviations and the
 184 calibration curve equations. The linearity of the calibration curves was assessed for each PCB
 185 by calculating the coefficients of determination (R^2). The limit of detection (LOD), using the
 186 definition $3s/m$ (s is the standard deviation of the intercept, and m is the slope of the linear
 187 calibration curve), was determined from the calibration curves for each individual PCB studied.

188 Data treatment

189 Data were processed with the TIBCO's Statistica software (version 13.0, TIBCO Software Inc.)
 190 and the R software (version 3.5.1, The R Foundation for Statistical Computing). The “lm”
 191 (linear model) function of R was used on the calibration curve data for the determination of R^2
 192 as well as s and m requested for LOD calculation. Student's t test or one-way analyses of
 193 variance (ANOVA) followed by *post hoc* Newman–Keuls test were performed to compare
 194 means data. The differences were considered significant when $p < 0.05$.

195 3. Results

196 PCB analysis

197
 198
 199
 200 The recovery rates of the 6 nDL-PCBs measured after spiking in dried powder of *T.*
 201 *molitor* (at 200 ng/g dry matter), extraction, purification and concentration are presented in
 202 Table 1. All the recovery rates lay in the classically accepted range of 70–130% according to
 203 the EPA Method 8000C (2003), with RSD between 10% and 18%.

204
 205 **Table 1: Recovery rates (%) obtained after spiking, extraction, purification,**
 206 **concentration and analysis of PCBs in dried powder of *Tenebrio molitor* larvae ($n=3$).**
 207

208	209	210	211
Compound name	PCB congener	m/z	Recovery rates \pm RSD ¹ (%)
2,4,4'-Trichlorobiphenyl	28	256	108 \pm 17
2,2',5,5'-Tetrachlorobiphenyl	52	292	110 \pm 16
2,2',4,5,5'-Pentachlorobiphenyl	101	326	98 \pm 18
2,2',3,4,4',5'-Hexachlorobiphenyl	138	360	107 \pm 14
2,2',4,4',5,5'-Hexachlorobiphenyl	153	360	108 \pm 16
2,2',3,4,4',5,5'-Heptachlorobiphenyl	180	394	129 \pm 10

¹ Relative standard deviation (%)

The LOD measured with GC- μ ECD for the 6 nDL-PCB congeners in dried powder of *T. molitor* larvae varied from 1.67 ng/g for PCB 153 to 2.98 ng/g for PCB 101, with limits of quantification (LOQ) from 5.56 to 9.93 ng/g respectively (Supplementary Table S1). Coefficients of determination ranged between 0.94 for PCB 101 and 0.98 for PCBs 138, 153 and 180.

Impact of feed contamination with PCBs on larval growth

Table 2 and Figure S2 present the weight of *T. molitor* larvae at days 1, 7, 10, 14, 17 and 20 of rearing depending on their feeding substrate: wheat bran unspiked, spiked with neat solvent without PCB or spiked with nDL-PCBs at 0.67, 4.0 or 24.4 ng/g corresponding to 0.4, 2.4 and 14.6 times the maximum level (ML) for the nDL-PCBs in feed materials of plant origin, respectively (Commission regulation (EU) No 277/2012). One-way ANOVA ($p < 0.05$) revealed no significant difference between the different conditions with a mean weight of 54.1 mg per larvae at the beginning of the study and a continuous growth throughout the rearing up to 95.6 mg per larvae on average after 20 days of rearing. Visual observations throughout the rearing period did not reveal differences in terms of larval mortality between the different conditions.

Table 2: Weight of *Tenebrio molitor* larvae (mg) throughout their rearing on wheat bran either unspiked (Control), spiked with neat solvent (Control + acetone) or spiked at 0.67, 4.0 or 24.4 ng PCB/g wheat bran. Data represent mean larvae weight (mg) \pm SD which is determined from the 20 larvae measurement for each rearing glass jar ($n=3$ glass jars for each exposure condition).

Day	Control	Control + acetone	0.67 ppb	4.00 ppb	24.40 ppb	Significance ¹
1	54.5 \pm 2.8	50.2 \pm 1.8	54.6 \pm 1.5	56.5 \pm 4.5	54.6 \pm 6.8	NS
7	67.1 \pm 3.5	67.2 \pm 9.2	64.1 \pm 3.3	65.3 \pm 3.8	73.0 \pm 6.4	NS
10	85.6 \pm 5.3	76.5 \pm 3.3	75.4 \pm 5.4	80.7 \pm 5.4	82.1 \pm 14.4	NS
14	83.4 \pm 4.9	81.3 \pm 0.8	85.1 \pm 3.3	88.9 \pm 7.0	86.0 \pm 2.5	NS
17	94.1 \pm 5.0	92.9 \pm 5.8	96.8 \pm 3.0	86.7 \pm 2.1	93.7 \pm 9.4	NS
20	93.1 \pm 7.2	92.2 \pm 8.6	97.4 \pm 5.8	97.2 \pm 4.7	98.0 \pm 2.4	NS

NS: not significant ($p > 0.05$)

¹ p values were calculated for one-way analyses of variance (ANOVA) followed by *post hoc* Newman–Keuls test.

Bioaccumulation factors

Bioaccumulation factors (BAF) were calculated for each congener of PCB in fresh (Table 3) and dried (Table S4) *T. molitor* larvae according to Truzzi *et al.* (2019) and Van der Fels-

242 Klerx *et al.* (2020), with BAF = concentration of PCB in the organism (fresh or dried larvae) /
 1 243 concentration of PCB in the feed provided (dried wheat bran).

2 244 For a spiking concentration of 0.67 ng PCB/g wheat bran, concentrations in dried larvae
 3 245 extracts of the 6 nDL-PCBs were below the detection limits detailed in Supplementary Table
 4 246 S1. For a spiking concentration of 4 ng PCB/g wheat bran, the concentrations of PCB 28, 52
 5 247 and 101 were between LOD and LOQ (Supplementary Table S2). For all other conditions, PCB
 6 248 congener concentrations are above the LOQ. Regarding these last values, Supplementary Table
 7 249 S4 shows that BAF obtained in dried larvae ranged from 1.0 to 2.8 (raw data enabling BAF
 8 249 calculation given in Table S3). For PCB 138, there is no significant difference in BAF obtained
 9 250 between the two highest spiking levels whereas for PCBs 153 and 180, BAF significantly
 10 251 increased with the spiking level. Moreover, BAF obtained for PCB 52 are significantly lower
 11 252 than for other congeners.
 12 253

13 254 Table 3 shows the BAF of the 6 nDL-PCBs congeners in fresh larvae. BAF in fresh larvae
 14 255 are 2.8 times lower than in dried larvae.
 15 256

16 256
 17 256
 18
 19 257 **Table 3: Bioaccumulation factors (BAF) of PCB congeners in *Tenebrio molitor* larvae on**
 20 258 **a fresh weight basis after 20 days of rearing on wheat bran spiked at 0.67, 4.0 or 24.4 ng**
 21 259 **PCB/g.** These values were estimated from data obtained on dried larvae, taking into account
 22 259 the impact of drying on larval weight. Data represent mean BAF \pm SD which is determined
 23 260 from 3 analytical replicates of each rearing glass jar ($n=3$ glass jars for each exposure
 24 261 condition).
 25 262
 26 263

Congener	Wheat bran contamination with PCBs		
	0.67 ppb	4.0 ppb	24.4 ppb
PCB 28	NA	0.80 \pm 0.17 ^{a*}	0.68 \pm 0.03 ^b
PCB 52	NA	0.39 \pm 0.07 ^{a*}	0.36 \pm 0.03 ^a
PCB 101	NA	0.66 \pm 0.10 ^{a*}	0.67 \pm 0.02 ^a
PCB 138	NA	0.86 \pm 0.16 ^a	0.96 \pm 0.05 ^a
PCB 153	NA	0.83 \pm 0.13 ^a	0.99 \pm 0.05 ^b
PCB 180	NA	0.81 \pm 0.12 ^a	0.97 \pm 0.08 ^b

27 264 NA: non-available (PCBs non-detected in larvae extracts)

28 265 a–b: different superscript letters within the same row indicate significant differences among values
 29 266 ($p<0.05$). BAF are determined from quantification values in larvae extracts either between LOD and
 30 267 LOQ (*) or greater than LOQ.
 31 268
 32 269
 33 270

34 271 4. Discussion

35 272
 36 273 In order to assess the fate of the 6 nDL-PCBs in *T. molitor* larvae during rearing with a
 37 274 contaminated substrate, the first step of this study was to determine their recovery rates after
 38 275 the spiking, extraction, purification, concentration and analysis protocol. Based on a protocol
 39 276 adapted from Planche *et al.* (2015) with a delipidation strengthened by the addition of fat
 40 277 retainers, the PCB recoveries in larvae samples (from 98% to 129%) are of the same order of
 41 278 magnitude as those in raw beef samples (from 87% to 133%) (Planche *et al.*, 2015). This
 42 279 demonstrates that the high fat level of *T. molitor* larvae (56–61% in whole dried *T. molitor*
 43 280 (EFSA, 2021) vs 11% in raw ground beef) does not decrease the recovery rates obtained even
 44 280
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65

281 if PCBs are hydrophobic compounds (Log Kow = 4.09–8.18 according to Hawker and Connell
282 (1988)), thereby confirming the relevance of the protocol used.

283 Based on this protocol, the impact of a contamination by PCBs on *T. molitor* was assessed
284 using a pilot experimental farming where larvae were fed for 20 days with wheat bran either
285 unspiked, spiked with neat solvent without PCB or spiked with nDL-PCBs at 0.67, 4.0 or 24.4
286 ng g⁻¹. No significant differences were observed in terms of larval growth between the group
287 fed with unspiked wheat bran and the group fed with wheat bran spiked with neat solvent
288 suggesting that the spiking step had no impact on the further dietary intake of larvae. This
289 confirms that the use of acetone as an inert PCB “vehicle” volatile solvent is relevant contrary,
290 for example, to a mix of chloroform and methanol which has led to a lower *T. molitor* body
291 weight when it was used as a vehicle to spike their feed with Aflatoxin B1 (Bosch *et al.*, 2017).

292 Moreover, a high PCB concentration in feed has no more impact than a low concentration
293 on the larvae body weight of *T. molitor* contrary to previous results obtained with antimicrobials
294 by Gao *et al.* (2019). These authors showed that the weight of *H. illucens* larvae declined
295 gradually with the increase in sulfonamide concentration from 0 to 10 mg/kg feed (Gao *et al.*,
296 2019). The present study indicates that a contamination of *T. molitor* by PCBs in an insect farm
297 could not be revealed by monitoring the larvae weight, suggesting that a systematic control of
298 the quality of insect feed substrates used is essential. It would be interesting to determine if
299 these results would be similar with dioxin-like PCBs that have a different mechanism of action
300 than nDL-PCBs on cellular pathways (Elnar *et al.*, 2012).

301 In order to determine the capacity of the 6 nDL-PCBs to be transferred from feed to *T.*
302 *molitor* larvae, bioaccumulation factors (BAF) were determined for each PCB congener. If we
303 take into consideration BAF determined from quantification values in larvae extracts greater
304 than LOQ, only BAF results for PCBs 138, 153 and 180 can be discussed for a spiking
305 concentration of 4 ng PCB/g wheat bran (Tables 3 and S4). For PCBs 153 and 180, BAF related
306 to the highest spiking concentration (24.4 ng PCB/g wheat bran) are significantly higher
307 compared to lower spiking concentration (4.0 ng PCB/g wheat bran), suggesting that, for these
308 congeners, BAF increases with the contamination level.

309 Knowing that a BAF greater than 1 suggests a bioaccumulation of PCBs from wheat bran
310 into *T. molitor* larvae, the BAF obtained with fresh larvae (from 0.4 to 1.0; Table 3) indicate
311 that there is no bioaccumulation of PCBs in larvae. However, PCBs are transferred from the
312 diet to the larvae during rearing and are then concentrated in larvae during the drying process.
313 This concentration effect explains why the BAF calculated on the basis of dried larvae are 2.8
314 times higher (from 1.0 to 2.8; Supplementary Table S4).

315 Regarding previous data on nDL-PCBs, Van der Fels-Klerx *et al.* (2020) reported BAF in
316 *H. illucens* dried larvae from 0.8 to 1.2 (based on upper bound values) after rearing on meat or
317 vegetarian substrate containing 3-6% of plastic or paperboard carton packaging material. Their
318 lower values of BAF compared to those obtained in the present study may be explained by the
319 differences in terms of insect species and feeding substrate used. Moreover, the starvation
320 period of 24h set up by Van der Fels-Klerx *et al.* (2020) before harvesting may allow to renew
321 the digestive tract content and thus may decrease the concentration of nDL-PCBs in larvae and
322 therefore the BAF.

323 PCBs are hydrophobic compounds (Log Kow = 4.09–8.18 according to Hawker and
324 Connell (1988)) which implies that they may be transferred in the lipid fraction of larvae during
325 rearing. Since the drying process does not impact this lipid fraction, it leads to a concentration
326 of PCBs in dried larvae and therefore an increase of BAF. When a PCB contamination is
327 detected in larvae feed substrate, it could therefore be envisaged not to use the whole dried
328 larvae for food and feed but to exploit separately the protein and lipid fractions. In order to limit
329 the risks linked to the presence of chemical contaminants, the lipid fraction (where PCBs are

330 accumulated) could then be used for non-food applications (biofuel production for example)
331 while the protein fraction could possibly be used for food and feed.

332 333 **5. Conclusion**

334
335 The pilot experimental farming set up enabled to assess the bioaccumulation of PCBs in *T.*
336 *molitor* larvae during rearing. A feed substrate contamination with PCBs doesn't seem to have
337 any significant impact on larval development without negative effects on growth. It could be
338 supposed that a much higher PCB level than those assessed in this study could have a
339 significant impact on the development of *T. molitor*, but these concentrations would be
340 unrealistic in insect feed substrate. In addition, it would be interesting to assess the impact on
341 larval development of a longer rearing period on a contaminated substrate. The present study
342 revealed that there is no bioaccumulation of PCBs in *T. molitor* larvae during rearing.
343 However, PCBs can be transferred from the diet to the larvae and are then concentrated in
344 larvae during the drying process. This highlights the significant impact of the drying process
345 which induces a concentration by a factor of almost 3 of PCBs in larvae. Knowing that the
346 nDL-PCB concentration maximum limit is set at 10 µg/kg either for feed materials of plant
347 origin and for feed materials of land animal origin (Commission regulation, EU No 277/2012),
348 the use as larvae feeding substrate of wheat bran containing a residual amount of nDL-PCBs
349 could result, after the rearing period, in dried larvae with an almost 3-fold higher nDL-PCB
350 concentration that could potentially exceed the maximum limit for its use as feed. This
351 highlights the importance of monitoring the quality of the feed substrates used for farming
352 insects for food and feed in order to guarantee a safe content in chemical contaminants
353 including persistent organic pollutants like PCBs. When a PCB contamination is detected in
354 larvae feed substrate, it could therefore be envisaged to use the larvae lipid fraction (where
355 PCBs are accumulated) for non-food applications. Finally, in order to ensure food safety
356 throughout the food chain, it would be interesting to assess the PCB biomagnification factors
357 if *T. molitor* larvae are used as feed, for example for aquaculture since it is expected that most
358 insect production will be targeted towards aquafeed by 2030 (Van Huis, 2022).

359 360 **Acknowledgements**

361 This study has been implemented in the R&D Booster program funded by the Auvergne Rhône
362 Alpes Region (INSECT2FEED project).

363 364 **References**

- 365
366 Bordiean, A., Krzyżaniak, M., Stolarski, M.J., Czachorowski, S. and Peni, D., 2020. Will
367 yellow mealworm become a source of safe proteins for Europe? Agriculture 10: 233.
368 <https://doi.org/10.3390/agriculture10060233>
369 Bosch, G., Fels-Klerx, H., Rijk, T. and Oonincx, D., 2017. Aflatoxin B1 tolerance and
370 accumulation in black soldier fly larvae (*Hermetia illucens*) and yellow mealworms
371 (*Tenebrio molitor*). Toxins 9(6): 185. <https://doi.org/10.3390/toxins9060185>
372 Bulak, P., Polakowski, C., Nowak, K., Waśko, A., Wiącek, D. and Bieganski, A., 2018.
373 *Hermetia illucens* as a new and promising species for use in entomoremediation.
374 Science of the Total Environment 633: 912-919.
375 <https://doi.org/10.1016/j.scitotenv.2018.03.252>
376 Cai, M., Ma, S., Hu, R., Tomberlin, J.K., Yu, C., Huang, Y., Zhan, S., Li, W., Zheng, L., Yu,
377 Z. and Zhang, J., 2018. Systematic characterization and proposed pathway of

- 378 tetracycline degradation in solid waste treatment by *Hermetia illucens* with intestinal
 1 379 microbiota. Environmental Pollution 242: 634-642.
 2 380 <https://doi.org/10.1016/j.envpol.2018.06.105>
 3 381 De Almeida, L.G., De Moraes, L.A.B., Trigo, J.R., Omoto, C. and Cônsoli, F.L., 2017. The
 4 382 gut microbiota of insecticide-resistant insects houses insecticide-degrading bacteria: a
 5 383 potential source for biotechnological exploitation. PLOS ONE 12(3): e0174754.
 6 384 <https://doi.org/10.1371/journal.pone.0174754>
 7 385 EFSA Panel on Nutrition, Novel Foods and Food Allergens (NDA), Turck, D., Castenmiller,
 8 386 J., De Henauw, S., Hirsch- Ernst, K.I., Kearney, J., Maciuk, A., Mangelsdorf, I.,
 9 387 McArdle, H.J., Naska, A., Pelaez, C., Pentieva, K., Siani, A., Thies, F., Tsabouri, S.,
 10 388 Vinceti, M., Cubadda, F., Frenzel, T., Heinonen, M., Marchelli, R., Neuhäuser-
 11 389 Berthold, M., Poulsen, M., Prieto Maradona, M., Schlatter, J.R., Van Loveren, H.,
 12 390 Ververis, E. and Knutsen, H.K., 2021. Safety of dried yellow mealworm (*Tenebrio*
 13 391 *molitor* larva) as a novel food pursuant to Regulation (EU) 2015/2283. EFSA Journal
 14 392 19. <https://doi.org/10.2903/j.efsa.2021.6343>
 15 393 Elnar, A.A., Diesel, B., Desor, F., Feidt, C., Bouayed, J., Kiemer, A.K. and Soulimani, R.,
 16 394 2012. Neurodevelopmental and behavioral toxicity via lactational exposure to the sum
 17 395 of six indicator non-dioxin-like-polychlorinated biphenyls (Σ 6 ND-L-PCBs) in mice.
 18 396 Toxicology 299: 44-54. <https://doi.org/10.1016/j.tox.2012.05.004>
 19 397 Engel, E., Ratel, J., Bouhrel, J., Planche, C. and Meurillon, M., 2015. Novel approaches to
 20 398 improving the chemical safety of the meat chain towards toxicants. Meat Science 109:
 21 399 75-85. <https://doi.org/10.1016/j.meatsci.2015.05.016>
 22 400 EU, 2021. Commission Regulation (EU) 2021/1372 of 17 August 2021 amending Annex IV
 23 401 to Regulation (EC) No 999/2001 of the European Parliament and of the Council as
 24 402 regards the prohibition to feed non-ruminant farmed animals, other than fur animals,
 25 403 with protein derived from animals.
 26 404 EU, 2017. Commission Regulation (EU) 2017/893 of 24 May 2017 amending Annexes I and
 27 405 IV to Regulation (EC) No 999/2001 of the European Parliament and of the Council
 28 406 and Annexes X, XIV and XV to Commission Regulation (EU) No 142/2011 as
 29 407 regards the provisions on processed animal protein.
 30 408 EU, 2012. Commission Regulation (EU) No 277/2012 of 28 March 2012 amending Annexes I
 31 409 and II to Directive 2002/32/EC of the European Parliament and of the Council as
 32 410 regards maximum levels and action thresholds for dioxins and polychlorinated
 33 411 biphenyls.
 34 412 Gao, Q., Deng, W., Gao, Z., Li, M., Liu, W., Wang, X. and Zhu, F., 2019. Effect of
 35 413 sulfonamide pollution on the growth of manure management candidate *Hermetia*
 36 414 *illucens*. PLOS ONE 14(5): e0216086. <https://doi.org/10.1371/journal.pone.0216086>
 37 415 Hawker, D.W. and Connell, D.W., 1988. Octanol-water partition coefficients of
 38 416 polychlorinated biphenyl congeners. Environmental Science & Technology 22: 382-
 39 417 387. <https://doi.org/10.1021/es00169a004>
 40 418 Meijer, N., de Rijk, T., Van Loon, J.J.A., Zoet, L. and Van der Fels-Klerx, H.J., 2021. Effects
 41 419 of insecticides on mortality, growth and bioaccumulation in black soldier fly
 42 420 (*Hermetia illucens*) larvae. PLOS ONE 16(4): e0249362.
 43 421 <https://doi.org/10.1371/journal.pone.0249362>
 44 422 Meyer, A.M., Meijer, N., Hoek-van den Hil, E.F. and Van der Fels-Klerx, H.J., 2021.
 45 423 Chemical food safety hazards of insects reared for food and feed. Journal of Insects as
 46 424 Food and Feed 7(5): 823-831. <https://doi.org/10.3920/JIFF2020.0085>
 47 425 Niermans, K., Woyzichovski, J., Kröncke, N., Benning, R. and Maul, R., 2019. Feeding study
 48 426 for the mycotoxin zearalenone in yellow mealworm (*Tenebrio molitor*) larvae-

- 427 investigation of biological impact and metabolic conversion. *Mycotoxin Research* 35:
1 428 231-242. <https://doi.org/10.1007/s12550-019-00346-y>
- 2 429 Planche, C., Ratel, J., Blinet, P., Mercier, F., Angénioux, M., Chafey, C., Zinck, J., Marchond,
3 430 N., Chevolleau, S., Marchand, P., Dervilly-Pinel, G., Guérin, T., Debrauwer, L. and
4 431 Engel, E., 2017. Effects of pan cooking on micropollutants in meat. *Food Chemistry*
5 432 232: 395-404. <https://doi.org/10.1016/j.foodchem.2017.03.049>
- 7 433 Planche, C., Ratel, J., Mercier, F., Blinet, P., Debrauwer, L. and Engel, E., 2015. Assessment
8 434 of comprehensive two-dimensional gas chromatography-time-of-flight mass
9 435 spectrometry based methods for investigating 206 dioxin-like micropollutants in
10 436 animal-derived food matrices. *Journal of Chromatography A* 1392: 74-81.
11 437 <https://doi.org/10.1016/j.chroma.2015.02.054>
- 13 438 Tran, G., Heuzé, V. and Makkar, H.P.S., 2015. Insects in fish diets. *Animal Frontiers* 5(2):
14 439 37-44. <https://doi.org/10.2527/af.2015-0018>
- 16 440 Truzzi, C., Illuminati, S., Girolametti, F., Antonucci, M., Scarponi, G., Ruschioni, S., Riolo,
17 441 P. and Annibaldi, A., 2019. Influence of feeding substrates on the presence of toxic
18 442 metals (Cd, Pb, Ni, As, Hg) in larvae of *Tenebrio molitor*: risk assessment for human
19 443 consumption. *International Journal of Environmental Research and Public Health* 16:
20 444 4815. <https://doi.org/10.3390/ijerph16234815>
- 22 445 United Nations, Department of Economic and Social Affairs, Population Division, 2019.
23 446 World population prospects 2019 highlights.
- 24 447 Van Broekhoven, S., Gutierrez, J.M., De Rijk, T.C., De Nijs, W.C.M. and Van Loon, J.J.A.,
25 448 2017. Degradation and excretion of the *Fusarium* toxin deoxynivalenol by an edible
26 449 insect, the yellow mealworm (*Tenebrio molitor* L.). *World Mycotoxin Journal* 10(2):
27 450 163-169. <https://doi.org/10.3920/WMJ2016.2102>
- 29 451 Van der Fels-Klerx, H.J., Meijer, N., Nijkamp, M.M., Schmitt, E. and Van Loon, J.J.A., 2020.
30 452 Chemical food safety of using former foodstuffs for rearing black soldier fly larvae
31 453 (*Hermetia illucens*) for feed and food use. *Journal of Insects as Food and Feed* 6(5):
32 454 475-488. <https://doi.org/10.3920/JIFF2020.0024>
- 34 455 Van Huis, A., 2022. Edible insects: challenges and prospects. *Entomological Research* 52:
35 456 161-177. <https://doi.org/10.1111/1748-5967.12582>
- 36 457 Van Huis, A., 2021. Prospects of insects as food and feed. *Organic Agriculture* 11: 301-308.
37 458 <https://doi.org/10.1007/s13165-020-00290-7>
- 39 459 Vandeweyer, D., De Smet, J., Van Looveren, N. and Van Campenhout, L., 2021. Biological
40 460 contaminants in insects as food and feed. *Journal of Insects as Food and Feed* 7(5):
41 461 807-822. <https://doi.org/10.3920/JIFF2020.0060>
- 42 462





Figure 2

[Click here to access/download;Figure;Figure 2 PDF.pdf](#)



Supplementary material

Fate of polychlorobiphenyls in ~~the insect~~ *Tenebrio molitor* larvae: consequences for further use as food and feed

J. Ratel¹, F. Mercier¹, J. Rivas¹, H. Wang¹, M. Angénieux¹, B. Calmont², S. Crépieux³, C. Planche^{1*}, E. Engel¹




Figure S1: Temperature (°C) and relative humidity (%) readings of each incubator during the larvae rearing.




Figure S2: Weight of *Tenebrio molitor* larvae (mg) throughout their rearing on wheat bran either unspiked (Control), spiked with neat solvent (Control + acetone) or spiked at 0.67, 4.0 or 24.4 ng PCB/g wheat bran. Data represent mean larvae weight (mg) \pm SD which is determined from the 20 larvae measurement for each rearing glass jar ($n=3$ glass jars for each exposure condition).

CONFIDENTIAL

Table S1: Performance of GC- μ ECD for quantification of the 6 nDL-PCB congeners in dried powder of *Tenebrio molitor* larvae (linearity range: 0.2-50.0 ng/g).

PCB congener	Coefficient of determination (R²)	Limit of detection (LOD) in ng/g	Limit of quantification (LOQ) in ng/g
28	0.95	2.80	9.32
52	0.96	2.40	8.00
101	0.94	2.98	9.93
138	0.98	1.76	5.88
153	0.98	1.67	5.56
180	0.98	1.68	5.61

CONFIDENTIAL

Table S2: Concentrations (ng/g) of PCB congeners in *Tenebrio molitor* larvae on a dried weight basis after 20 days of rearing on wheat bran spiked at 0.67, 4.0 or 24.4 ng PCB/g. For each exposure condition, three analytical replicates of each rearing glass jar were carried out ($n=3$ glass jars for each exposure condition).

Congener	Wheat bran contamination with PCBs		
	0.67 ppb	4.0 ppb	24.4 ppb
PCB 28	NA	5.26*	44.03
	NA	6.73*	51.78
	NA	9.98	48.67
	NA	8.98*	48.94
	NA	9.67	44.26
	NA	10.71	45.12
	NA	9.94	47.06
	NA	10.89	47.77
	NA	8.79*	46.21
PCB 52	NA	3.28*	20.82
	NA	3.28*	25.54
	NA	5.38*	25.38
	NA	4.25*	26.93
	NA	4.43*	25.05
	NA	5.09*	24.18
	NA	4.28*	25.57
	NA	5.12*	25.87
	NA	4.31*	25.44
PCB 101	NA	5.60*	44.90
	NA	5.77*	47.35
	NA	7.81*	46.75
	NA	6.75*	48.12
	NA	7.56*	44.12
	NA	8.85*	44.60
	NA	7.84*	47.83
	NA	8.88*	48.04
	NA	7.20*	46.44
PCB 138	<LOD	7.17	67.28
	<LOD	7.21	62.43
	<LOD	9.32	63.24
	<LOD	8.26	69.15
	<LOD	10.05	64.03
	<LOD	11.55	63.56
	<LOD	10.58	70.78
	<LOD	12.40	69.11
	<LOD	10.00	68.47
PCB 153	1.96*	7.65	68.68
	<LOD	7.70	64.16
	<LOD	8.84	65.71
	<LOD	7.43	70.40

	<LOD	9.56	65.13
	<LOD	10.69	65.32
	1.77*	10.71	73.80
	<LOD	11.61	73.50
	<LOD	9.60	71.97
	1.86*	7.89	71.41
	<LOD	8.23	59.18
	<LOD	7.58	61.84
	<LOD	7.46	65.84
PCB 180	<LOD	8.91	63.52
	<LOD	9.92	63.93
	1.71*	9.80	75.07
	<LOD	11.33	73.20
	<LOD	10.70	73.05

NA: non-available (non-detected in larvae extracts)

LOD: Limit of detection

* Quantification values between LOD and LOQ (see Table S1)

CONFIDENTIAL

Table S3: Raw data enabling the calculation of bioaccumulation factors (BAF) of PCB congeners in *Tenebrio molitor* larvae on a dry weight basis after 20 days of rearing on wheat bran spiked at 0.67, 4.0 or 24.4 ng PCB/g.

Wheat bran PCB conc. (ppb)	Replicate ^a	Sample (g)	IS ^b			PCB 28			PCB 52			PCB 101			PCB 138			PCB 153			PCB 180			
			Area	Area	Conc. (ppb) ^c	BAF ^d	Area	Area	Conc. (ppb)	BAF	Area	Area	Conc. (ppb)	BAF	Area	Area	Conc. (ppb)	BAF	Area	Area	Conc. (ppb)	BAF	Area	Area
0,67	1-R1	1,0019	5,56E+07	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	9,94E+07	1,69	2,53	1,46E+08	1,96	2,94	1,21E+08	1,86	2,79
0,67	1-R2	0,9993	6,60E+07	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	8,49E+07	1,17	1,75	1,37E+08	1,52	2,29	1,20E+08	1,54	2,31
0,67	1-R3	0,9991	7,14E+07	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1,13E+08	1,48	2,22	1,29E+08	1,31	1,97	1,11E+08	1,31	1,97
0,67	2-R1	1,0002	6,44E+07	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	9,85E+07	1,42	2,13	1,16E+08	1,30	1,95	9,84E+07	1,29	1,93
0,67	2-R2	1,0003	5,79E+07	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	8,16E+07	1,30	1,94	1,10E+08	1,39	2,08	1,05E+08	1,54	2,32
0,67	2-R3	1,0011	7,27E+07	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1,08E+08	1,37	2,06	1,43E+08	1,44	2,15	1,16E+08	1,35	2,02
0,67	3-R1	0,9982	6,14E+07	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1,05E+08	1,60	2,41	1,47E+08	1,77	2,66	1,23E+08	1,71	2,56
0,67	3-R2	0,9994	7,31E+07	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1,28E+08	1,66	2,49	1,37E+08	1,36	2,04	1,18E+08	1,37	2,05
0,67	3-R3	0,9984	6,51E+07	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1,11E+08	1,61	2,42	1,36E+08	1,53	2,30	1,14E+08	1,49	2,23
4	1-R1	1,0012	5,49E+07	3,95E+08	5,26	1,32	1,55E+08	3,28	0,82	3,32E+08	5,60	1,40	3,88E+08	7,17	1,79	5,38E+08	7,17	1,79	5,38E+08	7,65	1,91	4,95E+08	7,89	1,97
4	1-R2	1,0011	4,59E+07	4,16E+08	6,73	1,68	1,30E+08	3,28	0,82	2,85E+08	5,77	1,44	3,26E+08	7,21	1,80	4,52E+08	7,70	1,93	4,52E+08	7,70	1,93	4,31E+08	8,23	2,06
4	1-R3	1,0006	4,28E+07	5,65E+08	9,98	2,49	1,94E+08	5,38	1,34	3,58E+08	7,81	1,95	3,91E+08	9,32	2,33	4,83E+08	8,84	2,21	4,83E+08	8,84	2,21	3,71E+08	7,58	1,89
4	2-R1	1,0006	4,65E+07	5,54E+08	8,98	2,25	1,68E+08	4,25	1,06	3,36E+08	6,75	1,69	3,76E+08	8,26	2,06	4,42E+08	7,43	1,86	4,42E+08	7,43	1,86	3,96E+08	7,46	1,86
4	2-R2	1,0009	4,43E+07	5,67E+08	9,67	2,42	1,66E+08	4,43	1,11	3,58E+08	7,56	1,89	4,35E+08	10,05	2,51	5,40E+08	9,56	2,39	5,40E+08	9,56	2,39	4,50E+08	8,91	2,23
4	2-R3	1,0004	3,92E+07	5,54E+08	10,71	2,68	1,68E+08	5,09	1,27	3,70E+08	8,85	2,21	4,41E+08	11,55	2,89	5,34E+08	10,69	2,67	5,34E+08	10,69	2,67	4,43E+08	9,92	2,48
4	3-R1	1,0009	4,01E+07	5,28E+08	9,94	2,49	1,46E+08	4,28	1,07	3,36E+08	7,84	1,96	4,14E+08	10,58	2,65	5,47E+08	10,71	2,68	5,47E+08	10,71	2,68	4,48E+08	9,80	2,45
4	3-R2	1,0006	4,11E+07	5,90E+08	10,89	2,72	1,77E+08	5,12	1,28	3,89E+08	8,88	2,22	4,96E+08	12,40	3,10	6,07E+08	11,61	2,90	6,07E+08	11,61	2,90	5,30E+08	11,33	2,83
4	3-R3	1,0006	4,32E+07	5,05E+08	8,79	2,20	1,58E+08	4,31	1,08	3,33E+08	7,20	1,80	4,22E+08	10,00	2,50	5,29E+08	9,60	2,40	5,29E+08	9,60	2,40	5,27E+08	10,70	2,67
24,4	1-R1	1,0002	4,35E+07	2,47E+09	44,03	1,80	7,43E+08	20,82	0,85	2,05E+09	44,90	1,84	2,82E+09	67,28	2,75	3,77E+09	68,68	2,81	3,77E+09	68,68	2,81	3,52E+09	71,41	2,92
24,4	1-R2	1,001	4,37E+07	2,91E+09	51,78	2,12	9,13E+08	25,54	1,04	2,17E+09	47,35	1,94	2,63E+09	62,43	2,55	3,54E+09	64,16	2,62	3,54E+09	64,16	2,62	2,93E+09	59,18	2,42
24,4	1-R3	1,0006	3,80E+07	2,38E+09	48,67	1,99	7,89E+08	25,38	1,04	1,86E+09	46,75	1,91	2,31E+09	63,24	2,59	3,15E+09	65,71	2,69	3,15E+09	65,71	2,69	2,66E+09	61,84	2,53
24,4	2-R1	1,0009	4,20E+07	2,65E+09	48,94	2,00	9,26E+08	26,93	1,10	2,12E+09	48,12	1,97	2,80E+09	69,15	2,83	3,73E+09	70,40	2,88	3,73E+09	70,40	2,88	3,14E+09	65,84	2,69
24,4	2-R2	1,001	4,50E+07	2,56E+09	44,26	1,81	9,23E+08	25,05	1,02	2,08E+09	44,12	1,80	2,78E+09	64,03	2,62	3,70E+09	65,13	2,66	3,70E+09	65,13	2,66	3,24E+09	63,52	2,60
24,4	2-R3	1,0001	4,56E+07	2,65E+09	45,12	1,85	9,02E+08	24,18	0,99	2,13E+09	44,60	1,82	2,79E+09	63,56	2,60	3,75E+09	65,32	2,67	3,75E+09	65,32	2,67	3,30E+09	63,93	2,62
24,4	3-R1	1,0007	4,25E+07	2,58E+09	47,06	1,93	8,91E+08	25,57	1,05	2,13E+09	47,83	1,96	2,90E+09	70,78	2,90	3,96E+09	73,80	3,02	3,96E+09	73,80	3,02	3,62E+09	75,07	3,07
24,4	3-R2	0,9997	4,51E+07	2,77E+09	47,77	1,95	9,54E+08	25,87	1,06	2,27E+09	48,04	1,97	3,00E+09	69,11	2,83	4,18E+09	73,50	3,01	4,18E+09	73,50	3,01	3,74E+09	73,20	2,99
24,4	3-R3	1,0011	4,16E+07	2,47E+09	46,21	1,89	8,67E+08	25,44	1,04	2,03E+09	46,44	1,90	2,75E+09	68,47	2,80	3,78E+09	71,97	2,94	3,78E+09	71,97	2,94	3,44E+09	73,05	2,99

NA: non-available (PCBs non-detected in larvae extracts)

^a For each exposure condition, three analyses (R1 to R3) of approximately 1 g of dried larvae sampled in each of the 3 glass jars (1 to 3) were carried out.

^b Internal standard (5-F-PCB-126) used for the accurate quantification of the nDL-PCBs

^c PCB concentration in larvae was calculated according to the formula: Concentration = $\frac{((\text{Area PCB}_x / \text{Area IS}) - b) / a}{\text{Sample weight} \times \text{CF}}$ where CF is the concentration factor due to the extraction procedure (CF=10), and a and b are the slope and y-intercept, respectively, of the calibration curves:

	PCB 28	PCB 52	PCB 101	PCB 138	PCB 153	PCB 180
R ²	0,95	0,96	0,94	0,98	0,98	0,98
a	0,13	0,08	0,10	0,10	0,13	0,11
b	0,46	0,16	0,19	0,16	0,16	0,07

^d BAF was calculated according to the formula: BAF = $\frac{(\text{Larvae PCB}_x \text{ concentration})}{(\text{Wheat bran PCB}_x \text{ concentration})}$

Table S4: Bioaccumulation factors (BAF) of PCB congeners in *Tenebrio molitor* larvae on a dry weight basis after 20 days of rearing on wheat bran spiked at 0.67, 4.0 or 24.4 ng PCB/g. Data represent mean BAF \pm SD which is determined from 3 analytical replicates of each rearing glass jar ($n=3$ glass jars for each exposure condition).

Congener	Wheat bran contamination with PCBs		
	0.67 ppb	4.0 ppb	24.4 ppb
PCB 28	NA	2.25 \pm 0.47 ^{a*}	1.93 \pm 0.10 ^a
PCB 52	NA	1.10 \pm 0.19 ^{a*}	1.02 \pm 0.07 ^a
PCB 101	NA	1.84 \pm 0.29 ^{a*}	1.90 \pm 0.06 ^a
PCB 138	NA	2.40 \pm 0.45 ^a	2.72 \pm 0.13 ^a
PCB 153	NA	2.33 \pm 0.38 ^a	2.81 \pm 0.16 ^b
PCB 180	NA	2.27 \pm 0.35 ^a	2.76 \pm 0.24 ^b

NA: non-available (PCBs non-detected in larvae extracts)

a–b: different superscript letters within the same row indicate significant differences among values ($p<0.05$). BAF are determined from quantification values in larvae extracts between LOD and LOQ (*) or greater than LOQ.

CONFIDENTIAL

Supplementary material

Fate of polychlorobiphenyls in *Tenebrio molitor* larvae: consequences for further use as food and feed

J. Ratel¹, F. Mercier¹, J. Rivas¹, H. Wang¹, M. Angénieux¹, B. Calmont², S. Crépieux³, C. Planche^{1*}, E. Engel¹




Figure S1: Temperature (°C) and relative humidity (%) readings of each incubator during the larvae rearing.




Figure S2: Weight of *Tenebrio molitor* larvae (mg) throughout their rearing on wheat bran either unspiked (Control), spiked with neat solvent (Control + acetone) or spiked at 0.67, 4.0 or 24.4 ng PCB/g wheat bran. Data represent mean larvae weight (mg) \pm SD which is determined from the 20 larvae measurement for each rearing glass jar ($n=3$ glass jars for each exposure condition).

CONFIDENTIAL

Table S1: Performance of GC- μ ECD for quantification of the 6 nDL-PCB congeners in dried powder of *Tenebrio molitor* larvae (linearity range: 0.2-50.0 ng/g).

PCB congener	Coefficient of determination (R²)	Limit of detection (LOD) in ng/g	Limit of quantification (LOQ) in ng/g
28	0.95	2.80	9.32
52	0.96	2.40	8.00
101	0.94	2.98	9.93
138	0.98	1.76	5.88
153	0.98	1.67	5.56
180	0.98	1.68	5.61

CONFIDENTIAL

Table S2: Concentrations (ng/g) of PCB congeners in *Tenebrio molitor* larvae on a dried weight basis after 20 days of rearing on wheat bran spiked at 0.67, 4.0 or 24.4 ng PCB/g. For each exposure condition, three analytical replicates of each rearing glass jar were carried out ($n=3$ glass jars for each exposure condition).

Congener	Wheat bran contamination with PCBs		
	0.67 ppb	4.0 ppb	24.4 ppb
PCB 28	NA	5.26*	44.03
	NA	6.73*	51.78
	NA	9.98	48.67
	NA	8.98*	48.94
	NA	9.67	44.26
	NA	10.71	45.12
	NA	9.94	47.06
	NA	10.89	47.77
	NA	8.79*	46.21
PCB 52	NA	3.28*	20.82
	NA	3.28*	25.54
	NA	5.38*	25.38
	NA	4.25*	26.93
	NA	4.43*	25.05
	NA	5.09*	24.18
	NA	4.28*	25.57
	NA	5.12*	25.87
	NA	4.31*	25.44
PCB 101	NA	5.60*	44.90
	NA	5.77*	47.35
	NA	7.81*	46.75
	NA	6.75*	48.12
	NA	7.56*	44.12
	NA	8.85*	44.60
	NA	7.84*	47.83
	NA	8.88*	48.04
	NA	7.20*	46.44
PCB 138	<LOD	7.17	67.28
	<LOD	7.21	62.43
	<LOD	9.32	63.24
	<LOD	8.26	69.15
	<LOD	10.05	64.03
	<LOD	11.55	63.56
	<LOD	10.58	70.78
	<LOD	12.40	69.11
	<LOD	10.00	68.47
PCB 153	1.96*	7.65	68.68
	<LOD	7.70	64.16
	<LOD	8.84	65.71
	<LOD	7.43	70.40

	<LOD	9.56	65.13
	<LOD	10.69	65.32
	1.77*	10.71	73.80
	<LOD	11.61	73.50
	<LOD	9.60	71.97
	1.86*	7.89	71.41
	<LOD	8.23	59.18
	<LOD	7.58	61.84
	<LOD	7.46	65.84
PCB 180	<LOD	8.91	63.52
	<LOD	9.92	63.93
	1.71*	9.80	75.07
	<LOD	11.33	73.20
	<LOD	10.70	73.05

NA: non-available (non-detected in larvae extracts)

LOD: Limit of detection

* Quantification values between LOD and LOQ (see Table S1)

CONFIDENTIAL

Table S3: Raw data enabling the calculation of bioaccumulation factors (BAF) of PCB congeners in *Tenebrio molitor* larvae on a dry weight basis after 20 days of rearing on wheat bran spiked at 0.67, 4.0 or 24.4 ng PCB/g.

Wheat bran PCB conc. (ppb)	Replicate ^a	Sample (g)	IS ^b			PCB 28			PCB 52			PCB 101			PCB 138			PCB 153			PCB 180					
			Area	Area	Conc. (ppb) ^c	BAF ^d	Area	Area	Conc. (ppb)	BAF	Area	Area	Conc. (ppb)	BAF	Area	Area	Conc. (ppb)	BAF	Area	Area	Conc. (ppb)	BAF	Area	Area	Conc. (ppb)	BAF
0,67	1-R1	1,0019	5,56E+07	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	9,94E+07	1,69	2,53	1,46E+08	1,96	2,94	1,21E+08	1,86	2,79
0,67	1-R2	0,9993	6,60E+07	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	8,49E+07	1,17	1,75	1,37E+08	1,52	2,29	1,20E+08	1,54	2,31
0,67	1-R3	0,9991	7,14E+07	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1,13E+08	1,48	2,22	1,29E+08	1,31	1,97	1,11E+08	1,31	1,97
0,67	2-R1	1,0002	6,44E+07	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	9,85E+07	1,42	2,13	1,16E+08	1,30	1,95	9,84E+07	1,29	1,93
0,67	2-R2	1,0003	5,79E+07	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	8,16E+07	1,30	1,94	1,10E+08	1,39	2,08	1,05E+08	1,54	2,32
0,67	2-R3	1,0011	7,27E+07	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1,08E+08	1,37	2,06	1,43E+08	1,44	2,15	1,16E+08	1,35	2,02
0,67	3-R1	0,9982	6,14E+07	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1,05E+08	1,60	2,41	1,47E+08	1,77	2,66	1,23E+08	1,71	2,56
0,67	3-R2	0,9994	7,31E+07	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1,28E+08	1,66	2,49	1,37E+08	1,36	2,04	1,18E+08	1,37	2,05
0,67	3-R3	0,9984	6,51E+07	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1,11E+08	1,61	2,42	1,36E+08	1,53	2,30	1,14E+08	1,49	2,23
4	1-R1	1,0012	5,49E+07	3,95E+08	5,26	1,32	1,55E+08	3,28	0,82	3,32E+08	5,60	1,40	3,88E+08	7,17	1,79	5,38E+08	7,65	1,91	4,95E+08	7,89	1,97					
4	1-R2	1,0011	4,59E+07	4,16E+08	6,73	1,68	1,30E+08	3,28	0,82	2,85E+08	5,77	1,44	3,26E+08	7,21	1,80	4,52E+08	7,70	1,93	4,31E+08	8,23	2,06					
4	1-R3	1,0006	4,28E+07	5,65E+08	9,98	2,49	1,94E+08	5,38	1,34	3,58E+08	7,81	1,95	3,91E+08	9,32	2,33	4,83E+08	8,84	2,21	3,71E+08	7,58	1,89					
4	2-R1	1,0006	4,65E+07	5,54E+08	8,98	2,25	1,68E+08	4,25	1,06	3,66E+08	6,75	1,69	3,76E+08	8,26	2,06	4,42E+08	7,43	1,86	3,96E+08	7,46	1,86					
4	2-R2	1,0009	4,43E+07	5,67E+08	9,67	2,42	1,66E+08	4,43	1,11	3,58E+08	7,56	1,89	4,35E+08	10,05	2,51	5,40E+08	9,56	2,39	4,50E+08	8,91	2,23					
4	2-R3	1,0004	3,92E+07	5,54E+08	10,71	2,68	1,68E+08	5,09	1,27	3,70E+08	8,85	2,21	4,41E+08	11,55	2,89	5,34E+08	10,69	2,67	4,43E+08	9,92	2,48					
4	3-R1	1,0009	4,01E+07	5,28E+08	9,94	2,49	1,46E+08	4,28	1,07	3,36E+08	7,84	1,96	4,14E+08	10,58	2,65	5,47E+08	10,71	2,68	4,48E+08	9,80	2,45					
4	3-R2	1,0006	4,11E+07	5,90E+08	10,89	2,72	1,77E+08	5,12	1,28	3,89E+08	8,88	2,22	4,96E+08	12,40	3,10	6,07E+08	11,61	2,90	5,30E+08	11,33	2,83					
4	3-R3	1,0006	4,32E+07	5,05E+08	8,79	2,20	1,58E+08	4,31	1,08	3,33E+08	7,20	1,80	4,22E+08	10,00	2,50	5,29E+08	9,60	2,40	5,27E+08	10,70	2,67					
24,4	1-R1	1,0002	4,35E+07	2,47E+09	44,03	1,80	7,43E+08	20,82	0,85	2,05E+09	44,90	1,84	2,82E+09	67,28	2,75	3,77E+09	68,68	2,81	3,52E+09	71,41	2,92					
24,4	1-R2	1,001	4,37E+07	2,91E+09	51,78	2,12	9,13E+08	25,54	1,04	2,17E+09	47,35	1,94	2,63E+09	62,43	2,55	3,54E+09	64,16	2,62	2,93E+09	59,18	2,42					
24,4	1-R3	1,0006	3,80E+07	2,38E+09	48,67	1,99	7,89E+08	25,38	1,04	1,86E+09	46,75	1,91	2,31E+09	63,24	2,59	3,15E+09	65,71	2,69	2,66E+09	61,84	2,53					
24,4	2-R1	1,0009	4,20E+07	2,65E+09	48,94	2,00	9,26E+08	26,93	1,10	2,12E+09	48,12	1,97	2,80E+09	69,15	2,83	3,73E+09	70,40	2,88	3,14E+09	65,84	2,69					
24,4	2-R2	1,001	4,50E+07	2,56E+09	44,26	1,81	9,23E+08	25,05	1,02	2,08E+09	44,12	1,80	2,78E+09	64,03	2,62	3,70E+09	65,13	2,66	3,24E+09	63,52	2,60					
24,4	2-R3	1,0001	4,56E+07	2,65E+09	45,12	1,85	9,02E+08	24,18	0,99	2,13E+09	44,60	1,82	2,79E+09	63,56	2,60	3,75E+09	65,32	2,67	3,30E+09	63,93	2,62					
24,4	3-R1	1,0007	4,25E+07	2,58E+09	47,06	1,93	8,91E+08	25,57	1,05	2,13E+09	47,83	1,96	2,90E+09	70,78	2,90	3,96E+09	73,80	3,02	3,62E+09	75,07	3,07					
24,4	3-R2	0,9997	4,51E+07	2,77E+09	47,77	1,95	9,54E+08	25,87	1,06	2,27E+09	48,04	1,97	3,00E+09	69,11	2,83	4,18E+09	73,50	3,01	3,74E+09	73,20	2,99					
24,4	3-R3	1,0011	4,16E+07	2,47E+09	46,21	1,89	8,67E+08	25,44	1,04	2,03E+09	46,44	1,90	2,75E+09	68,47	2,80	3,78E+09	71,97	2,94	3,44E+09	73,05	2,99					

NA: non-available (PCBs non-detected in larvae extracts)

^a For each exposure condition, three analyses (R1 to R3) of approximately 1 g of dried larvae sampled in each of the 3 glass jars (1 to 3) were carried out.

^b Internal standard (5-F-PCB-126) used for the accurate quantification of the nDL-PCBs

^c PCB concentration in larvae was calculated according to the formula: Concentration = $\frac{((\text{Area PCB}_x / \text{Area IS}) - b) / a}{\text{Sample weight} \times \text{CF}}$ where CF is the concentration factor due to the extraction procedure (CF=10), and a and b are the slope and y-intercept, respectively, of the calibration curves:

	PCB 28	PCB 52	PCB 101	PCB 138	PCB 153	PCB 180
R ²	0,95	0,96	0,94	0,98	0,98	0,98
a	0,13	0,08	0,10	0,10	0,13	0,11
b	0,46	0,16	0,19	0,16	0,16	0,07

^d BAF was calculated according to the formula: BAF = (Larvae PCB_x concentration) / (Wheat bran PCB_x concentration)

Table S4: Bioaccumulation factors (BAF) of PCB congeners in *Tenebrio molitor* larvae on a dry weight basis after 20 days of rearing on wheat bran spiked at 0.67, 4.0 or 24.4 ng PCB/g. Data represent mean BAF \pm SD which is determined from 3 analytical replicates of each rearing glass jar ($n=3$ glass jars for each exposure condition).

Congener	Wheat bran contamination with PCBs		
	0.67 ppb	4.0 ppb	24.4 ppb
PCB 28	NA	2.25 \pm 0.47 ^{a*}	1.93 \pm 0.10 ^a
PCB 52	NA	1.10 \pm 0.19 ^{a*}	1.02 \pm 0.07 ^a
PCB 101	NA	1.84 \pm 0.29 ^{a*}	1.90 \pm 0.06 ^a
PCB 138	NA	2.40 \pm 0.45 ^a	2.72 \pm 0.13 ^a
PCB 153	NA	2.33 \pm 0.38 ^a	2.81 \pm 0.16 ^b
PCB 180	NA	2.27 \pm 0.35 ^a	2.76 \pm 0.24 ^b

NA: non-available (PCBs non-detected in larvae extracts)

a–b: different superscript letters within the same row indicate significant differences among values ($p<0.05$). BAF are determined from quantification values in larvae extracts between LOD and LOQ (*) or greater than LOQ.

CONFIDENTIAL