



HAL
open science

Representation and functions of shoot apical meristems in FSPMs

Romain Barillot, Frédéric Boudon, Didier Combes, Jean-Louis Durand,
Gaëtan Louarn

► **To cite this version:**

Romain Barillot, Frédéric Boudon, Didier Combes, Jean-Louis Durand, Gaëtan Louarn. Representation and functions of shoot apical meristems in FSPMs. FSPM2023, Mar 2023, Berlin, Germany. hal-04098500

HAL Id: hal-04098500

<https://hal.inrae.fr/hal-04098500>

Submitted on 16 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Representation and functions of shoot apical meristems in FSPMs

Romain Barillot¹, Frédéric Boudon^{2,3}, Didier Combes¹, Jean-Louis Durand¹, Gaëtan Louarn¹

¹ INRAE, UR P3F, Lusignan, F86600, France

² CIRAD, UMR AGAP Institut, F-34398 Montpellier, France

³ UMR AGAP Institut, Univ. Montpellier, CIRAD, INRAE, Institut Agro, Montpellier, France

For correspondence: romain.barillot@inrae.fr

Keywords: FSPM, shoot apical meristem, organogenesis, floral transition, scale integration

Functional Structural Plant Models (FSPMs) are individual-based models that explicitly account for the interactions between plant architecture and its abiotic and biotic environment. Plant morphogenesis, growth and grain or fruit production are among the most represented processes in FSPMs. These biological processes are largely determined at the Shoot Apical Meristem (SAM) level as it drives the rate of leaf initiation, phyllotaxy, leaf geometry, floral induction and the production of reproductive organs. Nevertheless, not all FSPMs are based on an explicit representation of SAMs, their functioning is then ignored in the case of non-dynamic models or embedded in parametric or statistical functions of plant morphogenesis in dynamic models. For FSPMs that include an explicit representation of SAMs, they are mainly constructed on the L-systems formalism, which is well suited to describe plant development (Boudon et al., 2012). First, one of the main functions of SAMs in these FSPMs is the production of new leaves or growth units, usually at a constant rate expressed in thermal unit. In some cases, the rate of leaf production by SAMs is coordinated with the rate of leaf emergence, leading to the concept of self-regulated architecture. However, the effect of substrate or water availability on the functioning of the SAM in terms of the size or properties of the emitted primordia is not accounting for. Secondly, the representation of apices in models is also used to simulate axillary bud break and the production of tillers, branches or new growth units. In these cases, the transition from a latent to an active SAM is controlled by light intensity and its spectrum (Verdenal et al., 2008; Faverjon et al., 2019), or temperature or hormones (Prusinkiewicz et al., 2009) or is based on stochastic approaches. Finally, some FSPMs also account for the production of reproductive organs like grains and fruits by SAMs (Boudon et al., 2020; Rouet et al., 2022). Nevertheless, this aspect of SAM functioning remains poorly described in FSPMs despite its importance in plant production, yield and ecology. We believe that a better integration of SAMs functioning and control in FSPMs is a promising way to assess functional hypotheses and predict plant plasticity to the environment.

References

- Boudon F et al. 2020. *Annals of Botany* 126: 745–763, doi: <https://doi.org/10.1093/aob/mcaa089> .
- Boudon F, Pradal C, Cokelaer T, Prusinkiewicz P, Godin C. 2012. *Frontiers in Plant Science* 3, doi: <https://doi.org/10.3389/fpls.2012.00076>
- Faverjon L, Escobar-Gutiérrez A, Litrico I, Julier B, Louarn G. 2019. *Journal of Experimental Botany* 70: 2491–2504, doi: <https://doi.org/10.1093/jxb/ery323>
- Prusinkiewicz P, Crawford S, Smith RS, Ljung K, Bennett T, Ongaro V, Leyser O. 2009. *Proceedings of the National Academy of Sciences* 106: 17431–17436, doi: <https://doi.org/10.1073/pnas.0906696106>
- Rouet S, Durand J-L, Leclercq D, Bernicot M-H, Combes D, Escobar-Gutiérrez A, Barillot R. 2022. *in silico Plants* 4: diac012, doi: <https://doi.org/10.1093/insilicoplants/diac012>
- Verdenal A, Combes D, Escobar-Gutiérrez AJ. 2008. *Functional Plant Biology* 35: 911–924, doi: <https://doi.org/10.1071/fp08050>