



# Time-Restricted Feeding Potentiates the Ability of *Lacticaseibacillus casei* to Enrich the Retina in Omega-3 Fatty Acids

Pierre Lapaquette, Anne-Sophie Boucard, Florian Chain, Stéphane Grégoire, Luis Bermúdez-Humarán, Niyazi Acar, Marie-Agnès Bringer

## ► To cite this version:

Pierre Lapaquette, Anne-Sophie Boucard, Florian Chain, Stéphane Grégoire, Luis Bermúdez-Humarán, et al.. Time-Restricted Feeding Potentiates the Ability of *Lacticaseibacillus casei* to Enrich the Retina in Omega-3 Fatty Acids. *Aging and disease*, 2023, 14 (6), pp.1945-1949. 10.14336/AD.2023.0324 . hal-04101285

**HAL Id: hal-04101285**

**<https://hal.inrae.fr/hal-04101285>**

Submitted on 19 May 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Letter to the Editor

# Time-Restricted Feeding Potentiates the Ability of *Lactocaseibacillus casei* to Enrich the Retina in Omega-3 Fatty Acids

Pierre Lapaquette<sup>1</sup>, Anne-Sophie Boucard<sup>2</sup>, Florian Chain<sup>2</sup>, Stéphane Grégoire<sup>3</sup>, Luis G. Bermúdez-Humarán<sup>2</sup>, Niyazi Acar<sup>3</sup>, Marie-Agnès Bringer<sup>3\*</sup>

<sup>1</sup>Univ. Bourgogne Franche-Comté, Agrosup Dijon, UMR PAM A 02.102, Dijon, France. <sup>2</sup>INRAE, AgroParisTech, Micalis Institute, Université Paris-Saclay, 78350, Jouy-en-Josas, France. <sup>3</sup>Centre des Sciences du Goût et de l'Alimentation, CNRS, INRAE, Institut Agro, Université de Bourgogne Franche-Comté, Dijon, France.

[Received February 12, 2023; Revised March 20, 2023; Accepted March 24, 2023]

To the Editor,

Current treatments for age-related macular degeneration (AMD) —the primary cause of visual impairment in western countries in people over 50 [1]— are limited to a single form of the disease and cannot always avoid severe visual loss. Among emerging preventive strategies, omega-3 (n-3) polyunsaturated fatty acids (PUFAs) are drawing interest as they promote normal retinal structure and function, reduce incidence, and slow progression of AMD, consistently with their abundance in retinal neurons. Evidence has also accumulated suggesting the influence of the gut microbiota in AMD pathophysiology [2]. In this work, we investigated whether strategies targeting the gut microbiota could be effective in modulating retinal PUFA content.

Changes in the gut microbiota of AMD patients have been reported and several experimental studies suggest the existence of a gut-retina axis, in which gut microbes would influence AMD-associated mechanisms (inflammation, pathological vascularization and neurodegeneration) and modulate retina physiology, including its lipid composition [2]. Approaches targeting the gut microbiota such as probiotic-based strategies and intermittent fasting —a dietary intervention that shapes the gut microbiota to benefit health— have been shown to influence lipid metabolism and to protect the retina against harmful conditions [2, 3]. In the urgency context to find new strategies to prevent or delay the onset and

progression of AMD, we evaluated the ability of supplementation with the probiotic *Lactocaseibacillus casei* (*L. casei*) strain BL23 and time-restricted feeding intervention to modulate the PUFA content in the mouse retina.

To do so, C57BL/6J mice were randomly distributed into two groups: one group had a permanent access to food (*ad libitum* group) whereas the other had a time-restricted access to food (isocaloric twice-a-day feeding (ITAD) group, [4]) (Supplementary Fig. 1A and Supplementary data). Half of mice in each group were supplemented with *L. casei* BL23 by daily oral gavage (Supplementary Fig. 1B and Supplementary data). To evaluate the impact of these treatments on lipids, amounts of fatty acid methyl esters (FAMES, which are derivatives of fatty acids carried by diacylglycerophospholipids and plasmalogens) and dimethylacetals (DMAs, which are derivatives of fatty alcohols on plasmalogens) were determined by gas chromatography (Supplementary data). For microbiota analysis, 16S rRNA gene sequencing was done on fecal material using the Illumina MiSeq technology (Supplementary data).


The analysis of fecal microbiota composition revealed that alpha diversity (Shannon and inverse Simpson indices) was significantly increased in mice supplemented with *L. casei* compared to ITAD fed mice (Fig. 1A). A trend to a decreased diversity (Shannon indice) was observed in gut microbiota of ITAD-fed mice compared to *ad libitum*-fed mice (Fig. 1A). Linear

\*Correspondence should be addressed to: Dr. Marie-Agnès Bringer, Centre des Sciences du Goût et de l'Alimentation, CNRS, INRAE, Institut Agro, Université de Bourgogne Franche-Comté, F-21000 Dijon, France. Email: [marie-agnes.bringer@inrae.fr](mailto:marie-agnes.bringer@inrae.fr).

**Copyright:** © 2023 Lapaquette P. et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

discriminant analysis effect size (LEfSe) approach showed that the feeding rhythm and *L. casei* supplementation were associated with specific changes in gut microbiota, including overrepresentation of: (i) *Prevotellaceae* —a family including producers of short chain fatty acids that are metabolites regulating of host lipid metabolism and health [5]— in *L. casei*-


supplemented mice; (ii) *Lactobacillaceae* —a family including species having probiotic potential [6]— in ITAD-fed mice; and (iii) *Eggerthellaceae* —a family with members involved in the production of health-promoting metabolites (urolithins) [7]— in ITAD-fed mice supplemented with *L. casei* (Fig. 1B and 1C).



**Figure 1. Effect of ITAD feeding and *Lactocaseibacillus casei* BL23 supplementation on gut microbiota. (A)** Four indices (observed, Chao1, Shannon and inverse Simpson) of alpha diversity were analyzed. **(B and C)** LEfSe analysis was performed to identify significantly altered taxa in one group compared to another. **(B)** LEfSe cladogram. **(C)** LDA scores. Alpha value for the factorial Kruskal-Wallis test among classes: 0.05. Threshold on the logarithmic LDA score for discriminative features: 2.0.  $n=8$  per group. \*  $p<0.05$ , \*\*  $p<0.01$ .

The combination of ITAD feeding with *L. casei* supplementation deeply impacted lipids in the retina by significantly enriching its PUFA content (Fig. 2). The high levels of PUFAs were balanced by decreased abundances of saturated and monounsaturated fatty acids (Fig. 2A and 2B). At the species level, this strategy significantly modified the relative abundance of 17/24 lipid species *versus* 4/24 in ITAD-fed mice and 7/24 *L. casei*-supplemented mice compared to *ad libitum* fed-mice (Fig. 2B). Particularly, the abundance of several n-6 PUFAs (C20:3n-6, C20:4n-6 (arachidonic acid, ARA) and C22:4n-6) and n-3 PUFAs (C22:5n-3 and C22:6n-3

(DHA)) was significantly increased in the retina of ITAD-fed and *L. casei*-supplemented mice (Fig. 2B). Moreover, ITAD feeding combined with *L. casei* supplementation significantly increased the plasmalogens content in the retina as shown by the decrease in total FAMES (fatty acid methyl esters) / total DMAs (dimethylacetals) ratio and the increased abundance of 2 DMA species (Fig. 2H and 2I). Plasmalogens are glycerophospholipids that concentrate PUFAs such as ARA and DHA. These PUFAs may be involved in the regulation of retinal vascular development after being released from plasmalogens [8].



**Figure 2. Effect of ITAD feeding and *Lactocaseibacillus casei* BL23 supplementation on retinal and liver lipid contents in mice.** (A) Retinal abundance of total saturated fatty acids (SFAs), total monounsaturated fatty acids (MUFAs) and total polyunsaturated fatty acids (PUFAs) relative to total fatty methyl esters (FAMES; 100%). (B) Heat map showing the retinal abundance of individual fatty acids relative to total FAMES (100%). Each column corresponds to the retina of one mouse. (C) Ratio of total n-6 PUFAs/total n-3 PUFAs in retinas. (D) Ratio of arachidonic acid (ARA, C20:4n-6)/docosahexaenoic acid (DHA, C22:6n-3) in retinas. (E) Fold increase in retinal DHA. The retinal amount of DHA of *ad libitum*-fed mice was defined as 1.0. (F) Ratio of ARA/DHA in livers. (G) Fold increase in liver DHA. The liver amount of DHA of *ad libitum*-fed mice was defined as 1.0. (H) Ratio of total FAMES/total dimethylacetals (DMAs) in retinas. (I) Heat map showing the retinal abundance of individual DMA species relative to total FAMES + total DMAs. Each column corresponds to the retina of one individual. Kruskal-Wallis test with Dunn's test for multiple comparisons. Significance: <sup>a</sup> *ad libitum* versus ITAD, <sup>b</sup> *ad libitum* versus *ad libitum* + *L. casei*, <sup>c</sup> *ad libitum* versus ITAD + *L. casei*, <sup>d</sup> ITAD versus *ad libitum* + *L. casei*, <sup>e</sup> ITAD versus ITAD + *L. casei*, and <sup>f</sup> *ad libitum* + *L. casei* versus ITAD + *L. casei*. \*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ . (C-H) Data are presented as mean  $\pm$  SEM.  $n = 8$  per group.

An excessive amount of n-6 PUFAs and a high n-6/n-3 ratio are promoting factors for many diseases including AMD [1]. We observed that *L. casei* supplementation alone or in combination with ITAD feeding significantly reduced the n-6/n-3 ratio in mouse retinas as well as that of ARA/DHA, which are the most abundant retinal n-6 and n-3 PUFAs (Fig. 2C and 2D). This phenotype is likely to be the consequence of the strong enrichment in DHA whose abundance was doubled in ITAD-fed mice supplemented with *L. casei* (Fig. 2E). Interestingly, this enrichment was higher than those reported in rodents supplemented with dietary DHA and/or its precursor, eicosapentaenoic acid (EPA) [9]. It may result from endogenous synthesis since the ARA/DHA ratio was increased and the amount of DHA was unchanged in the liver of ITAD-fed mice supplemented with *L. casei* (Fig. 2F and 2G).

In conclusion, this study provides evidence that time-restricted feeding combined to probiotics is a promising strategy to modulate retinal DHA content, and thus to sustain retinal homeostasis during aging. Further investigations are needed to decipher the molecular mechanisms involved in the microbiota-retina dialogue, as well as to evaluate the efficacy of such a strategy in protecting the retina against pathophysiological mechanisms related to AMD.

## Acknowledgements

We wish to thank the staff of the INRAE Infectiology of Fishes and Rodents Facility (IERP-UE907, Jouy-en-Josas Research Center, France) where animal experiments have been performed. IERP Facility belongs to the National Distributed Research Infrastructure for the Control of Animal and Zoonotic Emerging Infectious Diseases through *In Vivo* Investigation. This work was funded by Agence Nationale de la Recherche [ANR-11-LABX-0021-01]; Conseil Régional de Bourgogne, Franche-Comté [PARI grant]; FEDER (European Regional Development Fund).

## Supplementary Materials

The Supplementary data can be found online at: [www.aginganddisease.org/EN/10.14336/AD.2023.0324](http://www.aginganddisease.org/EN/10.14336/AD.2023.0324).

## Data and materials availability

For microbiota, the raw datasets are available in the Sequence Read Archive (SRA) database system under project accession number PRJNA927267. All other data supporting the findings reported herein are available on reasonable request to the corresponding author.

## References

- [1] van Leeuwen EM, Emri E, Merle BMJ, Colijn JM, Kersten E, Cougnard-Gregoire A, et al. (2018). A new perspective on lipid research in age-related macular degeneration. *Prog Retin Eye Res*, 67:56-86.
- [2] Bringer MA, Gabrielle PH, Bron AM, Creuzot-Garcher C, Acar N (2022). The gut microbiota in retinal diseases. *Exp Eye Res*, 214:108867.
- [3] Chaix A, Zarrinpar A, Miu P, Panda S (2014). Time-restricted feeding is a preventative and therapeutic intervention against diverse nutritional challenges. *Cell Metab*, 20:991-1005.
- [4] Martinez-Lopez N, Tarabra E, Toledo M, Garcia-Macia M, Sahu S, Coletto L, et al. (2017). System-wide Benefits of Intermeal Fasting by Autophagy. *Cell Metab*, 26:856-871 e855.
- [5] Precup G, Vodnar DC (2019). Gut Prevotella as a possible biomarker of diet and its eubiotic versus dysbiotic roles: a comprehensive literature review. *Br J Nutr*, 122:131-140.
- [6] Lebeer S, Vanderleyden J, De Keersmaecker SC (2008). Genes and molecules of lactobacilli supporting probiotic action. *Microbiol Mol Biol Rev*, 72:728-764, Table of Contents.
- [7] Selma MV, Beltran D, Luna MC, Romo-Vaquero M, Garcia-Villalba R, Mira A, et al. (2017). Isolation of Human Intestinal Bacteria Capable of Producing the Bioactive Metabolite Isovalinol from Ellagic Acid. *Front Microbiol*, 8:1521.
- [8] Saab S, Mazzocco J, Creuzot-Garcher CP, Bron AM, Bretillon L, Acar N (2014). Plasmalogens in the retina:

from occurrence in retinal cell membranes to potential involvement in pathophysiology of retinal diseases. *Biochimie*, 107 Pt A:58-65.

- [9] Schnebelen C, Gregoire S, Pasquis B, Joffre C, Creuzot-Garcher CP, Bron AM, et al. (2009). Dietary n-3 and n-6

PUFA enhance DHA incorporation in retinal phospholipids without affecting PGE(1) and PGE (2) levels. *Lipids*, 44:465-470.