



HAL
open science

Monitoring status and trends in genetic diversity for the Convention on Biological Diversity: An ongoing assessment of genetic indicators in nine countries

Sean Hoban, Jessica M da Silva, Alicia Mastretta-yanes, Catherine E
Grueber, Myriam Heuertz, Margaret E Hunter, Joachim Mergeay, Ivan
Paz-vinas, Keiichi Fukaya, Fumiko Ishihama, et al.

► To cite this version:

Sean Hoban, Jessica M da Silva, Alicia Mastretta-yanes, Catherine E Grueber, Myriam Heuertz, et al.. Monitoring status and trends in genetic diversity for the Convention on Biological Diversity: An ongoing assessment of genetic indicators in nine countries. Conservation Letters, 2023, 10.1111/conl.12953 . hal-04107368

HAL Id: hal-04107368

<https://hal.inrae.fr/hal-04107368>

Submitted on 26 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

PERSPECTIVE

Monitoring status and trends in genetic diversity for the Convention on Biological Diversity: An ongoing assessment of genetic indicators in nine countries

Sean Hoban^{1,2}  | Jessica M. da Silva^{3,4} | Alicia Mastretta-Yanes^{5,6} | Catherine E. Grueber⁷  | Myriam Heuertz⁸  | Margaret E. Hunter⁹ | Joachim Mergeay^{10,11} | Ivan Paz-Vinas¹²  | Keiichi Fukaya¹³ | Fumiko Ishihama¹³ | Rebecca Jordan¹⁴ | Viktoria Köppä¹⁵ | María Camilla Latorre-Cárdenas¹⁶ | Anna J. MacDonald¹⁷  | Victor Rincon-Parra¹⁸ | Per Sjögren-Gulve¹⁹  | Naoki Tani^{20,21} | Henrik Thurfjell²² | Linda Laikre¹⁵ 

¹The Morton Arboretum, Center for Tree Science, Lisle, Illinois

²The University of Chicago, Chicago, Illinois

³Foundational Biodiversity Sciences Division, South African National Biodiversity Institute, Pretoria, South Africa

⁴Department of Zoology, Centre for Ecological Genomics and Wildlife Conservation, University of Johannesburg, Johannesburg, South Africa

⁵Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, Ciudad de Mexico, Mexico

⁶Consejo Nacional de Ciencia y Tecnología, Ciudad de Mexico, Mexico

⁷Faculty of Science School of Life and Environmental Sciences, The University of Sydney, NSW, Australia

⁸INRAE, University of Bordeaux, Biogeco, Cestas, France

⁹U.S. Geological Survey, Wetland and Aquatic Research Center, Gainesville, Florida

¹⁰Research Institute for Nature and Forest, Geraardsbergen, Belgium

¹¹KU Leuven, Ecology, Evolution and Biodiversity Conservation, Leuven, Belgium

¹²Department of Biology, Colorado State University, Fort Collins, Colorado

¹³National Institute for Environmental Studies, Tsukuba, Ibaraki, Japan

¹⁴CSIRO, Sandy Bay, Tasmania, Australia

¹⁵Division of Population Genetics, Department of Zoology, Stockholm University, Stockholm, Sweden

¹⁶Instituto de Investigaciones en Ecosistemas y Sustentabilidad, Universidad Nacional Autónoma de México, Morelia, México

¹⁷Australian Antarctic Division, Department of Climate Change, Energy, the Environment and Water, Kingston, Tasmania, Australia

¹⁸Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, Programa de evaluación y monitoreo, Bogotá, Colombia

¹⁹Nordic Chapter of the Society for Conservation Biology, Uppsala, Sweden

²⁰Forestry Division, Japan International Research Center for Agricultural Sciences, Tsukuba, Ibaraki, Japan

²¹Faculty of Life and Environmental Sciences, University of Tsukuba, Tsukuba, Ibaraki, Japan

²²Swedish Species Information Centre, Swedish University of Agricultural Sciences, Uppsala, Sweden

Jessica M. da Silva and Alicia Mastretta-Yanes contributed equally.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. Conservation Letters published by Wiley Periodicals LLC.

Correspondence

Sean Hoban, The Morton Arboretum, Center for Tree Science, Lisle, Illinois; The University of Chicago, Chicago, Illinois
Email: shoban@mortonarb.org

Linda Laikre, Division of Population Genetics, Department of Zoology, Stockholm University, SE10691 Stockholm, Sweden.
Email: linda.laikre@popgen.su.se

Funding information

European Commission, Grant/Award Number: CA18134; COST (European Cooperation in Science and Technology); Formas, Grant/Award Number: 2020-01290; VR, Grant/Award Number: 2019-05503; the U.S. Geological Survey Powell Center for Synthesis and Analysis

Abstract

Recent scientific evidence shows that genetic diversity must be maintained, managed, and monitored to protect biodiversity and nature's contributions to people. Three genetic diversity indicators, two of which do not require DNA-based assessment, have been proposed for reporting to the Convention on Biological Diversity and other conservation and policy initiatives. These indicators allow an approximation of the status and trends of genetic diversity to inform policy, using existing demographic and geographic information. Application of these indicators has been initiated and here we describe ongoing efforts in calculating these indicators with examples. We specifically describe a project underway to apply these indicators in nine countries, provide example calculations, address concerns of policy makers and implementation challenges, and describe a roadmap for further development and deployment, incorporating feedback from the broader community. We also present guidance documents and data collection tools for calculating indicators. We demonstrate that Parties can successfully and cost-effectively report these genetic diversity indicators with existing biodiversity observation data, and, in doing so, better conserve the Earth's biodiversity.

KEYWORDS

adaptive capacity, conservation genetics, indicators, monitoring, policy, resilience

1 | REPORTING GENETIC DIVERSITY CHANGE IS VITAL AND IS FEASIBLE

Genetic diversity is variation at the DNA level, including differences among individuals and populations of each species. Because it contributes to the traits and survival of organisms, this intraspecific diversity is vital for helping species adapt to changing environments, including climate, pests, habitat changes, and disease. It also contributes to the stability and resilience of ecosystems including after extreme climate events (Raffard et al., 2019; Wernberg et al., 2018) and helps ensure successful ecological restoration (Breed et al., 2019), as has been documented in forest, grassland, streams, coral, and seagrass ecosystems (Booth & Grime, 2003; Des Roches et al., 2021; Ehlers et al., 2008). Unfortunately, genetic diversity has already declined substantially due mostly to habitat loss, habitat fragmentation, over-harvest, and other human activities (Exposito-Alonso et al., 2021; Leigh et al., 2019).

Although it is recognized as one of three basic levels of biodiversity (along with species and ecosystems), genetic diversity has been neglected in public policy and management (Hoban et al., 2020; Laikre et al., 2010; Laikre, 2010). Vague or imprecise wording in policy language, insufficient indicators to track progress, expensive technology, and metrics that are inaccessible to policy makers and

conservationists have resulted in insufficient actions and minimal reporting on genetic diversity status and trends (Hoban et al., 2013; Laikre et al., 2020; Taylor et al., 2017; Vernesi et al., 2008). For instance, a recent analysis of 57 national biodiversity reports revealed that countries primarily use indicators which are not well connected to genetic erosion, primarily focus on breeds of economically important species or crop wild relatives, neglect genetic monitoring, and focus on ex situ rather than in situ gene conservation (Hoban et al., 2021). Clear policy on genetic diversity conservation is also hampered by complicated concepts and terminology, a disconnect between genetic statistics and practical conservation actions, and a need for more meaningful engagement between genetic scientists and decision makers.

It is a critical time to conserve genetic diversity, particularly through the United Nations Convention on Biological Diversity (CBD) post-2020 Kunming-Montreal Global Biodiversity Framework (GBF), which was approved in December 2022 at COP15 (15th Conference of the Parties). During the next decade, Parties to the CBD will be legally required to report on their GBF progress using the Kunming-Montreal indicators. Indicators calculated at national scales and disaggregated by taxonomic groups, habitats or other categories also help countries understand and mitigate biodiversity loss. Through in person

and virtual engagement with CBD negotiations over the past two years, geneticists worked to help improve GBF Goals and Targets to avoid confusing, unclear, or indecisive wording (Hoban et al., 2023). The final Goal A states "...genetic diversity within populations of wild and domesticated species, is maintained, safeguarding their adaptive potential," and Target 4 states "...maintain and restore the genetic diversity within and between populations of native, wild and domesticated species to maintain their adaptive potential, including through in situ and ex situ conservation and sustainable management practices."

Safeguarding genetic diversity is feasible based on the finalized CBD Goals and Targets, but will depend on affordable, accessible, and relevant genetic diversity indicators (Frankham, 2022; Hoban et al., 2021). During COP-15 negotiations, many Parties mentioned they will need support in understanding and implementing indicators in practice, especially in developing countries.

Indicators of genetic diversity also provide useful measures of biodiversity for conservation and management efforts beyond the CBD. For instance, progress under national and international endangered species legislation (e.g., the U.S. Endangered Species Act and similar legislation in South Africa, Australia, and the European Union) is typically measured by numbers of species delisted. Indicators of genetic vulnerability within and among populations can provide finer assessment of changes in species' status, potentially improved allocation of resources, and targeted management. Similarly, simple indicators of genetic status may be useful for government and nongovernment organizations when communicating to the public about conservation threats to flagship species. Lastly, genetic diversity indicators highlight the importance of local populations for adaptive potential and resilience, which provides empowerment and leverage for local communities and indigenous peoples.

In this paper, we summarize significant advancements in indicators of genetic diversity that were ultimately recommended by the CBD at COP15 to assess status and trends in genetic diversity, and their application at national scales. Specifically, we

- reiterate the need for and purpose of three indicators, two of which do not require DNA-based analysis.
- summarize and address concerns from policy makers.
- describe ongoing deployment of indicators in 9 countries on 6 continents.
- describe indicator calculation, including with examples.
- address other challenges and describe a roadmap for uptake and use of genetic diversity indicators, including current and future support resources.




FIGURE 1 Example of genetic diversity indicators, for four hypothetical populations in Illinois, USA. Colors (red, green, yellow) indicate genetic differences among populations while shades within each color indicate genetic variations within each population. One tree = 1000 plants (five trees = 5000 plants). In 2020, 2 of 3 extant populations' census size is < 5000 ($N_e < 500$ considering an effective to census size ratio of $N_e/N_c = 0.1$) and thus too small to maintain genetic diversity (indicator 1). Three of four historic populations are maintained (indicator 2). DNA-based methods have been used to monitor genetic diversity in two populations (indicator 3—a value of 1 means that the species is monitored with DNA-based methods). DNA and tree images used are in public domain from ctker.com.

We are confident that nations and subnational actors can successfully report these genetic diversity indicators, and in doing so, better conserve the world's biodiversity.

Game et al. (2015) emphasize that policy-relevant science should clearly identify "actors who could take policy or management action." Our audience for this article is policy makers at international and national levels, agency personnel, and scientists or practitioners engaged in determining relative vulnerability of species—stakeholders who work toward allocating resources among such species, determining baselines for restoration, and outlining action plans for recovery.

2 | NEED AND PURPOSE FOR EACH INDICATOR

In 2020, three genetic diversity indicators were proposed and discussed (Figure 1) (Frankham, 2022; Hoban et al., 2020; Laikre et al., 2020). The indicators have several motivations: to assess or approximate genetic diversity status

without requiring new DNA data; to be affordable and feasible with existing data and with limited time investment; to use simple calculations; to allow for easy translation into policy and management of species; and to be applicable and relevant in all countries, taxonomic groups, and ecosystems. It is also desirable to use concepts that are intuitive or accessible to nongeneticists (e.g., genetic losses due to small populations and loss of populations). Assessing genetic status without DNA-based genetic data is vital since relatively few species have DNA-based research, especially in biodiversity hotspots. With these attributes, the indicators meet the first of three criteria for policy relevance outlined by Game et al. (2015)—salience or relevancy.

The proposed indicators of genetic diversity relate to central conservation genetics concepts:

1. conserving genetic diversity within large populations for rapid response to changing environmental conditions,
2. conserving genetic diversity among populations to provide diverse “options” for the future adaptation of the species,
3. assessing genetic data directly to guide conservation actions and sustainable use.

Indicator 1 (adopted as a Headline indicator A.4 for Goal A and Target 4 of the monitoring framework for the CBD Kunming-Montreal GBF) is “the proportion of populations within species with a genetically effective size $N_e > 500$.” Effective population size (N_e) is a concept that quantifies the rate of genetic erosion within a population; genetic erosion is the loss of genetic diversity, increase of inbreeding, and reduction of population ability to adapt. Past CBD indicators do not reflect genetic erosion within populations—though they did reflect genetic erosion in agricultural breeds (Hoban et al., 2020). Large N_e can help avoid population's and species' extinction, and supports high genetic diversity. (Genetic diversity statistics, such as “nucleotide diversity,” are directly calculable from N_e and from how often changes in the DNA are known to occur [“mutation rate”].) Specifically, contemporary effective size of 500 is well regarded as a minimum threshold for populations to maintain genetic diversity (Frankham et al., 2013; Franklin, 1980) (though see Frankham et al., 2013). Using a ratio of effective to census size, N_e/N_c (0.1 to 0.3 for most species; Hoban et al., 2021) translates to comparing census population size (N_c) to a threshold of about 5000. In this way, demographic information—census size—is translated to information on genetic status. For example, *Cupressus abramsiana*, a USA endemic gymnosperm, has five populations, and two exceed a census of 5000. Indicator 1 for this species is 0.4 (2 of 5). We caution that

past population size changes can impact N_e and N_e is not always correlated to N_c ; past demographic changes, as well as population connectivity, should be considered when interpreting genetic-based N_e .

Indicator 2 (adopted as complementary indicator for Goal A and Target 4 of the monitoring framework for the CBD Kunming-Montreal GBF) is “the proportion of populations maintained within species.” This indicator is needed because past CBD indicators do not reflect loss of genetic distinctiveness of each population. Each population may hold unique traits and genetic adaptations supporting species' survival (Karell et al., 2011; Palumbi et al., 2014). This concept is already recognized in distinct agricultural breeds, which are analogous to populations, each with unique traits or characters. Genetic, geographic and ecological variation allows future options for adaptation. Losses of species' populations and geographic range change are often quantifiable (Ceballos et al., 2017; Jetz et al., 2012). For example, *Capensibufo rosei* (Rose's mountain toadlet), endemic to the Cape Peninsula of South Africa, is known from five historical populations; only one of which exists today, along with a newly discovered population. Indicator 2 for this species is 0.33 (2 of 6 populations maintained). A caution with this indicator is that determining the baseline of historic range can be difficult and/or contentious. Where good information exists or can be estimated, this indicator estimates how much unique genetic diversity, available for adaptation, has been maintained or lost.

Indicator 3 (not currently an indicator of the monitoring framework) is “the number of species (and populations) monitored using DNA-based methods.” Although not common in all countries, DNA-based genetic monitoring, when available, can guide conservation actions and policy (Posledovich et al., 2021; Schwartz et al., 2007). For example, DNA-based monitoring of mountain pygmy possums (*Burrhamys parvus*) revealed inbreeding, very low N_e of ~ 10 and loss of over 80% of the population's genetic diversity. Based on this knowledge, a genetic rescue program introduced six genetically healthy males from a nearby population. This binary indicator counts any monitoring program using DNA data to help managers assess genetic status and choose appropriate actions, including studies of genetic connectivity, hybridization, adaptation, etc. Indicator 3 for this species is 1. By emphasizing the importance of conservation genetic studies, and quantifying them, as a complement to the proxy-based indicators, this indicator could incentivize more countries to start implementing DNA-based monitoring; conservation genetic studies should work with stakeholders to ensure the data can inform management.

These indicators more directly assess genetic erosion than the Red List Index or Living Planet Index. Neither

Index focuses on the threshold of N_e 500 within populations; below N_e 500, genetic erosion is exponentially faster. Also, the Red List Index typically assesses the entire species, which may be safe from extinction but still suffer important losses of genetic diversity; wide-ranging species may lose entire populations, and their unique genetic diversity, without affecting Red List status. Meanwhile the Living Planet Index primarily focuses on vertebrate species and does not consider losses before the 1970s, though many populations were lost or declined prior to this time.

3 | ADDRESSING CONCERNS FROM POLICY MAKERS

The Coalition for Conservation Genetics (Kershaw et al., 2022) presented these indicators to policy and management personnel globally through 10+ webinars and attendance at CBD meetings such as the Subsidiary Body on Science Technology and Technical Advice, SBSTTA (Supplemental Material). Five concerns about indicator uptake were common (Table 1): (1) necessity of DNA data, (2) feasibility of obtaining sufficient amounts of data, (3) realistic limitations on time, (4) limitations on skills or knowledge, and (5) concerns over data sharing, particularly Digital Sequence Information (DSI) (Scholz et al., 2022).

We address these concerns in Table 1, highlighting that: no new DNA collection is needed, most countries have sufficient data for reporting on these indicators, a small team can complete analysis in a fraction of one year without specialized expertise, and no DSI is shared in indicator reporting. Preliminary results of this assessment were presented to Parties in a side-event of the COP-15. By addressing these concerns through a dialogue with stakeholders, the indicators better meet the legitimacy criteria outlined by Game et al. (2015).

In the remainder of this article, we describe the transparent process of further developing the indicators, and address another of Game et al. (2015) credibility criteria.

4 | TOWARD INDICATOR DEPLOYMENT

While the indicators were being discussed by the CBD (CBD/WG2020/3/5, CBD/ID/OM/2022/1/2), a trial was initiated by the Swedish Environmental Protection Agency. Examining Swedish National Red List assessments, this trial (1) assessed data quality/ availability for 22,000 species and (2) calculated indicator values for 79 species (Thurfjell et al., 2022). Approximately 33% of species had census size estimates and 20% of species had historic and modern population information—data which

are the basis for indicators 1 and 2. The trial also found that 70% of herptile and 49% of mammal populations likely had $N_e > 500$ while 32% of herptiles and 84% of mammals are maintaining their geographic range. This study concluded that

- Data for the indicators are likely available for thousands of species.
- A large proportion of species are already experiencing genetic erosion within and among populations.

A new pilot project described here is implementing the indicators in nine countries across six continents, in order to:

1. create and refine a standard workflow, definitions, methodology, and data collection device, and
2. evaluate > 100 species for each country to:
3. determine how many species have necessary data to calculate indicators,
4. when possible, extract data and perform indicator calculations, and
5. identify challenges so guidance and calculations can be improved for use by more countries.

We describe the approach to demonstrate that data are available to calculate the indicators in countries around the world, and to show that data collation is practical, achievable and adaptable.

Gathering data on populations of species can be challenging because there is no global, standard database of population census size or changes in populations. However, population level data of many species are available in different reports, atlases, and databases, and with local and expert knowledge holders.

We identified three approaches to gathering data, which suit different countries' needs and can be used in combination (Figure 2).

- “Manual data extraction” involves reviewing national or subnational documents (management or recovery plans, status assessments, environmental impact reports), scientific literature, country flora or natural history descriptions, IUCN Red List assessment or NatureServe assessment text, etc.
- “Expert consultation” involves facilitated discussion and knowledge sharing among local experts and traditional knowledge holders with firsthand (but perhaps unpublished) information—similar to Red List assessment workshops which elicit quantitative information on species.
- “Automated data extraction” from existing databases on national surveys, where species' occurrences or pop-

TABLE 1 Resolving common concerns regarding genetic diversity indicators

Concerns	Solutions or clarifications
<p><i>Data needed</i></p> <p>Is genetic data (DNA-based analysis) needed?</p>	<p>Indicators 1 and 2 do not need genetic data (DNA sequencing), genetic expertise, or laboratory facilities to generate data.</p> <p>Genetic data can be used to estimate N_e, but <i>in most cases existing census size data</i> (counts of individuals using camera traps, surveys, etc.) along with an N_e/N_c ratio can be applied to obtain a valid proxy of N_e, and similar records for measuring “populations maintained.”</p>
<p><i>Achievable</i></p> <p>Is there enough census or geographic data?</p>	<p>Yes, many countries monitor priority species, maintain biodiversity databases or citizen science tools, and contribute Red List assessments which may have geographic and census data.</p> <p>Data collection can include local knowledge or expert consultation, or categorical or imperfect data.</p> <p>Genetic diversity indicator reporting is not necessary for all species; it would be for a relatively small, representative sample of species per country (we recommend ≥ 100) (Frankham, 2022; Hoban et al., 2021; Laikre et al., 2021). Representative = from a diversity of ecosystems, taxonomic groups, rarity, and lifespan</p>
<p><i>Realistic</i></p> <p>Is indicator calculation too time-consuming?</p>	<p>Collection of census sizes and number of populations is similar in scope to compiling other information for CBD National Reports. From pilot tests, 3–5 persons could complete analyses on a total of 100 species in 2–4 weeks, or 1 person could do it in 2–4 months per country.</p>
<p><i>Realistic</i></p> <p>Does indicator calculation require special skills or tools?</p>	<p>No. An undergraduate biology education, and short training sessions, should provide sufficient foundation for gathering data. Our guidance documents explain how to choose representative species, how to find and record data, and how to resolve challenges (Supplemental Material; see also Hvilsom et al., 2022). A data collection device allows for standard recording and storage of data, and analysis (Supplemental Material).</p>
<p><i>Data sharing</i></p> <p>Does reporting genetic indicators require reporting or sharing genetic data or Digital Sequence Information (DSI)?</p>	<p>Reporting on genetic indicators does not involve sharing or reporting any DSI. Indicators 1 and 2 will typically be based on demographic and geographic data (e.g., census sizes, population distributions), which do not use DNA data (DSI) at all. If DNA data were used to calculate N_e or define population boundaries, only the count of populations and estimates of effective size would be recorded; <i>no DSI is reported or shared</i>. Even raw census data and population data could be retained by the country, reporting only the proportions for the indicators</p> <p>Although indicator 3 does assess genetic studies in the country, it is only a count of studies. It does not assess DNA data or share DSI. Indicating DNA data availability is at the discretion of each country (Scholz et al., 2022).</p>

ulation sizes are regularly assessed and stored, often in a gridded spatial database (common in some fish, plants, birds, and mammals), as with stock assessments, some National Red List databases, or forest inventories. Population presence can also be obtained from citizen science databases (e.g., iNaturalist). Automated analysis can compare census sizes per population to the N_e (or N_c) threshold, and compare historic data, atlases, or range maps to current population presence or projections of habitat change to assess “populations maintained.”

Examples from different countries illustrate the diverse options available (Table 2). Recovery plans for dozens to thousands of threatened species are mandated by national legislation (e.g., Australia—the Environment Protection

and Biodiversity Conservation Act 1999, <https://www.dceew.gov.au/environment/epbc>; South Africa—Biodiversity Management Plans, https://www.dffe.gov.za/content/management_plans/biodiversity; USA—the Endangered Species Act, <https://www.fws.gov/law/endangered-species-act>). These documents typically detail species biology and demographic status. In Japan, many threatened vascular plants have been surveyed for census size for over two decades by the Japanese Society for Plant Taxonomy, while for common trees, statistical estimates for population size (Fukaya et al., 2020) were estimated from vegetation survey data. In Mexico, taxonomic experts who recently helped validate distribution models for crop wild relatives will be consulted for indicator values (Goettsch et al., 2021). In France, Belgium, and Sweden, much biodiversity data




FIGURE 2 Process of assessing indicators for a set of species, using data from different sources (manual, expert, automated). Indicators 1 and 2 are primarily obtained from demographic and geographic information. Note that genetic information is optional for indicators 1 and 2 as shown in grey dashed lines. For indicator 1, census sizes (N_c) can be converted to effective sizes (N_e) by applying a species-specific effective to census size ratio, and/or a ratio of $N_e/N_c = 0.1$. Indicator assessment leads to reporting, policy outcomes, and management action.

from experts and diverse sources are collected in easy to access web-based portals (France—<https://inpn.mnhn.fr/>; Belgium—<https://www.biodiversity.be/1767>; Sweden—<https://www.artdatabanken.se/en/>). In Colombia, the Biodiversity Information System (SIB) repository compiles species surveys from throughout the country (<https://biodiversidad.co/>), which is mandated by many public and private organizations. These data are reviewed by national experts for validation and used to create freely available species distribution models (<http://biomodelos.humboldt.org.co/>) and for conservation prioritization.

5 | CALCULATING INDICATOR VALUES AT NATIONAL LEVELS

Genetic indicator calculation is straightforward. First, indicators are calculated for each species. When data are available as a range (e.g., N_c is 10,000 to 20,000), the mean

is used. Qualitative data such as “a few hundred” or “at least 5000” are often sufficient for comparison to the N_c 5000 threshold. In the case of differing estimates from multiple sources, either the most recent source is used, or a mean of values based on the different sources is taken (see Table 3). We add that in most practical applications, indicator 1 will typically be based on an N_c estimate and an N_e/N_c ratio, but N_e can be calculated directly from DNA-based genetic studies. In such cases the contemporary N_e value should be used, and reported with confidence intervals.

While indicator 3 is a simple sum (species for which one or more populations are monitored using DNA), the country indicator value for indicators 1 and 2 is the mean of values across species (a median could be used for skewed distributions). If taxonomic groups are not represented evenly, the indicator value is the mean of each taxonomic group’s means, which down-weights overly represented taxonomic groups (e.g., mammals). Additionally, each species can be weighted by the proportion of its geographic

TABLE 2 Countries participating in testing of genetic diversity indicators, showcasing variation in overall approach

Country	Number of people	Taxonomic groups*	Approach	Sources**
South Africa	7 or 8	B, F, H, I, M, Ma, P	Manual + Expert	AH, EGI, NRL, GRL, SMP
USA	10	B, F, H, I, M, P	Manual	AH, GRL, SMP
Japan	4 or 5	P	Manual + Auto	AH, FG, ND, NRL
Mexico	10	B, CW, F, H, M, P	Expert + Auto	AH, EGI, ND, NRL
Australia	8	B, F, H, I, M, P	Manual	AH, GRL, ND, SMP
Sweden	2+	B, F, H, I, M, P	Manual + Expert	AH, ND, NRL, SMP
Belgium	2+	B, I, M, P	Manual + Expert	AH, ND, NRL, SMP
Colombia	3+	B, H, M, P	Manual + Expert	AH, EGI, GRL, ND, NRL
France	4	B, F, H, I, M, P	Manual	AH, ND, SMP, NRL

*B = Birds, CW = crop wild relatives, F = freshwater fish, H = herptiles, I = invertebrates, M = mammals, Ma = marine, P = plants.

**AH = ad hoc/ other (websites, email experts, scientific literature), EGI = expert group input, FG = field guide, GRL = Global Red List, ND = national data set or database of species information and/or occurrences, NRL = National Red List, SMP = species-specific management, action or recovery plan.

TABLE 3 Example indicator values for select species

Species	Taxonomic group	Country	Indicator 1 (Ne)	Indicator 2 (populations)	Indicator 3 (studies)*
<i>Pelobates fuscus</i>	Amphibian	Belgium	0	0.33	1
<i>Rana arvalis</i>	Amphibian	Belgium	0.18	0.84	1
<i>Angelica heterocarpa</i>	Angiosperm	France	0.5	1	1
<i>Zingel asper</i>	Fish	France	0.2	1	1
<i>Tetrao urogallus</i>	Bird	France	0.33**	0.6	1
<i>Taraxacum yuparense</i>	Angiosperm	Japan	0	0.5	0
<i>Carex cinerascens</i>	Angiosperm	Japan	0.5	0.8	0
<i>Capensibufo rosei</i>	Amphibian	South Africa	0.5	0.33	1
<i>Clinus spatulatus</i>	Fish	South Africa	N/A***	1	0
<i>Syncerus caffer caffer</i>	Mammal	South Africa	0.3	1	1
<i>Alces alces</i>	Mammal	Sweden	0.67	1	1
<i>Silurus glanis</i>	Fish	Sweden	0	0.5	1
<i>Cupressus abramsiana</i>	Gymnosperm	USA	0.4	1	0
<i>Rana muscosa</i>	Amphibian	USA	0	0.76**	1
<i>Charadrius melodus</i>	Bird	USA	0.39**	1	1
<i>Erigeron maguirei</i>	Angiosperm	USA	0.5	1	0

*Indicator 3 is binary for each species, 0 or 1 (1 = one or more populations of the species is monitored with DNA-based method; 0 = no DNA-based monitoring for the species).

**Different reports suggest different values or different interpretations of population boundaries; these values are means of the different interpretations.

***Indicator 1 could not be calculated because no N_c or N_e data are available.

range in the country, from 0 to 1, to reflect national responsibility, with full weight for endemic species (Domisch et al., 2016). Transboundary populations can be weighted similarly.

Equation (1): simple indicator value (IV) mean across species (s)

$$\frac{\sum_{s=1}^{s=100} IV_s}{s}$$

Equation (2): indicator value (IV) weighted by proportion of geographic range in country (W)

$$\frac{\sum_{s=1}^{s=100} IV_s W_s}{\sum_{s=1}^{s=100} W_s}$$

Equation (3): indicator value (IV) giving equal weight to birds ($b = 20$ species), plants ($p = 30$ species), and

mammals ($m = 50$ species)

$$\frac{\sum_{b=1}^{b=20} IV}{b} + \frac{\sum_{p=1}^{p=30} IV}{p} + \frac{\sum_{m=1}^{m=50} IV}{m}$$

3

6 | OVERCOMING CHALLENGES

In trials of the methodology outlined above we have resolved several challenges: biases, uncertainty in data, and difficulty in defining populations.

Biases

Ideally, the 100+ species assessed by a country reflect diverse ecosystems, taxonomic groups, rarity categories, and life history traits (e.g., lifespan). In reality, biases exist due to a country's habitat types, biogeography, latitude, and investment priorities in biodiversity monitoring and thus data availability. The Red List Index and Living Planet Index also have such biases (Bachman et al., 2019; Puurtinen et al., 2022). Weighting by taxonomy or ecosystem can help address bias (previous section). Additionally, biases should be noted by displaying counts per category in a matrix, as is common in Red List summary tables.

Uncertainty or qualitative information

Sometimes, census sizes are recorded as a range of values rather than point values; using vague wording such as "several hundred"; or census at the species but not population level. Our draft guidance (Supplemental Material) provides the assessor with advice on translating these into quantitative information, and recording uncertainty.

In addition, for indicator 1, if census size is not available, it may be possible to use known occupied area multiplied by mean density (number of individuals supported per unit area) to estimate the number of individuals. This allows evaluating whether an area is capable of housing a population N_c larger than the threshold; if the area is smaller, the N_e will likely be smaller than 500 (Mergeay et al., 2020). For indicator 2, if information on the number of historic populations is not known, the assessor may record overall decline in area, which will ultimately result in lost populations and genetic diversity (Exposito-Alonso et al., 2021).

Definition of population boundaries (required for indicators 1 and 2)

Ecologists and geneticists have worked for decades to understand population distinctions (Jost et al., 2018;

Waples & Gaggiotti, 2006). For the indicators, available knowledge can be used to assess genetic distinction, typically less than one migrant per generation from other populations (Mills & Allendorf, 1996), such as using population designations from the report or experts consulted, which reflect knowledge of the species biology, history, and dispersal; discrete patches such as forest or lakes; ecoregions, geographic (and migration) barriers such as mountains/ valleys, or hydrological zones, which may promote local adaptation; phylogeographic studies (i.e., "proxies of genetic differentiation"; Tobón-Niedfeldt et al., 2022); or grid cells based on species' dispersal.

Other challenges remain and will require capacity building across the biodiversity conservation sector, including from the CBD, national agencies, NGO partners, scientists, and local monitoring initiatives. These challenges include significant time (at least several person months per country, see Table 1), translation of guidance and infrastructure to non-English languages, and continued scientific work to make sure indicators are appropriate in challenging situations such as small countries, highly managed species or species with large ex situ populations, and species with naturally small populations. However, as detailed in Section 7, we believe these challenges can be overcome.

7 | ROADMAP

We have presented the purpose and straightforward methodology of each indicator, addressed concerns of policy makers, showed that data are available and usable, and described ongoing work in nine countries. We demonstrated that genetic diversity indicators are ready for use in biodiversity conservation and reporting, with existing data.

Increased uptake of these indicators by Parties to the CBD and other users (environmental agencies, wildlife managers, national legislation, etc.) will require further successful demonstrations, published step-by-step workflows, and training workshops, ideally in multiple languages (Kershaw et al., 2022). We have created an online data collection form using Kobotoolbox (www.kobotoolbox.org/) and a guidance document (Supplemental Material, and <https://github.com/AliciaMstt/GeneticIndicators>) for anyone to use. Kobotoolbox is a free open source tool for data collection in a standardized structured format, with connections to analysis in Excel, R, Python or GIS software. Our GitHub encourages interactive engagement with stakeholders who can offer suggestions or ask questions.

Future development of online resources can enhance data storage, managing, and sharing. An online portal could accept and store submitted data such as N_e or N_c values, population designations, and references over multiple reporting cycles at global, national and subnational levels,

which would make compilation of the indicators easier and increase transparency. Other cyber infrastructure would be connections to resources like species distribution models (Jetz et al., 2012), which could help with defining populations, calculating N_e based on area and density, loss of distinct populations etc. Finally, as noted by others (Garner et al., 2020; Thurfjell et al., 2022), it would be advantageous for Red List assessment workshops to include gathering raw data and information needed for calculation of the indicators, including population boundaries. Red List assessors scour literature and consult experts for demographic and geographic information, and indicators could be recorded systematically in the Red List database.

Lastly, we note that the indicators presented here complement other useful indicators for tracking genetic diversity which other authors have suggested. For example, the “genetic scorecard” assesses genetic diversity threats such as hybridization and low reproduction (Hollingsworth et al., 2020), while another indicator assesses the extent to which geographic ranges are protected in situ or ex situ (Khoury et al., 2019). Additionally, indicators using direct assessments of genomic health based on DNA data are available for some species, and collection of DNA data for species management remains important (van Oosterhout, 2020; Andersson et al., 2022; O’Brien et al., 2022). In the future, genetic indicators could be synthesized together for comprehensive genetic assessment (Frankham, 2022).

8 | CLOSING REMARKS

The indicators support and enable CBD Goal A and Target 4 wording (Frankham, 2022; Hoban et al., 2020; Laikre et al., 2021).

We note that many Parties (and non-Parties) will also set National Targets, which are typically in line with, but may be more ambitious, than the global targets, and the setting of these targets offers further opportunity for scientists to work with policy makers.

We close by reiterating that scientific evidence shows that genetic diversity is a basic pillar of all biodiversity that must be *maintained (not lost)*, *protected (via legislation and policy)*, *managed (through interventions such as restoring gene flow and genetic rescue)*, and *monitored (with DNA-based and non-DNA-based metrics like the indicators)* (Des Roches et al., 2021; Hoban et al., 2021). This is necessary to enable all species to adapt to environmental change, ensure resilient ecosystems, and benefit humanity.

AUTHOR CONTRIBUTIONS

SH, JdS, AM-Y, and LL conceived the study. SH wrote the first draft and conceived Figure 1. All authors contributed

to discussions and revisions. IP-V conceived Figure 2. MH and AM-Y conceived Table 1.

ACKNOWLEDGMENTS

We thank Peter Galbusera, Irene Ramos, and Gernot Segelbacher for comments on the manuscript, and the Coalition for Conservation Genetics for general support. We thank GEO BON and the CBD Secretariat, especially Maria Cecilia Londoño, Andrew Gonzalez, and Katie Millette, for feedback and the opportunity to present indicators at policy webinars. We would like to dedicate this paper to our dear colleague, Dr Michael Bruford, who recently passed away. He was a driving force behind conservation genetic policy and supported and encouraged this work on genetic diversity indicators. We are grateful for his leadership, vision, and inspiration.

DISCLAIMER


Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

DATA AVAILABILITY STATEMENT

The only data for the article are in Table 2; no other data were used.


ORCID


Sean Hoban  <https://orcid.org/0000-0002-0348-8449>

Catherine E. Grueber  <https://orcid.org/0000-0002-8179-1822>

Myriam Heuertz  <https://orcid.org/0000-0002-6322-3645>

Ivan Paz-Vinas  <https://orcid.org/0000-0002-0043-9289>

Anna J. MacDonald  <https://orcid.org/0000-0003-2972-200X>

Per Sjögren-Gulve  <https://orcid.org/0000-0001-6584-128X>

Linda Laikre  <https://orcid.org/0000-0001-9286-3361>

REFERENCES

- Andersson, A., Karlsson, S., Ryman, N., & Laikre, L. (2022). Monitoring genetic diversity with new indicators applied to an alpine freshwater top predator. *Molecular Ecology*, 31, 6422–6439.
- Bachman, S. P., Field, R., Reader, T., Raimondo, D., Donaldson, J., Schatz, G. E., & Lughadha, E. N. (2019). Progress, challenges and opportunities for Red Listing. *Biological Conservation*, 234, 45–55.
- Booth, R. E., & Grime, J. P. (2003). Effects of genetic impoverishment on plant community diversity. *Journal of Ecology*, 91, 721–730.
- Breed, M. F., Harrison, P. A., Blyth, C., Byrne, M., Gaget, V., Gellie, N. J. C., Groom, S. V. C., Hodgson, R., Mills, J. G., Prowse, T. A. A., Steane, D. A., & Mohr, J. J. (2019). The potential of genomics for restoring ecosystems and biodiversity. *Nature Reviews Genetics*, 20, 615–628.
- Ceballos, G., Ehrlich, P. R., & Dirzo, R. (2017). Biological annihilation via the ongoing sixth mass extinction signaled by vertebrate population losses and declines. *PNAS*, 114, E6089–E6096.

- Des Roches, S., Pendleton, L. H., Shapiro, B., & Palkovacs, E. P. (2021). Conserving intraspecific variation for nature's contributions to people. *Nature Ecology & Evolution*, 5, 574–582.
- Domisch, S., Wilson, A. M., & Jetz, W. (2016). Model-based integration of observed and expert-based information for assessing the geographic and environmental distribution of freshwater species. *Ecography*, 39, 1078–1088.
- Ehlers, A., Worm, B., & Reusch, T. (2008). Importance of genetic diversity in eelgrass *Zostera marina* for its resilience to global warming. *Marine Ecology Progress Series*, 355, 1–7.
- Exposito-Alonso, M., Booker, T. R., Czech, L., Gillespie, L., Hateley, S., Kyriazis, C. C., Lang, P. L. M., Leventhal, L., Nogues-Bravo, D., Pagowski, V., Ruffley, M., Spence, J. P., Toro Arana, S. E., Weis, C. L., & Zess, E. (2021). Genetic diversity loss in the Anthropocene. *Science*, 377, 1431–1435.
- Frankham, R. (2022). Evaluation of proposed genetic goals and targets for the Convention on Biological Diversity. *Conservation Genetics*, 23, 865–870. <https://doi.org/10.1007/s10592-022-01459-1>
- Frankham, R., Brook, B. W., Bradshaw, C. J. A., Traill, L. W., & Spielman, D. (2013). 50/500 rule and minimum viable populations: Response to Jamieson and Allendorf. *Trends in Ecology & Evolution*, 28, 187–188.
- Franklin, I. R. (1980). Evolutionary change in small populations. Sinauer Associates.
- Fukaya, K., Kusumoto, B., Shiono, T., Fujinuma, J., & Kubota, Y. (2020). Integrating multiple sources of ecological data to unveil macroscale species abundance. *Nature Communications*, 11, 1695.
- Game, E. T., Schwartz, M. W., & Knight, A. T. (2015). Policy relevant conservation science. *Conservation Letters*, 8, 309–311.
- Garner, B. A., Hoban, S., & Luikart, G. (2020). IUCN Red List and the value of integrating genetics. *Conservation Genetics*, 21, 795–801.
- Goettsch, B., Urquiza-Haas, T., Koleff, P., Acevedo Gasman, F., Aguilar-Meléndez, A., Alavez, V., Alejandre-Iturbide, G., Aragón Cuevas, F., Azurúa Pérez, C., Carr, J. A., Castellanos-Morales, G., Cerén, G., Contreras-Toledo, A. R., Correa-Cano, M. E., De La Cruz Larios, L., Debouck, D. G., Delgado-Salinas, A., Gómez-Ruiz, E. P., González-Ledesma, M., ... Jenkins, R. K. B. (2021). Extinction risk of Mesoamerican crop wild relatives. *Plants, People, Planet*, 3, 775–795.
- Hoban, S., Bruford, M., D'urban Jackson, J., Lopes-Fernandes, M., Heuertz, M., Hohenlohe, P. A., Paz-Vinas, I., Sjögren-Gulve, P., Segelbacher, G., Vernesi, C., Aitken, S., Bertola, L. D., Bloomer, P., Breed, M., Rodríguez-Correa, H., Funk, W. C., Grueber, C. E., Hunter, M. E., Jaffe, R., ... Laikre, L. (2020). Genetic diversity targets and indicators in the CBD post-2020 Global Biodiversity Framework must be improved. *Biological Conservation*, 248, 108654.
- Hoban, S., Bruford, M. W., Funk, W. C., Galbusera, P., Griffith, M. P., Grueber, C. E., Heuertz, M., Hunter, M. E., Hvilsom, C., Stroil, B. K., Kershaw, F., Khoury, C. K., Laikre, L., Lopes-Fernandes, M., Macdonald, A. J., Mergéay, J., Meek, M., Mittan, C., Mukassabi, T. A., ... Vernesi, C. (2021). Global commitments to conserving and monitoring genetic diversity are now necessary and feasible. *Bioscience*, 71, 964–976.
- Hoban, S., Bruford, M. W., da Silva, J. M., Chris Funk, W., Frankham, R., Gill, M. J., Grueber, C. E., Heuertz, M., Hunter, M. E., Kershaw, F., Lacy, R. C., Lees, C., Lopes-Fernandes, M., MacDonald, A. J., Mastretta-Yanes, A., McGowan, P. J. K., Meek, M. H., Mergéay, J., Millette, K. L., ... Laikre, L. (2023). Genetic diversity goals and targets have improved, but remain insufficient for clear implementation of the post-2020 global biodiversity framework. *Conservation Genetics*, 1–11.
- Hoban, S., Campbell, C. D., da Silva, J. M., Ekblom, R., Funk, W. C., Garner, B. A., Godoy, J. A., Kershaw, F., Macdonald, A. J., Mergéay, J., Minter, M., O'Brien, D., Paz-Vinas, I., Pearson, S. K., Pérez-Espona, S., Potter, K. M., Russo, I.-R. M., Segelbacher, G., Vernesi, C., & Hunter, M. E. (2021). Genetic diversity is considered important but interpreted narrowly in country reports to the Convention on Biological Diversity: Current actions and indicators are insufficient. *Biological Conservation*, 261, 109233.
- Hoban, S., Paz-Vinas, I., Aitken, S., Bertola, L. D., Breed, M. F., Bruford, M. W., Funk, W. C., Grueber, C. E., Heuertz, M., Hohenlohe, P., Hunter, M. E., Jaffé, R., Fernandes, M. L., Mergéay, J., Moharrek, F., O'Brien, D., Segelbacher, G., Vernesi, C., Waits, L., & Laikre, L. (2021). Effective population size remains a suitable, pragmatic indicator of genetic diversity for all species, including forest trees. *Biological Conservation*, 253, 108906.
- Hoban, S. M., Hauffe, H. C., Pérez-Espona, S., Arntzen, J. W., Bertorelle, G., Bryja, J., Frith, K., Gaggiotti, O. E., Galbusera, P., Godoy, J. A., Hoelzel, A. R., Nichols, R. A., Primmer, C. R., Russo, I.-R., Segelbacher, G., Siegismund, H. R., Sihvonen, M., Vernesi, C., Vilà, C., & Bruford, M. W. (2013). Bringing genetic diversity to the forefront of conservation policy and management. *Conservation Genetics Resources*, 5, 593–598.
- Hollingsworth, P., O'Brien, D. R., Ennos, R., Yahr, R., Neaves, L., Ahrends, A., Ballingall, K. T., Brooker, R. W., Burke, T., Cavers, S., Dawson, I. K., Elston, D. A., Kerr, J., Marshall, D. F., Pakeman, R. J., Trivedi, C., Wall, E., Wright, F., & Ogden, R. (2020). *Scotland's biodiversity progress to 2020 Aichi targets: Conserving genetic diversity – development of a national approach for addressing Aichi Biodiversity Target 13 that includes wild species* (Vol. 32). Scottish Natural Heritage.
- Hvilsom, C., Ekblom, R., Fischer, M. C., & Segelbacher, G. (2022). Selecting species and populations for monitoring of genetic diversity. International Union for Conservation of Nature. <https://doi.org/10.2305/iucn.ch.2022.07.en>
- Jetz, W., McPherson, J. M., & Guralnick, R. P. (2012). Integrating biodiversity distribution knowledge: Toward a global map of life. *Trends in Ecology & Evolution*, 27, 151–159.
- Jost, L., Archer, F., Flanagan, S., Gaggiotti, O., Hoban, S., & Latch, E. (2018). Differentiation measures for conservation genetics. *Evolutionary Applications*, 11, 1139–1148.
- Karell, P., Ahola, K., Karstinen, T., Valkama, J., & Brommer, J. E. (2011). Climate change drives microevolution in a wild bird. *Nature Communications*, 2, 208.
- Kershaw, F., Bruford, M. W., Funk, W. C., Grueber, C. E., Hoban, S., Hunter, M. E., Laikre, L., Macdonald, A. J., Meek, M. H., Mittan, C., O'Brien, D., Ogden, R., Shaw, R. E., Vernesi, C., & Segelbacher, G. (2022). The Coalition for Conservation Genetics: Working across organizations to build capacity and achieve change in policy and practice. *Conservation Science and Practice*, 4, e12635.
- Khoury, C. K., Amariles, D., Soto, J. S., Diaz, M. V., Sotelo, S., Sosa, C. C., Ramírez-Villegas, J., Achicanoy, H. A., Velásquez-Tibatá, J., Guarino, L., León, B., Navarro-Racines, C., Castañeda-Álvarez, N. P., Dempewolf, H., Wiersema, J. H., & Jarvis, A. (2019). Comprehensiveness of conservation of useful wild plants: An operational indicator for biodiversity and sustainable development targets. *Ecological Indicators*, 98, 420–429.

- Laikre, L. (2010). Genetic diversity is overlooked in international conservation policy implementation. *Conservation Genetics*, *11*, 349–354.
- Laikre, L., Allendorf, F. W., Aroner, L. C., Baker, C. S., Gregovich, D. P., Hansen, M. M., Jackson, J. A., Kendall, K. C., Mckelvey, K., Neel, M. C., Olivieri, I., Ryman, N., Schwartz, M. K., Bull, R. S., Stetz, J. B., Tallmon, D. A., Taylor, B. L., Vojta, C. D., Waller, D. M., & Waples, R. S. (2010). Neglect of genetic diversity in implementation of the convention of biological diversity. *Conservation Biology*, *24*, 86–88.
- Laikre, L., Hoban, S., Bruford, M. W., Segelbacher, G., Allendorf, F. W., Gajardo, G., Rodríguez, A. G., Hedrick, P. W., Heuertz, M., Hohenlohe, P. A., Jaffé, R., Johannesson, K., Liggins, L., Macdonald, A. J., Orozcoerwengel, P., Reusch, T. B. H., Rodríguez-Correa, H., Russo, I.-R. M., Ryman, N., & Vernesi, C. (2020). Post-2020 goals overlook genetic diversity. *Science*, *367*, 1083–1085.
- Laikre, L., Hohenlohe, P. A., Allendorf, F. W., Bertola, L. D., Breed, M. F., Bruford, M. W., Funk, W. C., Gajardo, G., González-Rodríguez, A., Grueber, C. E., Hedrick, P. W., Heuertz, M., Hunter, M. E., Johannesson, K., Liggins, L., Macdonald, A. J., Mergeay, J., Moharrek, F., O'Brien, D., & Hoban, S. (2020). Authors' reply to letter to the editor: Continued improvement to genetic diversity indicator for CBD. *Conservation Genetics*, *22*, 533–536.
- Leigh, D. M., Hendry, A. P., Vázquez-Domínguez, E., & Friesen, V. L. (2019). Estimated six per cent loss of genetic variation in wild populations since the industrial revolution. *Evolutionary Applications*, *12*, 1505–1512.
- Mergeay, J., Felix, L., Neyrinck, S., & Cox, K. (2020). *Conservation Genetic Status of Moor Frog (Rana arvalis) in France*. Reports of the Research Institute for Nature and Forest 2020 (43). Research Institute for Nature and Forest, Brussels. <https://doi.org/10.21436/inbor.19362192>
- Mills, L. S., & Allendorf, F. W. (1996). The one-migrant-per-generation rule in conservation and management. *Conservation Biology*, *10*, 1509–1518.
- O'Brien, D., Laikre, L., Hoban, S., Bruford, M. W., Ekblom, R., Fischer, M. C., Hall, J., Hvilsom, C., Hollingsworth, P. M., Kershaw, F., Mittan, C. S., Makussabi, T. A., Ogden, R., Segelbacher, G., Shaw, R. E., Vernesi, C., & MacDonald, A. (2022). Bringing together approaches to reporting on within species genetic diversity. *Journal of Applied Ecology*, *59*, 2227–2233.
- Palumbi, S. R., Barshis, D. J., Traylor-Knowles, N., & Bay, R. A. (2014). Mechanisms of reef coral resistance to future climate change. *Science*, *344*, 895–898.
- Posledovich, D., Ekblom, R., & Laikre, L. (2021) Mapping and monitoring genetic diversity in Sweden: A proposal for species, methods and costs. (Report of the Swedish Environmental Protection Agency; No. 6959). Retrieved from <https://urn.kb.se/resolve?urn=nbn:se:naturvardsverket:diva-8964>
- Puurinen, M., Elo, M., & Kotiaho, J. S. (2022). The Living Planet Index does not measure abundance. *Nature*, *601*, E14–E15.
- R Taylor, H., Dussex, N., & Van Heezik, Y. (2017). Bridging the conservation genetics gap by identifying barriers to implementation for conservation practitioners. *Global Ecology and Conservation*, *10*, 231–242.
- Raffard, A., Santoul, F., Cucherousset, J., & Blanchet, S. (2019). The community and ecosystem consequences of intraspecific diversity: A meta-analysis. *Biological Reviews of the Cambridge Philosophical Society*, *94*, 648–661.
- Scholz, A. H., Freitag, J., Lyal, C. H. C., Sara, R., Cepeda, M. L., Cancio, I., Sett, S., Hufton, A. L., Abebaw, Y., Bansal, K., Benbouza, H., Boga, H. I., Brisse, S., Bruford, M. W., Clissold, H., Cochrane, G., Coddington, J. A., Deletoille, A.-C., García-Cardona, F., ... Overmann, J. (2022). Multilateral benefit-sharing from digital sequence information will support both science and biodiversity conservation. *Nature Communications*, *13*, 1086.
- Schwartz, M., Luikart, G., & Waples, R. (2007). Genetic monitoring as a promising tool for conservation and management. *Trends in Ecology & Evolution*, *22*, 25–33.
- Thurfjell, H., Laikre, L., Ekblom, R., Hoban, S., & Sjögren-Gulve, P. (2022). Practical application of indicators for genetic diversity in CBD post-2020 Global Biodiversity Framework implementation. *Ecological Indicators*, *142*, 109167.
- Tobón-Niedfeldt, W., Mastretta-Yanes, A., Urquiza-Haas, T., Goettsch, B., Cuervo-Robayo, A. P., Urquiza-Haas, E., Orjuela-R, M. A., Gasman, F. A., Oliveros-Galindo, O., Burgeff, C., Rivera-Rodríguez, D. M., de Jesús Sánchez González, J., Alarcón-Guerrero, J., Aguilar-Meléndez, A., Cuevas, F. A., Alavez, V., Alejandre-Iturbide, G., Avendaño-Arrazate, C.-H., Pérez, C. A., ... Koleff, P. (2022). Incorporating evolutionary and threat processes into crop wild relatives conservation. *Nature Communications*, *13*, 6254. <https://doi.org/10.1038/s41467-022-33703-0>
- Van Oosterhout, C. (2020). Mutation load is the spectre of species conservation. *Nature Ecology & Evolution*, *4*, 1004–1006.
- Vernesi, C., Bruford, M. W., Bertorelle, G., Pecchioli, E., Rizzoli, A., & Hauffe, H. C. (2008). Where's the conservation in conservation genetics? *Conservation Biology*, *22*, 802–804.
- Waples, R. S., & Gaggiotti, O. (2006). What is a population? An empirical evaluation of some genetic methods for identifying the number of gene pools and their degree of connectivity. *Molecular Ecology*, *15*, 1419–1439.
- Wernberg, T., Coleman, M. A., Bennett, S., Thomsen, M. S., Tuya, F., & Kelaher, B. P. (2018). Genetic diversity and kelp forest vulnerability to climatic stress. *Scientific Reports*, *8*, 1851.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Hoban, S., da Silva, J. M., Mastretta-Yanes, A., Grueber, C. E., Heuertz, M., Hunter, M. E., Mergeay, J., Paz-Vinas, I., Fukaya, K., Ishihama, F., Jordan, R., Köppä, V., Latorre-Cárdenas, M. C., MacDonald, A. J., Rincon-Parra, V., Sjögren-Gulve, P., Tani, N., Thurfjell, H., & Laikre, L. (2023). Monitoring status and trends in genetic diversity for the Convention on Biological Diversity: an ongoing assessment of genetic indicators in nine countries. *Conservation Letters*, e12953. <https://doi.org/10.1111/conl.12953>