



HAL
open science

Establishment and evolution of ruminotypes of lactating Lacaune ewes

Tiphaine Blanchard, Christel Marie-Etancelin, Yves Farizon, Charlotte Allain,
Annabelle Meynadier

► **To cite this version:**

Tiphaine Blanchard, Christel Marie-Etancelin, Yves Farizon, Charlotte Allain, Annabelle Meynadier.
Establishment and evolution of ruminotypes of lactating Lacaune ewes. EAAP2023, Aug 2023, Lyon,
France. hal-04116731

HAL Id: hal-04116731

<https://hal.inrae.fr/hal-04116731>

Submitted on 5 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Establishment and evolution of ruminotypes of lactating Lacaune ewes

Tiphaine Blanchard¹, Christel Marie-Etancelin¹, Yves Farizon¹, Charlotte Allain², Annabelle Meynadier¹

¹ GenPhySE, Université de Toulouse, INRAE, ENVT, 31326, Castanet Tolosan, France

² INRAE, Experimental Unit of La Fage, F-12250 Saint-Jean et Saint-Paul, France

Ruminal microbiota of sheep is being studied to evaluate its impact on animals' energy balances in early lactation stage. The establishment of ruminotypes could be a useful tool to summarize the large dataset of ruminal microbiota abundances. The feasibility of directional selection of gut microbiota enterotypes has been demonstrated in pigs, proving the existence of genetic determinism of the host on enterotypes. Sixty-five Lacaune dairy sheep were studied and their ruminal contents and blood were sampled at 2 weeks (D1), 3 weeks (D2), 4 weeks (D3) and 6 weeks (D4) after lambing. Ewes were weighed on each date and were fed the same diet with hay, grass silage and concentrates. During the first month, lambs remained with the ewes: milk recording (milk quantity, fat and protein contents, somatic cell count) were performed at D4. Ruminal microbiota was analysed by sequencing the 16s rRNA gene for bacteria and sequences were computed with the FROGS pipeline to obtain relative abundances of ASVs which were then labelled according to genus level. Clustering of microbiota genera abundances using Jensen-Shannon divergence and partition around medoids clustering algorithm has led to 2 groups. Spls-da of these two clusters revealed a first ruminotype characterised by *Mogibacterium*, *Lachnospiraceae NK3A20 group* and *Acetivomaculum* genera (RuM) and a second one by *Prevotella*, *Prevotella_9* and *Lachnospiraceae NK4A136 group* genera (RuP). Over time, a significant shift towards RuP is noted (D1: 22%, D2: 34%, D3: 58%) which becomes a minority again after weaning (D4: 14%). At D1, RuM is associated with higher non esterified fatty acids ($p=0,04$), beta-hydroxybutyrate ($p<0,01$), isobutyrate ($p=0,01$) and lower total volatile fatty acids ($p<0,01$) compared with RuP. At D2, RuM is associated with lower weekly weight loss ($p=0,03$) compared to RuP. At D3, ruminotypes were not associated with zootechnical traits. At D4, RuM was associated with lower milk protein and fat contents ($p<0,02$). Characterization of the specific metabolic pathways of these 2 clusters is underway via a functional inference approach, in order to make links with the zootechnical traits.