



HAL
open science

Improving feed efficiency in meat sheep increases CH₄ emissions measured indoor or at pasture

Flavie Tortereau, Jean-Louis Weisbecker, Camille Coffre-Thomain, Yannick Legoff, Dominique Francois, Quentin Le Graverand, Christel Marie-Etancelin

► To cite this version:

Flavie Tortereau, Jean-Louis Weisbecker, Camille Coffre-Thomain, Yannick Legoff, Dominique Francois, et al.. Improving feed efficiency in meat sheep increases CH₄ emissions measured indoor or at pasture. EAAP2023, Aug 2023, Lyon, France. hal-04116732

HAL Id: hal-04116732

<https://hal.inrae.fr/hal-04116732v1>

Submitted on 5 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Improving feed efficiency in meat sheep increases CH₄ emissions measured indoor or at pasture.

Flavie Tortereau¹, Jean-Louis Weisbecker¹, Camille Coffre-Thomain², Yannick Legoff², Dominique François¹, Quentin Le Graverand¹, Christel Marie-Etancelin¹

¹ GenPhySE, Université de Toulouse, INRAE, ENVT, 31326, Castanet Tolosan, France ; ² INRAE Experimental unit P3R, 18390 Osmoy, France

Ruminants are often criticized when it comes to environmental impacts, due to CH₄ production and to the part of concentrates in the diets. A major objective in breeding ruminants is to limit both feed-food competition and greenhouse gas (GHG) emissions. We divergently selected, over 4 generations, Romane meat sheep on their Residual Feed Intake (RFI) from 3 to 5 months old under a 100% concentrate diet. We present here two results of GHG measurements (performed with sheep GreenFeed, C-Lock) on males and females belonging to these divergent lines (animals were from the RFI- (efficient) or RFI+ (inefficient) line). A total of 124 males belonging to the 3rd and 4th generations of selection, bred indoor, had both GHG and forage intake measurements. On average, males weighed 64.6 kg and emitted 1345 g/d of CO₂ and 39.26 g/d of CH₄. Positive correlations were estimated between CO₂ and CH₄ (0.59) and between gases and forage intake (0.24 for CH₄ and 0.54 for CO₂) and body weight (0.38 for CH₄ and 0.78 for CO₂). No significant differences were observed between RFI- and RFI+ males on body weight. RFI- males ate less forage than RFI+ ones (1.13 kg/d for RFI- and 1.22 kg/d for RFI+) but they emitted significantly more CH₄ (38.86 g/d for RFI- and 37.50g/d for RFI+), and less CO₂ (1326g/d for RFI- and 1335 g/d for RFI+). We confirmed this result with 85 ewes from the 4th generation of selection, first fed indoor with dry forage and then fed at pasture: RFI- ewes emitted significantly more CH₄ than RFI+ ewes, whatever the diet. GHG emissions were positively correlated between both diets (0.48 for CH₄ and 0.56 for CO₂). Animal effects for GHG emissions, obtained from a repeatability model were significantly correlated under the two diets (from 0.66 for CH₄ to 0.78 for CO₂ intensity). Both studies highlight unfavourable relationship between feed efficiency and gas emissions: the more efficient animals ate less, but emitted more CH₄ than the less efficient ones. Moreover, phenotyping GHG indoor under a forage-based diet is of high interest when phenotyping GHG at pasture is not feasible.