



**HAL**  
open science

## Grapevine yield gap: identification of environmental limitations by soil and climate zoning in the region of Languedoc-Roussillon (South of France)

Hugo Fernandez-Mena, Nicolas Guilpart, Philippe Lagacherie, Renan Le Roux, Mayeul Plaige, Maxime Dumont, Marine Gautier, Nina Graveline, Jean-Marc Touzard, Hervé Hannin, et al.

### ► To cite this version:

Hugo Fernandez-Mena, Nicolas Guilpart, Philippe Lagacherie, Renan Le Roux, Mayeul Plaige, et al.. Grapevine yield gap: identification of environmental limitations by soil and climate zoning in the region of Languedoc-Roussillon (South of France). *OENO One*, 2023, 57 (2), pp.360-378. 10.20870/oeno-one.2023.57.2.7246 . hal-04122556

**HAL Id: hal-04122556**

**<https://hal.inrae.fr/hal-04122556>**

Submitted on 8 Jun 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



ORIGINAL RESEARCH ARTICLE

# Grapevine yield gap: identification of environmental limitations by soil and climate zoning in the region of Languedoc-Roussillon (South of France)

Hugo Fernández-Mena<sup>1,2</sup>, Nicolas Guilpart<sup>3</sup>, Philippe Lagacherie<sup>4</sup>, Renan Le Roux<sup>5</sup>, Mayeul Plaige<sup>1</sup>, Maxime Dumont<sup>1,6</sup>, Marine Gautier<sup>1</sup>, Nina Graveline<sup>7</sup>, Jean-Marc Touzard<sup>7</sup>, Hervé Hannin<sup>8</sup> and Christian Gary<sup>1</sup>

<sup>1</sup> UMR ABSys – INRAE, CIRAD, Montpellier SupAgro, U. Montpellier – Montpellier, France

<sup>2</sup> UMR EMMAH – INRAE, U. Avignon – Avignon, France

<sup>3</sup> UMR Agronomie – AgroParisTech, INRAE, Université Paris-Saclay – Thiverval-Grignon, France

<sup>4</sup> UMR LISAH – INRAE, CIRAD, IRD – Montpellier, France

<sup>5</sup> U. AgroClim – INRAE – Avignon, France

<sup>6</sup> UMR ITAP – INRAE, Montpellier SupAgro, U. Montpellier – Montpellier, France

<sup>7</sup> UMR Innovation – INRAE, CIRAD, Montpellier SupAgro, U. Montpellier – Montpellier, France

<sup>8</sup> UMR MOISA – INRAE, CIRAD, Montpellier SupAgro, U. Montpellier – Montpellier, France



\*correspondence:

[hugo.fernandez-mena@supagro.fr](mailto:hugo.fernandez-mena@supagro.fr)

Associate editor:

José Miguel Martínez Zapater



Received:

3 Novembre 2022

Accepted:

25 March 2023

Published:

7 June 2023



This article is published under the **Creative Commons licence (CC BY 4.0)**.

*Use of all or part of the content of this article must mention the authors, the year of publication, the title, the name of the journal, the volume, the pages and the DOI in compliance with the information given above.*

## ABSTRACT

Often unable to fulfill theoretical production potentials and to obtain the maximum yields set by wine quality labels, many vineyards and cellars need to solve the issue of so-called grapevine yield gaps in order to assure their durability. These yield gaps particularly occur in Mediterranean wine regions, where extreme events have intensified because of climate change. Yield gaps at the regional level have been widely studied in arable crops using big datasets, but much less so in perennial crops, such as grapevine. Understanding the environmental factors involved in yield gaps, such as those linked to climate and soil, is the first step in grapevine yield gap analysis. At a regional scale, there are numerous studies on ‘terroir’ linked to wine typicity and quality; however, none have classified spatial zones based on environmental factors identified as being involved in grapevine yield. In the present study, we aggregated into one big dataset information obtained from producers at the municipality level in the wine region Languedoc-Roussillon (South of France) between 2010 and 2018. We used a backward stepwise model selection process using linear mixed-effect models to discriminate and select the statistically significant indicators capable of estimating grapevine yield at the municipality level. We then determined spatial zones by using the selected indicators to create clusters of municipalities with similar soil and climate characteristics. Finally, we analysed the indicators of each zone related to the grapevine yield gap, as well as the variations among the grapevine varieties in the zones. Our selection process evidenced 6 factors that could explain annual grapevine yield annually ( $R^2 = 0.112$ ) and average yield for the whole period ( $R^2 = 0.546$ ): Soil Available Water Capacity (SAWC), soil pH, Huglin Index, the Climate Dryness Index, the number of Very Hot Days and Days of Frost. The clustering results show seven different zones with two marked yield gap levels, although all the zones had municipalities with no or high yield gaps. On each zone, grapevine yield was found to be driven by a combination of climate and soil factors, rather than by a single environmental factor. The white wine varieties showed larger yield gaps than the red and rosé wine varieties. Environmental factors at this scale largely explained yield variability across the municipalities, but they were not performant in terms of annual yield prediction. Further research is required on the interactions between environmental variables and plant material and farming practices, as well as on vineyard strategies, which also play an important role in grapevine yield gaps at vineyard and regional scale.

**KEYWORDS:** viticulture, classification, mapping, cartography, big-data, terroir, climate change

## INTRODUCTION

### 1. The importance of grapevine yield gaps

Compared to the production of other types of crop, grapevine yield has been historically overlooked, assuming a trade-off between grape yield and wine quality (Bisson *et al.*, 2002; Jackson and Lombard, 1993; Poni *et al.*, 2018). Indeed, in viticulture, the issue of yield is treated in a particular way, since many production regions set a limit on grapevine yields within the framework of geographical indications (Stranieri and Tedeschi, 2019). However, in many vineyards, producers do not manage to reach the maximum grapevine yield authorised within the framework of the quality label. In France, Schauburger *et al.* (2018) showed that historical grapevine yields have stagnated since the 1980s. Some authors have even diagnosed a “vineyard decline” that could relate to environmental and management drivers (Riou *et al.*, 2016). Furthermore, in winegrowing regions with typically warm temperatures and low rainfall, such as the Mediterranean, climate change phenomena (i.e., increasing temperatures, droughts and extreme events) are expected to have a negative impact on grape yield (Bernardo *et al.*, 2018; Droulia and Charalampopoulos, 2021; Hannin *et al.*, 2021; Ollat *et al.*, 2017; Touzard *et al.*, 2017). Therefore, it seems necessary to increase knowledge about the factors involved in grapevine yield and explore potential adaptation measures to maintain or increase them. In the case of the *Pays d’Oc* Protected Geographical Indication (PGI), the label is particularly concerned about the long-term stability of its production levels, because (i) the yield is obviously a main driver of individual wine estate performance and its long-term sustainability (Tintinger, 2020), and (ii) as a product, the amount of *Pays d’Oc* on the market should be constant to guarantee satisfactory prices. In light of this, the label supports the increasing effort to explore avenues for stabilising yields at satisfactory individual and collective levels.


Yield gaps can be defined as the differences between the targeted yield and the obtained yield, and they result from a combination of environmental, genetic and management factors (Cooper *et al.*, 2021; Edreira *et al.*, 2017; Van Ittersum *et al.*, 2013), of which we focused on the environmental components in the present study. Environmental factors contribute to ‘resource yield gaps’, (e.g., the need for nutrients and water), but they also interact with management and genetic factors to create ‘efficiency yield gaps’ in the use of those resources (Silva *et al.*, 2017). Edreira *et al.* (2018) created the concept of ‘technology extrapolation domains’ to study zones with similar environmental resources for crop production. Andrade *et al.* (2019) added socio-economic variables to this framework to study the application of agricultural technologies in different countries. At the regional scale, defining zones with similar environmental variables is a first step towards understanding the local factors involved in yield gaps in large datasets with spatial variability (Beza *et al.*, 2017; Liu *et al.*, 2017; Van Wart *et al.*, 2013).

While numerous studies have been carried out on yield gaps in arable crops (Anderson *et al.*, 2016; Guilpart *et al.*, 2017; Pradhan *et al.*, 2015), very little is known about grapevine yield gaps, with only some very recent studies carried out in the Barossa and Eden valleys by Bonada *et al.* (2022), in which water-limited grapevine yield gaps were analysed. In contrast to other crops, most European wine production must comply with maximum yield requirements set by the geographical quality labels; this is the target yield for many growers (Stranieri and Tedeschi, 2019). In some cases, grapevine yields are intentionally limited to improve wine quality (Poni *et al.*, 2018), by applying specific farming practices, such as leaf removal, reduced fertilisation or deficit irrigation - or to limit other production factors, such as workforce or capital (e.g., if they do not optimise the maintenance of their vine stock capital by renewing the vineyard); however, in accordance with the quality label experts, we assumed these factors to be less significant for the study region and wine quality label studied. We therefore defined grapevine yield gaps as the difference between the yield maximum limit established by the corresponding quality label and the actual yield obtained.

### 2. Environmental factors involved in grapevine yield

Grapevine yield is known to develop during the course of two years and is linked to a combination of environmental and management factors associated with several key phenological stages (Guilpart *et al.*, 2014). As has been shown by studies at the plant scale, both soil and climate factors contribute to grapevine yield (Gerós *et al.*, 2015). Soil and climate factors are involved in several key grapevine stages, particularly those determining the availability of water resources (Gaudin *et al.*, 2014; Pellegrino *et al.*, 2005; Simonneau *et al.*, 2017). Furthermore, soil and climate indicators can be used not only at vineyard scale, but also at regional scale to determine and map key environmental characteristics involved in grapevine yield (Carbonneau *et al.*, 2015).

Different soil indicators have been used in vineyards to spatially define zones, such as erosion fragility (Chevigny *et al.*, 2014; Rodrigo-Comino and Cerdà, 2018), degree of soil compaction (Lagacherie *et al.*, 2006), carbon content (Bonfatti *et al.*, 2016) and mineral nutrition (Arnó *et al.*, 2012). Moreover, climate indicators have often been used to estimate grapevine phenology (García de Cortazar Aauri *et al.*, 2017; Gavrilescu *et al.*, 2018), identify suitable grapevine growing areas (Anderson *et al.*, 2012; Fraga *et al.*, 2014a; Moral *et al.*, 2016), determine cultivar growing conditions (Parker *et al.*, 2013; Santos *et al.*, 2019) and assess or predict the climatic events impacting grapevine yields, such as heatwaves, hail or frost (Fraga *et al.*, 2020; Petoumenou *et al.*, 2019; Sgubin *et al.*, 2018). Numerous climatic models predict (Quénol *et al.*, 2017) that climate change will threaten grapevine yields by climatic extreme events or worsened water scarcity in specific regions. For instance, simulations performed by Fraga *et al.* (2016) have indicated that climate change is projected to impact grapevine yield in southern Europe as a result of dryness.



**FIGURE 1.** (A) The former Languedoc-Roussillon French administrative region until 2015, now part of Occitanie region (Wikialine., 2009). (B) Land-use dedicated to grapevine in Languedoc-Roussillon, in 2018. Adapted from IGN (2020).

Santos *et al.* (2011) explored climatic evolution in several zones of northern Portugal, helping to identify potential local adaptation strategies. While the prediction of climate evolution in vineyards is useful when developing adaptation strategies to mitigate climate change impacts (Mosedale *et al.*, 2016; Naulleau *et al.*, 2021; Santillán *et al.*, 2019), precise local adaptation can only be applied through a fine spatial description of the environmental factors involved (Naulleau *et al.*, 2022).

### 3. Soil and climate zoning in viticulture

In viticulture, zoning is traditionally based on the concept of ‘terroir’, which refers to the relationship between the climate, soil and vine compartments, and farming practices, with a final objective of ensuring wine quality and typicity (van Leeuwen and Bois, 2018). Recently, many authors have suggested that terroir studies should become more unbiased by including precision agriculture techniques for measuring environmental data (Bonfante and Brillante, 2022; Bramley and Hamilton, 2007; Brillante *et al.*, 2020; Vaudour *et al.*, 2015). Numerous terroir classifications have thus been proposed to try and better explain terroir in terms of the environmental variables of a given wine growing region (usually trying to find a link with wine composition), these classifications have sometimes used grapevine yield data to explain terroir (Bramley *et al.*, 2011; Bramley *et al.*, 2020; Peng *et al.*, 2021). However, little attention has been paid to spatial classifications that consider the environmental factors involved in grapevine yield.

At regional and sub-regional scales, climatic indicators have been used to classify viticultural production. Tonietto and Carbonneau (2004) proposed the MCC (Multicriteria Climatic Classification) system based on three climate indicators that also proved to be correlated with the typicity of wine sensory characteristics (Tonietto *et al.*, 2014). Bois *et al.* (2018) proposed a temperature-based zoning of the Bordeaux wine region to predict maturity dates. Resco *et al.* (2016) classified the Spanish viticultural regions helping to exploring adaptation to future climate, and Blanco-Ward *et al.* (2019) used bioclimatic indicators to study potential climate change effects on the quality of Portuguese Douro wines.

Regarding soil indicators, van Leeuwen *et al.* (2010) proposed a soil-based zoning methodology for viticultural terroirs. Although a high diversity of soil indicators in vineyards exists (vanLeeuwen *et al.*, 2018), some authors prefer to consider geological soil factors (Ferretti, 2019). Ultimately, since the availability of soil sample data is limited, resulting in imprecise upscaled cartography, soil data is often the bottleneck in zoning, (White, 2020). Therefore, only a few studies have combined both soil and climatic indicators for the zoning of wine regions; for example, the classification of viticultural terroirs in the Iberian Peninsula by Fraga *et al.* (2014b); the classification of the north-west zones in Iberian Peninsula by Cardoso *et al.* (2019); the zoning of terroirs in Denmark by Peng *et al.* (2021); and the classification of the Australian Barossa sub-zones by Bramley and Ouzman (2022). However, no methods exist for the spatial classification of soil and climate indicators directly related to grapevine yield.

### 4. Purpose of this study

In this study, we used climate and soil indicators linked to grapevine yield gap to determine spatial zones within the winegrowing region of ‘Languedoc-Roussillon’ (South of France, south-eastern part of the French Occitanie region). We collected 9 years’ worth of data from grapevine producers under the *Pays d’Oc* PGI quality label - who are hence all subject to the same maximum yield requirements established by the label - and aggregated them at the municipality level. We calculated, at municipality level, the soil and climate indicators that influence grapevine yield according to the scientific literature. We then kept only those indicators proving to have a significant effect on the grapevine yield at municipality level. Then, we clustered the zones presenting similar indicators, thus facilitating the identification of environmental resources. We hypothesised that (i) the combination of climate and soil can result in different yield levels (e.g., low-yielding and high-yielding environments), (ii) the same yield level can be obtained with different combinations of climate and soils, and (iii) some grape varieties are preferentially cultivated in specific combinations of climate and soils with higher associated yields.

## MATERIALS AND METHODS

### 1. Data sources

#### 1.1. Yield data

Grapevine yield data were obtained from harvest customs declaration data provided by producers under the *Pays d'Oc* PGI in the former Languedoc-Roussillon region (Figure 1A). This label is the largest wine label in France in terms of cultivated area, wine produced and number of winegrowers, with over 1100 wine cellars. Between 70,000 and 120,000 ha are declared under this label every year, which represents over 50 % of the grapevine cultivated area in Languedoc-Roussillon (Figure 1B). The label sets a maximum red and white wine production limit at 90 hl·ha<sup>-1</sup>·year<sup>-1</sup>. Although the maximum yield for rosé wine was increased from 90 to 100 hl·ha<sup>-1</sup>·year<sup>-1</sup> in 2015, we considered 90 hl·ha<sup>-1</sup>·year<sup>-1</sup> as the generic yield objective and threshold for all years and wine colours. The dataset contains a total of 96,667 yield data for a period of 9 years (from 2010 to 2018), 58 grapevine varieties and 606 municipalities. The yield data (hl·ha<sup>-1</sup>·year<sup>-1</sup>) was aggregated on a yearly basis at municipality level for all grapevine varieties and vineyards, resulting in 4455 annual municipality yield data. These data are available in the following repository: <https://doi.org/10.57745/THCVRE> (Fernández-Mena, 2022).

#### 1.2 Weather data

We used weather data from the SAFRAN reanalysis (Vidal *et al.*, 2010). The SAFRAN data cover the whole of the French territory at a spatial resolution of 8km by 8km and provide weather variables relevant to crop growth. These data have already been used to investigate the effects of climate on crop yield (e.g., Ben-Ari *et al.*, 2018). We used SAFRAN data provided by *MétéoFrance* and extracted from the INRAE SICLIMA platform at a daily time step between 1 January 2010 and 31 December 2018 for the following variables: daily average (*tmean*, °C), maximum (*tmax*, °C) and minimum temperature (*tmin*, °C), daily rainfall (mm), and daily reference evapotranspiration (*ETo*, mm).


#### 1.3 Soil data

Two soil indicators were considered, as they can have an important effect on grapevine yield: soil available water holding capacity (*SAWC*, mm) and soil pH (*pH*, no units). High *SAWC* can buffer against transient water deficits, which is particularly relevant in Mediterranean areas during dry periods in spring and summer (Coipel *et al.*, 2006; van Leeuwen *et al.*, 2018). Soil pH can affect grapevine yield negatively, particularly at very low pH levels (Himelrick, 1991; Tagliavini and Rombola, 2001).

We used the latest update of the Soil Available Water Capacity (*SAWC*) cartography in Languedoc-Roussillon (Styc and

**TABLE 1.** Variables analysed in this study.

	Variable name	Acronym	Units	Minimum	Maximum	Average
Grapevine	Municipality average grapevine yield	Yield	hl·ha <sup>-1</sup> ·year <sup>-1</sup>	6.00	100.00	67.03
Soil	Soil Available Water Capacity	SAWC	mm	55.00	133.26	88.79
	Soil pH	soil_pH	-	5.30	8.44	7.91
Climate	Climatic Dryness Index	CDI	mm	-346.3	146.2	-221.6
	Huglin Index	HI	°C	1406	2709	2270
	Days of Frost	DoF	days	0	65	16.76
	Frequency of Late Frost	FoLF	days	0	2	0.01
	Severity of Late Frost	SoLF	°C	0	0.4	0.0002
	Hot Days	HD	days	0	62	31.83
	Very Hot Days	VHD	days	0	12	1.8
	Severity of Heat Stress	SoHS	°C	0	29.5	2.16



**FIGURE 2.** Municipality average grapevine yield (weighted by their area for all grapevine varieties) in Languedoc-Roussillon of Pays d’Oc PGI labelled wines between 2010 and 2018 ( $n = 606$  municipalities). Top left: histogram of the displayed data for all municipalities and 9 years between 2010 and 2018 ( $n = 4455$ ) averaged from individual yield declarations ( $n = 96677$ ).

Lagacherie, 2019), which provides SAWC values at three maximum rooting depths (60, 100 and 200 cm). Validation by comparing with local SAWC observations revealed a substantial level of uncertainty in this map (Styc and Lagacherie, 2021). However, the map displays pedologically sound spatial patterns of predicted SAWC that justify its use for large scale studies, such as the present study (Styc and Lagacherie, 2021). We used soil pH data from the Languedoc-Roussillon *GlobalSoilMap* (Vaysse and Lagacherie, 2015) at a spatial resolution of 90m, following the *GlobalSoilMap* specifications (Arrouays *et al.*, 2019). The map had been created using legacy-measured soil profiles associated with a set of soil covariates using random forest and by kriging. The data covers four soil layers at depths of 0-5 cm, 5-15 cm, 15-30 cm and 30-60 cm and shows high prediction performance ( $R^2 = 0.71 - 0.73$ ). Both datasets (SAWC and pH) are publicly available for download at <https://www.openig.org/>.

## 2. Data preparation

### 2.1 Data aggregation at the municipality level

Yield, climate and soil data were aggregated at the municipality level. Average grapevine yield ( $\text{hl}\cdot\text{ha}^{-1}\cdot\text{year}^{-1}$ ) was calculated as the area-weighted average of yields over all grapevine varieties in each municipality. SAWC map raster was intersected with the municipalities of the region studied and an area weighted average of SAWC was calculated for each municipality. We calculated the municipality soil pH as the average pH of several soil layers at depths of 0-5, 5-15, 15-30 and 30-60 cm. The weather data were aggregated at municipality level using the nearest neighbour method.

### 2.2 Calculation of climatic indicators

Drawing from current knowledge of grapevine physiology, a total of eight climatic indicators were initially selected for potentially being able to explain annual and spatial variations in grapevine yield. These can be grouped into four types. The first type of climate indicator provides information about temperature suitability for grapevine development (Tonietto *et al.*, 2014; Tonietto and Carbonneau, 2004), and comprises a single indicator: the Huglin Index (*HI*, degree Celsius), proposed by Huglin (1978). This index is calculated as the sum of temperatures over a threshold of 10 °C, using both the mean and maximum daily air temperatures, during the grapevine vegetative period; i.e., from the beginning of April to the end of September in the northern hemisphere. The sum is multiplied by a length-of-day coefficient ( $d$ ). We used a  $d$  equal to 1.03 for the latitude of Languedoc-Roussillon region (42°01' – 44°00').

The second type describes water availability and includes a single indicator: the Climatic Dryness Index (*CDI*, mm), calculated every year as the difference between cumulated rainfall and ETo from 1 April to 30 August. The CDI is similar to the Standardised Precipitation-Evaporation Index of Vicente-Serrano *et al.* (2010) that was used to categorise viticultural areas by Santillán *et al.* (2019). However, we limited the calculation of the indicator to up to 30 August to exclude any potential extreme rainfall events in the study region close to harvest during the month of September.

The third type of climate indicator describes heat stress and comprises three indicators: (i) Hot Days (*HD*, days),

calculated as the number of days per year with maximum temperatures over 30 °C. Temperatures over 30 °C hamper carbon assimilation and limit yield by reducing the number of berries per vine (Greer and Weedon, 2014; Petrie and Clingeleffer, 2005; Torregrosa *et al.*, 2017). (ii) Very Hot Days (VHD, *days*), calculated as the number of days per year with maximum temperatures over 35 °C. Very high temperatures can have negative effects on grapevine photosynthesis and fruit set above 35 °C (Luo *et al.*, 2011; Pagay and Collins, 2017) and can also cause leaf scorch and fruit wilt above 40 °C (Galet, 2000; Liu *et al.*, 2019; Pagay and Collins, 2017; Venios *et al.*, 2020). (iii) Severity of Heat Stress (SHS, *degree Celsius*), calculated as the accumulation of daily mean temperature over 28 °C from budburst (considered as 1 April) to harvest (considered as 30 September). Around 28 °C is considered as the optimum daily mean temperature for grapevine photosynthetic activity in a Mediterranean context (Schultz, 2000; Xiao *et al.*, 2017). Heat stress will increase proportionally to the degree days accumulated over this optimum, reducing grapevine photosynthetic activity and thus yield.

The fourth type of climate indicator describes cold stress and comprises three indicators: (i) Days of Frost (DF, *days*), calculated as the total number of days per year with minimum temperature below 0 °C. Low winter temperatures can cause seasonal changes to fruit structure and a decrease in yield; extreme cold events (i.e., < -10 °C) can create injuries in vines (Buztepe *et al.*, 2017; Kaya and Köse, 2017). This indicator is also correlated with the probability of larger-scale local episodes of late frosts not being detected by climate stations. (ii) Frequency of Late Frost (FLF, *days*), calculated as the number of days with minimal temperature below 0 °C from 1 April to 30 September. Late frosts are a significant hazard for grapevines as they cause considerable damage to plant tissue (Molitor *et al.*, 2014; Trought *et al.*, 1999). (iii) Severity of Late Frost (SLF, *degree Celsius*), calculated as the accumulation of daily minimum temperatures below 2 °C from 1 April to 30 September. The most severe frost impact increases with extremely low temperatures below 0 °C (Poling, 2008), but some temperatures between 0 and 3 °C have proven to significantly impact grapevine growth

(Hendrickson *et al.*, 2004), and local frost episodes can also occur when downscaling the average temperature to specific vineyards.


### 2.3 Selection of climate and soil indicators that best explain variations in grapevine yield

After identifying and calculating an initial set of climate and soil indicators relevant to grapevine yield (Table 1), we used linear modelling to select indicators that best explained the variations in grapevine yield at the municipality level. Although some indicators, such as the Huglin Index and the number of hot days, were correlated (SI: Figure 2), we did not eliminate any indicators based on correlations. Instead, a backward stepwise model selection process was used to select indicators the most related to grapevine yield, following Zuur *et al.* (2009), Gareth *et al.* (2013) and Cayuela (2018).


Our yield prediction model tested all the soil and climate variables listed in Table 1 as predictors for each municipality yield and year ( $n = 4455$ ), as well as for the average grapevine yield of the municipality for the whole period studied ( $n = 606$ ). These models were fitted using linear mixed-effect models (LMM). LMM are commonly employed in analyses of grouped data where observations cannot reasonably be assumed to be independent of one another (Pinheiro and Bates, 2006). In our case, climate and soil indicators were considered as fixed effects for the prediction of average grapevine yield at the municipality level, and the municipality was assumed to have a random effect, since there is a spatial dependence between among interannual yield data for the same municipality (Bonansea *et al.*, 2015). In the backward stepwise selection process, we started with the most complex model and dropped one variable at each step depending on the p-values, until the remaining variables had significant p-values (< 0.05). Then, we selected the most parsimonious model using both Akaike information criterion (AIC) and Bayesian information criterion (BIC). This technique of model selection is also used in earth science studies, such as that of Cremona *et al.* (2018), to discriminate factors involved in ecological processes. Random forest was used to discriminate the importance of variables by ranking

**TABLE 2.** Fixed effects with Estimates, Confidence Interval at 95 % (CI) and p-values of the most parsimonious mixed-model results to estimate annual grapevine yield at the municipality level ( $n = 4455$ ). Predictors other than intercept are ordered by random forest partial dependence plot of variable importance (Figure S2). P-values are flagged: \* for < 0.05 ; \*\* for < 0.01 and \*\*\* for < 0.001.


Predictors	Estimates	CI	p-values
Intercept	13.57	-2.61 – 29.75	
Soil Available Water Capacity (SAWC)	0.27	0.23 – 0.32	***
Huglin Index (HI)	0.02	0.01 – 0.02	***
Soil pH	0.169	-2.48 – 3.86	**
Climatic Dryness Index (CDI)	0.03	0.02 – 0.05	***
Days of Frost (DF)	-0.15	-0.33 – -0.18	***
Very Hot Days (VHD)	-0.55	-0.77 – -0.33	**



**FIGURE 3.** Average Soil Available Water Capacity (SAWC) of each municipality in Languedoc-Roussillon. Adapted from Styc and Lagacherie (2021).



**FIGURE 4.** Average Climatic Dryness Index (CDI) of each municipality in Languedoc-Roussillon between 2010 and 2018.



**FIGURE 5.** Average Huglin Index (HI) of each municipality in Languedoc-Roussillon between 2010 and 2018.

the predictors of the selected model as a function of their contribution to reducing prediction error (Strobl *et al.*, 2008).

Statistical analyses were performed using the R software (R Core Team., 2014) with ‘nlme’ package (Pinheiro *et al.*, 2021) for LMM and ‘corrplot’ package for correlation matrix calculations (Wei and Simko, 2017). Datamining, plots and tables were done using the packages ‘dplyr’ (Wickham *et al.*, 2018), ‘sjPlot’ (Lüdecke, 2021), and ‘ggplot2’ (Wickham, 2016).

### 3. Spatial clustering and zone assessment

Based on the set of selected indicators as described in Section 2.2, we clustered the municipalities with similar soil and climate, helping us to create groups of municipalities that we refer to as zones. For the clustering, we used a combination of principal components analysis (PCA) and ascendant hierarchical classification (AHC). PCA and AHC on principal components were performed using the ‘FactomineR’ R package (Lê *et al.*, 2008). The number of clusters in the AHC was defined in collaboration with local viticultural experts (grapevine growers and wine label managers of Pays d’Oc PGI) to capture a representative range of the pedoclimatic regional variability. The Ward’s method suggested using three clusters, but this number of clusters was too low and did not include some important factors such as pH, CDI or VHD.

We assessed and described the characteristics of the defined zones in terms of the grapevine yield gap (as the difference between the average grapevine yield and the label production limit; i.e., 90 hl·ha<sup>-1</sup>·year<sup>-1</sup>). For every zone, we calculated the selected indicators in Section 2.2, the distribution of grapevine varieties, the number of municipalities and the total grapevine cultivated area. Tukey’s range test was then used to compare the average values between zones using

‘multcomp’ R package (Hothorn *et al.*, 2009). Anova test (R Core Team., 2014) was used to estimate the variability of the grapevine yield distribution explained by the zone classification grapevine yield, and to compare it to the unexplained variability inside each zone. Chi-Square test of goodness of fit using ‘summarytools’ R package (Comtois, 2021) was used to analyse the proportions of grapevine varieties in each zone (with a significance level alpha = 0.05), comparing their distribution to an expected equilibrated distribution among the zones.

## RESULTS

### 1. Descriptive statistics of grapevine yield data

In our database, the vineyard cultivated area and wine volume as declared by the Pays d’Oc quality label were, on average, 73718 ha·year<sup>-1</sup> and 5 950 302 hl·year<sup>-1</sup>. However, we aggregated individual yields weighted by their area at the municipality level, obtaining an average yield per municipality and year. Yield values for all the municipalities were as follows: a mean of 65.3 hl·ha<sup>-1</sup>·year<sup>-1</sup>, a median of 67.03 hl·ha<sup>-1</sup>·year<sup>-1</sup>, a minimum of 6 hl·ha<sup>-1</sup>·year<sup>-1</sup>, and a maximum of 100 hl·ha<sup>-1</sup>·year<sup>-1</sup> (the latter corresponding to a municipality with rosé wines - permitted maximum yield being 100 hl·ha<sup>-1</sup>·year<sup>-1</sup>). If the yield gap was filled, the extra production expected could be 684 318 hl·year<sup>-1</sup> (i.e., over 11.5 % of the current wine volume) with regards the label’s maximum yield requirements for red and white wines (90 hl·ha<sup>-1</sup>·year<sup>-1</sup>).

Spatially, the estimation of average grapevine yield at the municipality level between 2010 and 2018 revealed localised yield gaps in numerous municipalities (Figure 2).

Temporally, no declining trend was observed within this time frame, although some years had lower yields (Figure S1), in particular 2010 and 2017, which were linked to severe drought conditions.

The most planted grapevine varieties in the region were, in order of cultivated area, Merlot, Cabernet-Sauvignon, Syrah, Chardonnay, Grenache noir, Sauvignon blanc and Cinsault. In total, 10 varieties were grown over 70 % of the cultivated area (Table ST1). The yield of white wine varieties including Chardonnay, Grenache blanc, Viognier and Muscat petit grain were generally lower than 70 hl·ha<sup>-1</sup>·year<sup>-1</sup> and lower than that of red and rosé wine varieties. Yield differences between red wine varieties were low.

## 2. Identification of six relevant climate and soil indicators


Of the models tested in Table ST2, we selected a mixed model that maximised AIC and BIC performance, for which 6 (Table 2) of the 10 calculated indicators proved to have a significant effect on the annual grapevine yield of the municipalities (n = 4455). This method obtained a low marginal  $R^2$  (0.112), thus showing low potential for annual yield prediction. Yet, the same predictors proved to be more relevant for the prediction of average grapevine yield for the whole period (n = 606), with a marginal  $R^2$  of 0.546 and a conditional  $R^2$  of 0.627. The variables that were found to have a significant effect on grapevine yield at the municipality level were, in order of increasing significance: Soil Available Water Capacity (Figure 3), Climatic Dryness Index (Figure 4), Huglin Index (Figure 5), Days of Frost (Figure S3), soil

pH (Figure S4) and the Very Hot Days (Figure S5). Despite their theoretical impact on grape yield, four indicators were excluded from our model: SLF, FLF, HD and SHS (Figures S6, S7, S8 and S9). A random forest partial dependence plot of variable importance (Figure S2) ranged the variables according to their predicting capacity, as shown in Table 2. The Pearson correlation matrix showed a high positive correlation between the indicators related to extreme heat events (HD, VHD and SHS), also with temperature accumulation in HI, whereas CDI was highly and positively correlated with DF and highly and negatively correlated with HI (Figure S2). These correlations helped to discriminate HD and VHD for improving model performance and verify the low correlations of all the factors with the Municipality as an aggregated variable.


## 3. Seven agroecological zones of municipalities with similar climate and soil conditions

The PCA and HCH statistical analyses helped to define seven clusters of municipalities using the selected soil and climate variables as listed in Section 3.2 (Figure S10). Each of those clusters represents an agroecological zone with similar soil and climate characteristics (Figure 6). The characteristics of the zones are significantly different from each other in terms of at least one index that favours or constrains grapevine yield (Figure 7).

Zone 1 is the ‘*Humid zone of the hinterland*’ and has the coolest temperatures due to its distance from the Mediterranean coast. As a consequence of having the lowest



**FIGURE 6.** Soil and climate zones related to grapevine yield at the municipality level in Languedoc-Roussillon.



**FIGURE 7.** Selected soil and climate indicator distribution for each of the seven zones (clusters) in the Languedoc-Roussillon region. The box-plots show the distribution in quartiles with median lines. Circles represent the mean value and filled dots are outliers. Letters correspond to Tukey's range test for means comparison.

Climate Dryness Index (CDI) (around -150 mm) and number of Very Hot Days (VHD) (from 0 to 2), this region benefits from high grapevine yield. Its main constraint is its Huglin Index (HI), which is the lowest, with 300 to 400 degree-days less than other zones. The Soil Available Water Capacity (SAWC) is relatively high (from 70 to 100 mm) and Days of Frost (DF) are average (from 10 to 20).

Zone 2 is the ‘Zone with acid and shallow soils in the mountains’, which is the only one with an acid soil pH (ranging from 5 to 7.5), which constrains grapevine yield. This zone also has the lowest SAWC (from 50 to 80 mm) and a low HI. The rest of the variables are average. The municipalities of this zone are located at the highest elevations with municipalities in the southern (Pyrenees mountains) and northern (Caroux Mountains) areas of the region.

Zone 3 is the ‘Zone of piedmont with constraining SAWC’. It has low temperature-related variables (HI; DF and VHD) similar to those in Zone 2, but municipalities in this zone have alkaline soil pH (from 7 to 8.3). Water-related indicators are also not very favourable, although SAWC is significantly higher than in Zone 2. CDI is lower, being linked to higher temperatures and the HI. The municipalities of this zone are located at mid-elevation and in the piedmont areas of the region.

Zone 4 is the ‘Cold and dry zone surrounding Pic St Loup’. This zone is constrained by numerous Days of Frost (DF), but high temperatures in summer (high HI and VHD). This zone is also constrained by low water availability from rainfall (low CDI) and soils (low SAWC). The municipalities

of this zone are located in high areas surrounding the peak Saint-Loup (north of Montpellier).

Zone 5 is the ‘Zone of average inland soils’. It comprises relatively average soils, although SAWC is very variable. The region is constrained by a high Climatic Dryness Index (CDI) and the highest Huglin Index. The municipalities of this zone are mainly located on the inland plains in the central and eastern parts of the region.


Zone 6 is the ‘Zone with deep soils in mild coasts’. It comprises the best soils (highest SAWC), compensating for having the highest water deficit (highest CDI) in the region. Extreme temperatures are rare in this zone due to the proximity of the sea.

Zone 7 is the ‘Highest number of very hot days but deep soils’. It is subject to the most extreme temperatures with the highest level of Very Hot Days (VHD) and many Days of Frost (DF). In contrast, water availability is favourable due to deep soils (high SAWC) and less dry climate (low CDI). The municipalities of this zone are located on several inland plains in the eastern part of the region.

#### 4. Assessment of the yield gap and varieties per agroecological zone

Depending on their yield gaps (i.e., the difference between label maximum yield (90 hl·ha<sup>-1</sup>·year<sup>-1</sup>) and obtained yield), the clustered zones can be divided into two main groups (Figure 8):

1. The group with the highest yield gaps, from 30 to 50 hl·ha<sup>-1</sup>·year<sup>-1</sup>; i.e., yields ranging from 50 to 60 hl·ha<sup>-1</sup>·year<sup>-1</sup>. This group corresponds to municipalities in



**FIGURE 8.** Distributions of average municipality grapevine yields (n = 606) in hl·ha<sup>-1</sup>·year<sup>-1</sup> in each of the seven clustered zones in Languedoc-Roussillon between 2010 and 2018. The boxplots represent the distribution in quartiles with median lines. Circles represent the mean and filled dots are outliers. Letters correspond to Tukey's range test for comparison of means. The dashed red line corresponds to the 90 hl·ha<sup>-1</sup>·year<sup>-1</sup> maximum label yield used for yield gap calculation. The percentage in brown corresponds to the coefficient of variation over time for each zone.

Zones 2, 3 and 4. Within this group, Zone 3 has a significantly higher yield.

2. The group with the lowest yield gaps, from 10 to 25 hl·ha<sup>-1</sup>·year<sup>-1</sup>; i.e., yields ranging from 65 to 80 hl·ha<sup>-1</sup>·year<sup>-1</sup>, which corresponds to the municipalities of Zones 5, 6 and 7.

Municipalities in Zone 1 show an intermediate yield gap between the two above-described groups.

The conditions for the application of the Anova test on grapevine yield depending on zone were validated by the Shapiro-Wilk normality test. The variability in grapevine yield explained by the clustered zones is significant according to the Anova test. The F-value obtained was 11.54, indicating more variability explained by the zones than not explained inside the zones.

We observed a high variation in yield levels depending on the zones as shown by the coefficient of variation in Figure 8. Although over time there was not a significant tendency towards lower yields in the zones, the zones with low yields (i.e., Zones 2, 3 and 4) drastically reduced their yields in occasional years (Figure S11). Zones 5 and 6 account for the highest cultivated areas (i.e., 20000 to 25000 ha) and Zones 4 and 2 for the lowest (i.e., 1000 to 2500 ha) (Figure S12). In addition, these zones account for the highest and the lowest number of municipalities respectively (Figure S12).

Concerning grapevine varieties, we found a total of 15 different grapevine varieties based on the distribution of the top 10 varieties in each of the seven pedo-climatic zones (Figure S13); the selected varieties represent over 81 % of the total diversity grown under the *Pays d'Oc* PGI label during the years studied. Of these varieties, 9 of them represented between 75 and 93 % of the planted area depending on the zone (Table ST3); these were, in order of increasing planted area: Merlot, Cabernet-Sauvignon, Syrah, Chardonnay, Grenache noir, Sauvignon blanc, Cinsault, Viognier and Pinot noir. We analysed the yield distribution of each of these nine wines per zone, resulting a similar overall distribution (Figure S14). The white wine varieties showed lower yields than the red wine varieties. As well as these popular varieties, only two original varieties were found in a few zones: Cabernet franc in Zone 1 and Muscat in Zone 2 and 3.

## DISCUSSION AND PERSPECTIVES

In the following sections, we first discuss the limitations of the data used and selection of indicators (Section 4.1). We then refer to the zoning results to give recommendations for management adaptations (Section 4.2), discuss other factors involved in grapevine yield gaps that would need to be studied to complete this work (Section 4.3), and, finally, give our conclusions (Section 4.4).

### 1. Limitations of climate and soil data and selected indicators

Soil data availability at the landscape scale is often the bottleneck when zoning terroir and in grapevine production

(White, 2020); in particular, the lack of an SAWC map and measurements constrains many classifications, such as that of the Italian Chianti, as suggested by Priori *et al.* (2019). As a result of the *GlobalSoilMap* initiative, new soil databases are available at the regional scale for Languedoc-Roussillon (Vaysse and Lagacherie, 2015). In this study, we showed that not only climate, but also soil variables (*SAWC* and *soil pH*) are relevant when studying regional grapevine yield gaps. However, *GlobalSoilMap* products are less useful for doing these analyses on more local scales. Although the *SAWC* map explained a small amount of the total SAWC variance (20 % according to Styc and Lagacherie (2021), the aggregation of SAWC at the municipality level reduced this variance, as observed by Vaysse and Lagacherie (2017).

The use of SAFRAN climate data raises a number of questions. On the one hand, the 8 km by 8 km resolution reduced out the local variability of the climate, which can sometimes be very important in magnitude over small distances and participate in a significant way to the heterogeneity of the vineyard environment (Le Roux *et al.*, 2017). On the other hand, SAFRAN is a reanalysis, which necessarily produces biases, especially in areas where relief is more pronounced (Quintana-Segui *et al.*, 2008). SAFRAN tends to minimise extreme temperatures (Ollat *et al.*, 2021), thus decreasing the number of *Very Hot Days* and *Days of Frost*. Because of the scale used, we could not consider certain local phenomena such as hail, that can drastically damage grapevine yield (González-Fernández *et al.*, 2020). The analysis of a longer climate data series (i.e., over 20 or 30 years) would help to improve the relevance of climatic indicators, but they would require grapevine yield data from the same period and length of time. Another aspect that could be considered is the integration of plant phenology into the calculation of the indicators. Indeed, the impacts of climate on the functioning of the plant differ depending on the phenological stage of the plant, and they therefore depend highly on the variety used (Morales-Castilla *et al.*, 2020). A possible approach to integrating these effects would be to calculate the indicators based not on calendar dates but on periods corresponding to key stages of the vine (budburst, flowering, ripening, etc.); this has already been applied to other crops (Caubel *et al.*, 2015).

Most of the proposed indicators proved to be significant determinants of grape yield. Despite the influence of high temperature being partially explained by the number of *Very Hot Days*, the number of *Hot Days* and *Severity of Heat Stress* were, surprisingly, not significant indicators in a region with numerous heat events (Garnier, 2010). In addition, the period studied did not account for the most severe heat wave in the South of France recorded in 2019, which severely impacted grapevine yield (Lopez-Fornieles *et al.*, 2022). The selection of the *Days of Frost* indicator may seem surprising from the point of view of extreme event impact, yet many *Days of Frost* in winter will delay budburst, which, for some cultivars, reduces the length of the vegetative cycle and therefore yield (Cameron *et al.*, 2022). The *Frequency of Late Frost* was not deemed significant, since frosts were very marginal during the

period studied (Figure S8). Although rare, some strong late frost events can occasionally occur in Languedoc-Roussillon; a noteworthy example is that of the historic early frost of April 2021 that caused much damage in all the southern viticultural regions of France and whose consequences are still being assessed (DDTM du GARD., 2022). Still under discussion are some projections made by authors of a reduction in frost events in Europe (Leolini *et al.*, 2018); meanwhile, a recent modelling study in France has predicted a higher risk of them occurring (Vautard *et al.*, 2022).

## 2. The role of 7 pedoclimatic zones to guide R&D in viticulture in Languedoc-Roussillon

The 7 pedoclimatic zones explained more than 50 % of the grapevine yield spatial variations at the scales studied, despite being based on few variables. In each zone, we identified some indicators that exceeded the regional average and which may have limited grapevine yield, so we proposed some specific management adaptations. For instance, Zone 1 has the lowest *HI*; therefore, growing grapevine varieties adapted to lower cumulated temperatures, such as Merlot, Cabernet franc and Cabernet-Sauvignon, could help to improve or maintain grapevine yield (Morales-Castilla *et al.*, 2020; Parker *et al.*, 2013; Parker *et al.*, 2020). Zone 2 has the lowest *SAWC* and *soil pH*; therefore, rootstock adaptation to lower pH and water stress could help to improve productivity in this area (Himelrick, 1991; Serra *et al.*, 2014; Tagliavini and Rombola, 2001; Vrsic *et al.*, 2016). Zones 3, 4, 5 and 6 are affected by water stress, either having a low *SAWC* (especially in Zone 4), a high Climatic Dryness Index (in particular in Zone 6, as well as in Zone 5) or both the latter (in Zone 3). The combination of water and thermic stress are known to increase stomata closure, which negatively affects grapevine photosynthesis (Carvalho and Amâncio, 2019). Adaptations to water scarcity can include rootstock and cultivar choice, irrigation, decreasing planting density, weed early control and cover crop management (Duchêne, 2016; Lovisollo *et al.*, 2016; Naulleau *et al.*, 2021). Zone 7 also has limitations linked to extreme temperatures, which are low during the winter (i.e., high number of *Days of Frost*, thus shortening the grapevine cycle) and very high during the summer (i.e., high number of *Very Hot Days*, severely impacting grapevine yield as shown by Lopez-Fornieles *et al.* (2022)). The introduction of shading nets and tree shades in this zone could help to prevent extreme temperatures and scorching of the vines (Grimaldi *et al.*, 2019; Oliva Oller *et al.*, 2022; Villalobos-Soublett *et al.*, 2021; Williams *et al.*, 2022).

Furthermore, the adaptations to be implemented will depend on the total cultivated area of a given zone. Zones 5 and 6 are the most cultivated, followed by 1, 7 and 3. Within each zone, we observed that the municipality variability of the yield gaps was higher than their indicator variability. Consequently, it would be worth investigating the causes of municipalities in a high yield zone having a low average yield gap, and, if relevant, improving farm management practices to fulfill the environmental potential of the zone.

## 3. Other non-environmental factors are involved in grapevine yield gaps

Indeed, climate and soil alone do not determine yield at the municipal level. It is very important to consider other factors, including the grapevine variety, clones and rootstocks that have been planted. Little is known about the yield potential of each variety and rootstock, particularly linked to growing conditions (Palacios *et al.*, 2022). In our study, the choice of varieties had less effect on the yield gap than environmental factors, with smaller differences found between the most grown varieties (i.e., 5-10 hl·ha<sup>-1</sup>·year<sup>-1</sup>) compared to within-zone yield variation (i.e., 10-30 hl·ha<sup>-1</sup>·year<sup>-1</sup>). In addition, their spatial distribution is quite homogeneous, with the nine most grown varieties planted in 75 to 93 % of the zone areas, as shown in Table ST3. In other case studies exhibiting large differences between varieties, the zoning method proposed here can also be applied by zoning per grapevine variety. Further research should be carried out on the yield potential of grapevine varieties, clones and rootstock, as well as their adaptation to different climate and soil conditions (Duchêne, 2016; Gisbert *et al.*, 2022; Serra *et al.*, 2014).

Vineyard technology and management practices beyond the scope of this study also have an important effect on grapevine yields. Irrigation facilities are available in 17.8 % of grapevine cultivated area in Languedoc-Roussillon region (Cambrea *et al.*, 2020), and many winegrowers considering turning to irrigation (Graveline and Grémont, 2021), thus helping to reduce grapevine yield gaps linked to water stress. Regarding vineyard inter-row management, different strategies exist already in Languedoc-Roussillon that can help to reduce competition between inter-row plants and the grapevine for water and soil nutrients (Fernández-Mena *et al.*, 2021). Soil fertility preservation practices by mineral fertilisation and the incorporation of compost into soils can enhance grapevine growth and grape production (Vrignon-Brenas *et al.*, 2019). The crop protection strategy to avoid the impact of pests and diseases also has an important effect on grapevine yields depending on the climate and pest pressure within each zone, powdery mildew being the most common disease in the region (Fouillet *et al.*, 2022; Guilpart, 2014).

For each of the seven zones, the yield gaps were divided into two groups representing i) high yield gaps (30-50 hl·ha<sup>-1</sup>·year<sup>-1</sup>), ii) low yield gaps (10-25 hl·ha<sup>-1</sup>·year<sup>-1</sup>). All the zones contained municipalities with high yield gaps (i.e., higher than 30 hl·ha<sup>-1</sup>·year<sup>-1</sup>), and municipalities without any yield gaps; in the latter case, irrigation in these municipalities may explain the closing of the yield gap. In addition, insight into good management practices could be gained by investigating the management practices of winegrowers who obtain high yields in low yield municipalities or zones, as did Andrade *et al.* (2022) and Mourtzinis *et al.* (2018). The exploration of such avenues is a future perspective for this research; for instance, by collecting extra data or by ‘innovation tracking’ (Salembier *et al.*, 2021).

Low grapevine yield has often been related to high quality wine and therefore yield has been restricted by quality labels to ensure a minimum quality standard (Stranieri and Tedeschi, 2019). The grapevine yield data used was provided by *Pays d'Oc* PGI, the most cultivated wine label in Languedoc-Roussillon region, which is a PGI (Protected Geographical Indication) label within the EU label framework (European Commission., 2022). Although other PGI labels exist in the region, as well as numerous *Protected Designation of Origin* labels (PDO) with more restricted vineyard practices and constrained target yields, the *Pays d'Oc* PGI label was more appropriate for studying a population of winegrowers and wineries with similar yield objectives. However, mixed label wineries are often located in low yield municipalities that apply business models that are not necessarily only based on the grapevine yield objective of this study.

## CONCLUSION

The present study proposes a method for selecting theoretical climate and soil factors that may have significant influence on grapevine yield at the municipality level. By working to such a scale, it is possible to gain more knowledge about winegrowing landscape characteristics that could contribute to future studies on vineyard management practices. Our analysis evidenced 6 relevant factors that explained grapevine yield at  $R^2 = 0.546$ , thus explaining only part of the grapevine yield. Further research should consider longer yield and the time-span of the climate database (e.g., 30 years) to improve the accuracy of the indicator selection. We opted to perform clustering to help analyse the types of municipality that have similar characteristics in terms of soil and climate. The choice to apply this zoning approach was also motivated by the fact that it provides a basis for formulating R&D recommendations. Based on statistical clustering carried out following the advice of regional wine label experts, we divided the Languedoc-Roussillon region into seven differential zones that had two contrasting yield gap levels associated with different combinations of indicators related to the limiting of grapevine yield. For each zone, we determined the extent to which the variability could be explained by pedoclimatic factors. Understanding the limiting factors linked to each zone could help local experts to implement adaptation measures in order to avoid or limit grapevine yield loss. In the present study, we showed that environmental factors at this scale can explain a small part of the annual variability of yield, but a large part (> 50 %) of average yield over time. Further research is needed to study the interactions between plant material and farming practices within each zone, as they may also play an important role in grapevine yield gaps at the regional scale.

## ACKNOWLEDGEMENTS

We thank all the *Pays d'Oc* PGI collaborators in this project, who provided funding, and shared data and expertise, in particular Laure Lacombe, Florence Barthès, Olivier Simonou, Jacques Gravegeal, Damien Onorré and

Gérard Bancillon. We highly appreciated the support of the Wine and Vine chair of the University of Montpellier, INRAE and Institut Agro for facilitating our partnership with the private sector and scientific exchanges.

## REFERENCES

- Anderson, J.D., Jones, G.V., Tait, A., Hall, A., Trought, M.C., (2012). Analysis of viticulture region climate structure and suitability in New Zealand. *OENO One* 46, 149–165. <https://doi.org/10.20870/oeno-one.2012.46.3.1515>
- Anderson, W., Johansen, C., Siddique, K.H., (2016). Addressing the yield gap in rainfed crops: a review. *Agron. Sustain. Dev.* 36, 18. <https://doi.org/10.1007/s13593-015-0341-y>
- Andrade, J.F., Mourtzinis, S., Edreira, J.I.R., Conley, S.P., Gaska, J., Kandel, H.J., Lindsey, L.E., Naeve, S., Nelson, S., Singh, M.P., (2022). Field validation of a farmer supplied data approach to close soybean yield gaps in the US North Central region. *Agric. Syst.* 200, 103434. <https://doi.org/10.1016/j.agry.2022.103434>
- Andrade, J.F., Rattalino Edreira, J.I., Farrow, A., van Loon, M.P., Craufurd, P.Q., Rurinda, J., Zingore, S., Chamberlin, J., Claessens, L., Adewopo, J., van Ittersum, M.K., Cassman, K.G., Grassini, P., (2019). A spatial framework for ex-ante impact assessment of agricultural technologies. *Glob. Food Secur.* 20, 72–81. <https://doi.org/10.1016/j.gfs.2018.12.006>
- Arnó, J., Rosell, J., Blanco, R., Ramos, M., Martínez-Casasnovas, J., (2012). Spatial variability in grape yield and quality influenced by soil and crop nutrition characteristics. *Precis. Agric.* 13, 393–410. <https://doi.org/10.1007/s11119-011-9254-1>
- Arrouays, D., McKenzie, N., Hempel, J., Richer de Forges, A., McBratney, A.B., (2019). GlobalSoilMap: basis of the global spatial soil information system. *CRC press*. <https://doi.org/10.1201/b16500>
- Ben-Ari, T., Boé, J., Ciais, P., Lecerf, R., Van der Velde, M., Makowski, D., (2018). Causes and implications of the unforeseen 2016 extreme yield loss in the breadbasket of France. *Nat. Commun.* 9, 1627. <https://doi.org/10.1038/s41467-018-04087-x>
- Bernardo, S., Dinis, L.-T., Machado, N., Moutinho-Pereira, J., (2018). Grapevine abiotic stress assessment and search for sustainable adaptation strategies in Mediterranean-like climates. A review. *Agron. Sustain. Dev.* 38, 1–20. <https://doi.org/10.1007/s13593-018-0544-0>
- Beza, E., Silva, J.V., Kooistra, L., Reidsma, P. (2017). Review of yield gap explaining factors and opportunities for alternative data collection approaches. *Eur. J. Agron., Farming systems analysis and design for sustainable intensification: new methods and assessments* 82, 206–222. <https://doi.org/10.1016/j.eja.2016.06.016>
- Bisson, L.F., Waterhouse, A.L., Ebeler, S.E., Walker, M.A., Lapsley, J.T., (2002). The present and future of the international wine industry. *Nature* 418, 696–699. <https://doi.org/10.1038/nature01018>
- Blanco-Ward, D., Ribeiro, A., Barreales, D., Castro, J., Verdial, J., Feliciano, M., Viceto, C., Rocha, A., Carlos, C., Silveira, C., (2019). Climate change potential effects on grapevine bioclimatic indices: A case study for the Portuguese demarcated Douro Region (Portugal). Presented at the BIO Web of Conferences, *EDP Sciences*. <https://doi.org/10.1051/bioconf/20191201013>
- Bois, B., Joly, D., Quénot, H., Pieri, P., Gaudillère, J.-P., Guyon, D., Saur, E., Van Leeuwen, C., (2018). Temperature-based

- zoning of the Bordeaux wine region. *OENO One* 52, 291–306. <https://doi.org/10.20870/oeno-one.2018.52.4.1580>
- Bonada, M., Phogat, V., Collins, C., Petrie, P.R., Sadras, V.O., (2022). Benchmarking water-limited yield potential and yield gaps of Shiraz in the Barossa and Eden Valleys. *bioRxiv*. <https://doi.org/10.1101/2022.10.25.513680>
- Bonasea, M., Rodriguez, M.C., Pinotti, L., Ferrero, S., (2015). Using multi-temporal Landsat imagery and linear mixed models for assessing water quality parameters in Río Tercero reservoir (Argentina). *Remote Sens. Environ.* 158, 28–41. <https://doi.org/10.1016/j.rse.2014.10.032>
- Bonfante, A., Brillante, L., (2022). Terroir analysis and its complexity: This article is published in cooperation with Terclim 2022 (XIVth International Terroir Congress and 2nd ClimWine Symposium), 3-8 July 2022, Bordeaux, France. *OENO One* 56, 375–388. <https://doi.org/10.20870/oenoone.2022.56.2.5448>
- Bonfatti, B.R., Hartemink, A.E., Giasson, E., Tornquist, C.G., Adhikari, K., (2016). Digital mapping of soil carbon in a viticultural region of Southern Brazil. *Geoderma* 261, 204–221. <https://doi.org/10.1016/j.geoderma.2015.07.016>
- Bramley, R. G. V., Ouzman, J., (2022). Underpinning terroir with data: on what grounds might subregionalisation of the Barossa Zone geographical indication be justified? *Aust. J. Grape Wine Res.* 28, 196–207. <https://doi.org/10.1111/ajgw.12513>
- Bramley, R. G. V., Ouzman, J., Boss, P. k., (2011). Variation in vine vigour, grape yield and vineyard soils and topography as indicators of variation in the chemical composition of grapes, wine and wine sensory attributes. *Aust. J. Grape Wine Res.* 17, 217–229. <https://doi.org/10.1111/j.17550238.2011.00136.x>
- Bramley, R.G.V., Ouzman, J., Trought, M.C.T., (2020). Making sense of a sense of place: precision viticulture approaches to the analysis of terroir at different scales: This article is published in cooperation with the XIIIth International Terroir Congress November 17-18 2020, Adelaide, Australia. Guest editors: Cassandra Collins and Roberta De Bei. *OENO One* 54, 903–917. <https://doi.org/10.20870/oeno-one.2020.54.4.3858>
- Brillante, L., Bonfante, A., Bramley, R.G.V., Tardaguila, J., Priori, S., (2020). Unbiased Scientific Approaches to the Study of Terroir Are Needed! *Front. Earth Sci.* 8. <https://doi.org/10.3389/feart.2020.539377>
- Buztepe, A., Kose, C., Kaya, O., (2017). Evaluation of cold tolerance of dormant buds according to position using thermal analysis in Karaerik (*V. vinifera* L.) grape. *Int J Res Rev* 4, 38–45.
- Cambrea, L., Lafon, C., Mayoux, L., (2020). En direct des territoires - L'irrigation de la vigne en ex-Languedoc-Roussillon : un potentiel de production maintenu par les économies d'eau. *Sci. Eaux Territ.* 28–33. <https://doi.org/10.14758/SET-REVUE.2020.5.05>
- Cameron, W., Petrie, P.R., Barlow, E.W.R., (2022). The effect of temperature on grapevine phenological intervals: Sensitivity of budburst to flowering. *Agric. For. Meteorol.* 315, 108841. <https://doi.org/10.1016/j.agrformet.2022.108841>
- Carbonneau, A., Deloire, A., Torregrosa, L., Jaillard, B., Pellegrino, A., Méta, A., Ojeda, H., Lebon, E., Abbal, P., (2015). *Traité de la vigne : physiologie, terroir, culture. Prat. Vitivinic. Dunod.* <https://hal.science/hal-02154373>
- Cardoso, A.S., Alonso, J., Rodrigues, A.S., Araújo-Paredes, C., Mendes, S., Valín, M.I., (2019). Agro-ecological terroir units in the North West Iberian Peninsula wine regions. *Appl. Geogr.* 107, 51–62. <https://doi.org/10.1016/j.apgeog.2019.03.011>
- Caubel, J., de Cortázar-Atauri, I.G., Launay, M., de Noblet-Ducoudré, N., Huard, F., Bertuzzi, P., Graux, A.I., (2015). Broadening the scope for ecoclimatic indicators to assess crop climate suitability according to ecophysiological, technical and quality criteria. *Agric. For. Meteorol.* 207, 94–106.
- Cayuela, L., (2018). Modelos lineales mixtos (LMM) y modelos lineales generalizados mixtos (GLMM) en R. Retrieved on 10 January 2020 on: <https://dl.orangedox.com/XpAzAQ0duvnoBmWZJ2/5-Modelos%20lineales%20mixtos%20en%20R.pdf>
- Chevigny, E., Quiquerez, A., Petit, C., Curmi, P., (2014). Lithology, landscape structure and management practice changes: Key factors patterning vineyard soil erosion at metre-scale spatial resolution. *Catena* 121, 354–364. <https://doi.org/10.1016/j.catena.2014.05.022>
- Coipel, J., Lovelle, B.R., Sipp, C., van Leeuwen, C., (2006). “Terroir” effect, as a result of environmental stress, depends more on soil depth than on soil type. *OENO One* 40, 177–185. <https://doi.org/10.20870/oeno-one.2006.40.4.867>
- Comtois, D., (2021). summarytools R package: Tools to Quickly and Neatly Summarize Data. Consulted on: <https://CRAN.R-project.org/package=summarytools>
- Cooper, M., Voss-Fels, K.P., Messina, C.D., Tang, T., Hammer, G.L., (2021). Tackling G× E× M interactions to close on-farm yield-gaps: creating novel pathways for crop improvement by predicting contributions of genetics and management to crop productivity. *Theor. Appl. Genet.* 1–20. <https://doi.org/10.1007/s00122-021-03812-3>
- Cremona, F., Tüvikene, L., Haberman, J., Nöges, P., Nöges, T., (2018). Factors controlling the three-decade long rise in cyanobacteria biomass in a eutrophic shallow lake. *Sci. Total Environ.* 621, 352–359. <https://doi.org/10.1016/j.scitotenv.2017.11.250>
- DDTM du GARD. (2022). Gel du 8 avril 2021 : procédure calamités agricoles. Consulted online the 10th of Mai 2022: <https://www.gard.gouv.fr/Politiques-publiques/Agriculture/Aidesagricoles/Calamites-agricoles2/Gel-du-8-avril-2021-procedure-calamites-agricoles>
- Droulia, F., Charalampopoulos, I., (2021). Future Climate Change Impacts on European Viticulture: A Review on Recent Scientific Advances. *Atmosphere* 12, 495. <https://doi.org/10.3390/atmos12040495>
- Duchêne, E., (2016). How can grapevine genetics contribute to the adaptation to climate change? *OENO One* 50, 113–124. <https://doi.org/10.20870/oeno-one.2016.50.3.98>
- Edreira, J.I.R., Cassman, K.G., Hochman, Z., van Ittersum, M.K., van Bussel, L., Claessens, L., Grassini, P., (2018). Beyond the plot: technology extrapolation domains for scaling out agronomic science. *Environ. Res. Lett.* 13, 054027. <https://doi.org/10.1088/1748-9326/aac092>
- Edreira, J.I.R., Mourtzinis, S., Conley, S.P., Roth, A.C., Ciampitti, I.A., Licht, M.A., Kandel, H., Kyveryga, P.M., Lindsey, L.E., Mueller, D.S., (2017). Assessing causes of yield gaps in agricultural areas with diversity in climate and soils. *Agric. For. Meteorol.* 247, 170–180. <https://doi.org/10.1016/j.agrformet.2017.07.010>
- European Commission. (2022). Aims of EU quality schemes. Consulted on 10th of May 2022: [https://agriculture.ec.europa.eu/farming/geographical-indications-and-quality-schemes/geographical-indications-and-quality-schemes-explained\\_en](https://agriculture.ec.europa.eu/farming/geographical-indications-and-quality-schemes/geographical-indications-and-quality-schemes-explained_en)
- Fernández-Mena, H., (2022). Dataset containing average climate, soil, and IGP Pays d’Oc grapevine yield in municipalities of Languedoc-Roussillon from 2010-2018. <https://doi.org/10.57745/THCVRE>
- Fernández-Mena, H., Frey, H., Celette, F., Garcia, L., Barkaoui, K., Hossard, L., Naulleau, A., Métral, R., Gary, C., Méta, A., (2021). Spatial and temporal diversity of service plant management strategies across vineyards in the south of France. *Analysis*

- through the Coverage Index. *Eur. J. Agron.* 123, 126191. <https://doi.org/10.1016/j.eja.2020.126191>
- Ferretti, C.G., (2019). Relationship between the geology, soil assessment, and terroir of Gewürtztraminer vineyards: A case study in the Dolomites of northern Italy. *Catena* 179, 74–84. <https://doi.org/10.1016/j.catena.2019.03.044>
- Fouillet, E., Delière, L., Chartier, N., Munier-Jolain, N., Cortel, S., Rapidel, B., Mérot, A., (2022). Reducing pesticide use in vineyards. Evidence from the analysis of the French DEPHY network. *Eur. J. Agron.* 136, 126503. <https://doi.org/10.1016/j.eja.2022.126503>
- Fraga, H., Malheiro, A., Moutinho-Pereira, J., Jones, G., Alves, F., Pinto, J.G., Santos, J., (2014a). Very high resolution bioclimatic zoning of Portuguese wine regions: present and future scenarios. *Reg. Environ. Change* 14, 295–306. <https://doi.org/10.1007/s10113-013-0490-y>
- Fraga, H., Malheiro, A.C., Moutinho-Pereira, J., Cardoso, R.M., Soares, P.M., Cancela, J.J., Pinto, J.G., Santos, J.A., (2014b). Integrated analysis of climate, soil, topography and vegetative growth in Iberian viticultural regions. *PLoS One* 9. <https://doi.org/10.1371/journal.pone.0108078>
- Fraga, H., García de Cortázar Atauri, I., Malheiro, A.C., Santos, J.A., (2016). Modelling climate change impacts on viticultural yield, phenology and stress conditions in Europe. *Glob. Change Biol.* 22, 3774–3788. <https://doi.org/10.1111/gcb.13382>
- Fraga, H., Molitor, D., Leolini, L., Santos, J.A., (2020). What Is the Impact of Heatwaves on European Viticulture? A Modelling Assessment. *Appl. Sci.* 10, 3030. <https://doi.org/10.3390/app10093030>
- Galet, P., (2000). Précis de viticulture : à l'usage des ingénieurs agronomes, des étudiants en agronomie et en oenologie, des techniciens des lycées agricoles et des professionnels de la viticulture, Dehan, Montpellier, 602p. ed.
- Garcia de Cortazar Atauri, I., Duchêne, E., Destrac, A., Barbeau, G., De Resseguier, L., Lacombe, T., Parker, A.K., Saurin, N., van Leeuwen, C., (2017). Grapevine phenology in France: from past observations to future evolutions in the context of climate change. *OENO One*, 51(2), 115-126. <https://doi.org/10.20870/oenone.2017.51.2.1622>
- Gareth, J., Daniela, W., Trevor, H., Robert, T., (2013). An introduction to statistical learning: with applications in R. Springer.
- Garnier, E., (2010). Exceptionnal Meanness Water and Hot Weather 500 years of drought and heat wave in France and neighboring countries. *Houille Blanche* 4, 26–42. <https://doi.org/10.1051/lhb/2010039>
- Gaudin, R., Kansou, K., Payan, J.-C., Pellegrino, A., Gary, C., (2014). A water stress index based on water balance modelling for discrimination of grapevine quality and yield. *J. Int. Sci. Vigne Vin* 48, 1–9. <https://doi.org/10.20870/oenone.2014.48.1.1655>
- Gavrilescu, C., Richard, Y., Joly, D., Bois, B., (2018). Climate zoning of the Burgundy winegrowing region. Presented at the E3S Web of Conferences, EDP Sciences, p. 01003.
- Gerós, H., Chaves, M.M., Gil, H.M., Delrot, S., (2015). Grapevine in a Changing Environment: A Molecular and Ecophysiological Perspective. John Wiley & Sons.
- González-Fernández, E., Piña-Rey, A., Fernández-González, M., Aira, M.J., Rodríguez-Rajo, F.J., (2020). Prediction of grapevine yield based on reproductive variables and the influence of meteorological conditions. *Agronomy* 10, 714. <https://doi.org/10.3390/agronomy10050714>
- Graveline, N., Grémont, M., (2021). The role of perceptions, goals and characteristics of wine growers on irrigation adoption in the context of climate change. *Agric. Water Manag.* 250, 106837. <https://doi.org/10.1016/j.agwat.2021.106837>
- Greer, D., Weedon, M., (2014). Temperature-dependent responses of the berry developmental processes of three grapevine (*Vitis vinifera*) cultivars. *N. Z. J. Crop Hortic. Sci.* 42, 233–246. <https://doi.org/10.1080/01140671.2014.894921>
- Grimaldi, J., Wang, Y.-J., Gastellu-Etchegorry, J.-P., Chavanon, E., Lauret, N., Guilleux, J., Bustillo, V., Houet, T., (2019). What are the impacts of tree shade on the absorption of light by grapevine in alleycropped vineyards? Presented at the 4th World Congress on Agroforestry.
- Guilpart, N., (2014). Relations entre services écosystémiques dans un agroécosystème à base de plantes pérennes : compromis entre rendement de la vigne et régulation de l'oïdium.
- Guilpart, N., Grassini, P., Sadras, V.O., Timsina, J., Cassman, K.G., (2017). Estimating yield gaps at the cropping system level. *Field Crops Res.* 206, 21–32. <https://doi.org/10.1016/j.fcr.2017.02.008>
- Guilpart, N., Metay, A., Gary, C., (2014). Grapevine bud fertility and number of berries per bunch are determined by water and nitrogen stress around flowering in the previous year. *Eur. J. Agron.* 54, 9–20. <https://doi.org/10.1016/j.eja.2013.11.002>
- Hannin, H., Touzard, J.-M., Aigrain, P., Bois, B., Brugière, F., Duchêne, E., de Cortázar-Atauri, I.G., Gautier, J., Héraud, E.G., Hammond, R., (2021). Le vignoble français face au changement climatique: l'élaboration d'une stratégie d'adaptation à partir de scénarios de prospective. <https://hal.inrae.fr/hal-03141778>
- Hendrickson, L., Ball, M., Wood, J., Chow, W., Furbank, R.T., (2004). Low temperature effects on photosynthesis and growth of grapevine. *Plant Cell Environ.* 27, 795–809. <https://doi.org/10.1111/j.1365-3040.2004.01184.x>
- Himelrick, D.G., (1991). Growth and nutritional responses of nine grape cultivars to low soil pH. *Hort. Science* 26, 269–271. <https://doi.org/10.21273/HORTSCI.26.3.269>
- Hothorn, T., Bretz, F., Hothorn, M.T., (2009). The multcomp package. *R Proj. Stat. Comput.*
- Huglin, M., (1978). Nouveau mode d'évaluation des possibilités héliothermiques d'un milieu viticole.
- IGN. (2020). Occupation du sol - usages - Viticulture. Geoportail. Consulted the 3th July 2020 on <https://www.geoportail.gouv.fr/>.
- Jackson, D., Lombard, P., (1993). Environmental and management practices affecting grape composition and wine quality—a review. *Am. J. Enol. Vitic.* 44, 409–430. <https://doi.org/10.5344/ajev.1993.44.4.409>
- Kaya, Ö., Köse, C., (2017). Determination of resistance to low temperatures of winter buds on lateral shoot present in Karaerik (*Vitis vinifera* L.) grape cultivar. *Acta Physiol. Plant.* 39, 1–9. <https://doi.org/10.1007/s11738-017-2513-7>
- Lagacherie, P., Coulouma, G., Ariagno, P., Virat, P., Boizard, H., Richard, G., (2006). Spatial variability of soil compaction over a vineyard region in relation with soils and cultivation operations. *Geoderma* 134, 207–216. <https://doi.org/10.1016/j.geoderma.2005.10.006>
- Le Roux, R., De Resseguier, L., Corpetti, T., Jégou, N., Madelin, M., van Leeuwen, C., Quénot, H., (2017). Comparison of two fine scale spatial models for mapping temperatures inside winegrowing areas. *Agric. For. Meteorol.* 247, 159–169. <https://doi.org/10.1016/j.agrformet.2017.07.020>
- Lê, S., Josse, J., Husson, F., (2008). FactoMineR: an R package for multivariate analysis. *J. Stat. Softw.* 25, 1– 18. <https://doi.org/10.18637/jss.v025.i01>

- Leolini, L., Moriondo, M., Fila, G., Costafreda-Aumedes, S., Ferrise, R., Bindi, M., (2018). Late spring frost impacts on future grapevine distribution in Europe. *Field Crops Res.* 222, 197–208. <https://doi.org/10.1016/j.fcr.2017.11.018>
- Liu, B., Chen, X., Meng, Q., Yang, H., van Wart, J., (2017). Estimating maize yield potential and yield gap with agro-climatic zones in China—Distinguish irrigated and rainfed conditions. *Agric. For. Meteorol.* 239, 108–117. <https://doi.org/10.1016/j.agrformet.2017.02.035>
- Liu, G.-T., Jiang, J.-F., Liu, X.-N., Jiang, J.-Z., Sun, L., Duan, W., Li, R.-M., Wang, Y., Lecourieux, D., Liu, C.-H., (2019). New insights into the heat responses of grape leaves via combined phosphoproteomic and acetylproteomic analyses. *Hortic. Res.* 6, 1–15. <https://doi.org/10.1038/s41438-019-0183-x>
- Lopez-Fornieles, E., Brunel, G., Devaux, N., Roger, J.-M., Tisseire, B., (2022). Is It Possible to Assess Heatwave Impact on Grapevines at the Regional Level with Time Series of Satellite Images? *Agronomy* 12, 563. <https://doi.org/10.3390/agronomy12030563>
- Lovisol, C., Lavoie-Lamoureux, A., Tramontini, S., Ferrandino, A., (2016). Grapevine adaptations to water stress: new perspectives about soil/plant interactions. *Theor. Exp. Plant Physiol.* 28, 53–66. <https://doi.org/10.1007/s40626-016-0057-7>
- Lüdecke, D., (2021). sjPlot: Data Visualization for Statistics in Social Science. R package version 2.8.10. <https://CRAN.R-project.org/package=sjPlot>
- Luo, H.-B., Ma, L., Xi, H.-F., Duan, W., Li, S.-H., Loescher, W., Wang, J.-F., Wang, L.-J., (2011). Photosynthetic responses to heat treatments at different temperatures and following recovery in grapevine (*Vitis amurensis* L.) leaves. *PLoS One* 6, e23033. <https://doi.org/10.1371/journal.pone.0023033>
- Molitor, D., Caffarra, A., Sinigoj, P., Pertot, I., Hoffmann, L., Junk, J., (2014). Late frost damage risk for viticulture under future climate conditions: a case study for the Luxembourgish winegrowing region. *Aust. J. Grape Wine Res.* 20, 160–168.
- Moral, F.J., Rebollo, F.J., Paniagua, L.L., García, A., de Salazar, E.M., (2016). Application of climatic indices to analyse viticultural suitability in Extremadura, south-western Spain. *Theor. Appl. Climatol.* 123, 277–289. <https://doi.org/10.1007/s00704-014-1363-0>
- Morales-Castilla, I., de Cortázar-Atauri, I.G., Cook, B.I., Lacombe, T., Parker, A., van Leeuwen, C., Nicholas, K.A., Wolkovich, E.M., (2020). Diversity buffers winegrowing regions from climate change losses. *Proc. Natl. Acad. Sci.* 117, 2864–2869. <https://doi.org/10.1073/pnas.1906731117>
- Mosedale, J.R., Abernethy, K.E., Smart, R.E., Wilson, R.J., Maclean, I.M., (2016). Climate change impacts and adaptive strategies: lessons from the grapevine. *Glob. Change Biol.* 22, 3814–3828. <https://doi.org/10.1111/gcb.13406>
- Mourtzinis, S., Edreira, J.I.R., Grassini, P., Roth, A.C., Casteel, S.N., Ciampitti, I.A., Kandel, H.J., Kyveryga, P.M., Licht, M.A., Lindsey, L.E., (2018). Sifting and winnowing: Analysis of farmer field data for soybean in the US North-Central region. *Field Crops Res.* 221, 130–141. <https://doi.org/10.1016/j.fcr.2018.02.024>
- Naulleau, A., Gary, C., Prévot, L., Berteloot, V., Fabre, J.-C., Crevoisier, D., Gaudin, R., Hossard, L., (2022). Participatory modeling to assess the impacts of climate change in a Mediterranean vineyard watershed. *Environ. Model. Softw.* 150, 105342. <https://doi.org/10.1016/j.envsoft.2022.105342>
- Naulleau, A., Gary, C., Prévot, L., Hossard, L., (2021). Evaluating Strategies for Adaptation to Climate Change in Grapevine Production—A Systematic Review. *Front. in Plant Sc.* <https://doi.org/10.3389/fpls.2020.607859>
- Oliva Oller, P., Notaro, M., Langer, E., Gary, C., (2022). Structure and management of traditional agroforestry vineyards in the high valleys of southern Bolivia. *Agrofor. Syst.* 96, 375–386. <https://doi.org/10.1007/s10457-021-00725-4>
- Ollat, N., van Leeuwen, C., de Cortazar Aauri, I.G., Touzard, J.-M., (2017). The challenging issue of climate change for sustainable grape and wine production. *OENO One* 51, 59–60. <https://doi.org/10.20870/oeno-one.2017.51.2.1872>
- Ollat, N., Zito, S., Richard, Y., Aigrain, P., Brugière, F., Duchêne, E., De Cortazar-Atauri, I.G., Gautier, J., Giraud-Héraud, E., Hannin, H., (2021). La diversité des vignobles français face au changement climatique: simulations climatiques et prospective participative. *Climatologie* 18, 3. <https://doi.org/10.1051/climat/202118003>
- Pagay, V., Collins, C., (2017). Effects of timing and intensity of elevated temperatures on reproductive development of field-grown Shiraz grapevines. *OENO One* 51. <https://doi.org/10.20870/oeno-one.2017.51.4.1066>
- Palacios, F., Diago, M. P., Melo-Pinto, P., & Tardaguila, J. (2022). Early yield prediction in different grapevine varieties using computer vision and machine learning. *Precision Agriculture*, 1-29. <https://doi.org/10.3390/agronomy12102463>
- Parker, A., de Cortázar-Atauri, I.G., Chuine, I., Barbeau, G., Bois, B., Boursiquot, J.-M., Cahurel, J.-Y., Claverie, M., Dufourcq, T., GénY, L., (2013). Classification of varieties for their timing of flowering and veraison using a modelling approach: A case study for the grapevine species *Vitis vinifera* L. *Agric. For. Meteorol.* 180, 249–264. <https://doi.org/10.1016/j.agrformet.2013.06.005>
- Parker, A.K., de Cortázar-Atauri, I.G., GénY, L., Spring, J.-L., Destrac, A., Schultz, H., Molitor, D., Lacombe, T., Graça, A., Monamy, C., (2020). Temperature-based grapevine sugar ripeness modelling for a wide range of *Vitis vinifera* L. cultivars. *Agric. For. Meteorol.* 285, 107902. <https://doi.org/10.1016/j.agrformet.2020.107902>
- Pellegrino, A., Lebon, E., Simonneau, T., Wery, J., (2005). Towards a simple indicator of water stress in grapevine (*Vitis vinifera* L.) based on the differential sensitivities of vegetative growth components. *Aust. J. Grape Wine Res.* 11, 306–315. <https://doi.org/10.1111/j.1755-0238.2005.tb00030>
- Peng, Y., Roell, Y.E., Odgers, N.P., Møller, A.B., Beucher, A., Greve, M.B., Greve, M.H., (2021). Mapping and describing natural terroir units in Denmark. *Geoderma* 394, 115014. <https://doi.org/10.1016/j.geoderma.2021.115014>
- Petoumenou, D.G., Biniari, K., Xyrafis, E., Mavronasios, D., Daskalakis, I., Palliotti, A., (2019). Effects of Natural Hail on the Growth, Physiological Characteristics, Yield, and Quality of *Vitis vinifera* L. cv. Thompson Seedless under Mediterranean Growing Conditions. *Agronomy* 9, 197. <https://doi.org/10.3390/agronomy9040197>
- Petrie, P.R., Clingeleffer, P.R., (2005). Effects of temperature and light (before and after budburst) on inflorescence morphology and flower number of Chardonnay grapevines (*Vitis vinifera* L.). *Aust. J. Grape Wine Res.* 11, 59–65. <https://doi.org/10.1111/j.1755-0238.2005.tb00279>
- Pinheiro, J. C., and Bates, D. M., (2006). Mixed-Effects Models in S and S-PLUS. *Springer science & business media.* <https://doi.org/10.1007/b98882>
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., R Core Team, (2021). nlme: Linear and Nonlinear Mixed Effects Models. R package version 3.1-152, <https://CRAN.R-project.org/package=nlme>.
- Poling, E.B., (2008). Spring cold injury to winegrapes and protection strategies and methods. *Hort. Science* 43, 1652–1662. <https://doi.org/10.21273/HORTSCI.43.6.1652>

- Poni, S., Gatti, M., Palliotti, A., Dai, Z., Duchêne, E., Truong, T.-T., Ferrara, G., Matarrese, A.M.S., Gallotta, A., Bellincontro, A., (2018). Grapevine quality: A multiple choice issue. *Sci. Hort.* 234, 445–462. <https://doi.org/10.1016/j.scienta.2017.12.035>
- Pradhan, P., Fischer, G., van Velthuizen, H., Reusser, D.E., Kropp, J.P., (2015). Closing yield gaps: how sustainable can we be? *PLoS One* 10, e0129487. <https://doi.org/10.1371/journal.pone.0129487>
- Priori, S., Pellegrini, S., Perria, R., Puccioni, S., Storchi, P., Valboa, G., Costantini, E.A.C., (2019). Scale effect of terroir under three contrasting vintages in the Chianti Classico area (Tuscany, Italy). *Geoderma* 334, 99–112. <https://doi.org/10.1016/j.geoderma.2018.07.048>
- Quénot, H., de Cortazar Atauri, I.G., Bois, B., Sturman, A., Bonnardot, V., Le Roux, R., (2017). Which climatic modeling to assess climate change impacts on vineyards? *OENO One* 5, 91–97. <https://doi.org/10.20870/oeno-one.2017.51.2.1869>
- Quintana-Seguí, P., Le Moigne, P., Durand, Y., Martin, E., Habets, F., Baillon, M., Canellas, C., Franchisteguy, L., Morel, S., (2008). Analysis of near-surface atmospheric variables: Validation of the SAFRAN analysis over France. *J. Appl. Meteorol. Climatol.* 47, 92–107. <https://doi.org/10.1175/2007JAMC1636.1>
- R Core Team. (2014). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- Resco, P., Iglesias, A., Bardají, I., Sotés, V., (2016). Exploring adaptation choices for grapevine regions in Spain. *Reg. Environ. Change* 16, 979–993. <https://doi.org/10.1007/s10113-015-0811-4>
- Riou, C., Agostini, D., Aigrain, P., Barthe, M., des Robert, M.-L., Gervais, J.-P., Jobard, E., Lurton, L., Moncomble, D., Prêtet-Lataste, C., (2016). Action plan against declining vineyards: An innovative approach. Presented at the BIO Web of Conferences, *EDP Sciences*, p. 01040. <https://doi.org/10.1051/bioconf/20160701040>
- Rodrigo-Comino, J., Cerdà, A., (2018). Improving stock unearthing method to measure soil erosion rates in vineyards. *Ecol. Indic.* 85, 509–517. <https://doi.org/10.1016/j.ecolind.2017.10.042>
- Salembier, C., Segrestin, B., Weil, B., Jeuffroy, M.-H., Cadoux, S., Cros, C., Favrelière, E., Fontaine, L., Gimaret, M., Noilhan, C., (2021). A theoretical framework for tracking farmers' innovations to support farming system design. *Agron. Sustain. Dev.* 41, 1–23. <https://doi.org/10.1007/s13593-021-00713-z>
- Santillán, D., Iglesias, A., La Jeunesse, I., Garrote, L., Sotés, V., (2019). Vineyards in transition: A global assessment of the adaptation needs of grape producing regions under climate change. *Sci. Total Environ.* 657, 839–852. <https://doi.org/10.1016/j.scitotenv.2018.12.079>
- Santos, J.A., Costa, R., Fraga, H., (2019). New insights into thermal growing conditions of Portuguese grapevine varieties under changing climates. *Theor. Appl. Climatol.* 135, 1215–1226. <https://doi.org/10.1007/s00704-018-2443-3>
- Santos, J.A., Malheiro, A.C., Karremann, M.K., Pinto, J.G., (2011). Statistical modelling of grapevine yield in the Port Wine region under present and future climate conditions. *Int. J. Biometeorol.* 55, 119–131. <https://doi.org/10.1007/s00484-010-0318-0>
- Schauberger, B., Ben-Ari, T., Makowski, D., Kato, T., Kato, H., Ciaï, P., (2018). Yield trends, variability and stagnation analysis of major crops in France over more than a century. *Sci. Rep.* 8, 1–12. <https://doi.org/10.1038/s41598-018-35351-1>
- Schultz, H., (2000). Climate change and viticulture: a European perspective on climatology, carbon dioxide and UV-B effects. *Aust. J. Grape Wine Res.* 6, 2–12. <https://doi.org/10.1038/s41598-018-35351-1>
- Serra, I., Strever, A., Myburgh, P., Deloire, A., (2014). The interaction between rootstocks and cultivars (*Vitis vinifera* L.) to enhance drought tolerance in grapevine. *Aust. J. Grape Wine Res.* 20, 1–14. <https://doi.org/10.1111/ajgw.12054>
- Sgubin, G., Swingedouw, D., Dayon, G., de Cortazar-Atauri, I.G., Ollat, N., Pagé, C., van Leeuwen, C., (2018). The risk of tardive frost damage in French vineyards in a changing climate. *Agric. For. Meteorol.* 250, 226–242. <https://doi.org/10.1016/j.agrformet.2017.12.253>
- Silva, J.V., Reidsma, P., Laborte, A.G., Van Ittersum, M.K., (2017). Explaining rice yields and yield gaps in Central Luzon, Philippines: An application of stochastic frontier analysis and crop modelling. *Eur. J. Agron.* 82, 223–241. <https://doi.org/10.1016/j.eja.2016.06.017>
- Simonneau, T., Lebon, E., Coupel-Ledru, A., Marguerit, E., Rosse Deutsch, L., Ollat, N., (2017). Adapting plant material to face water stress in vineyards: which physiological targets for an optimal control of plant water status? *OENO One* 51, 167. <https://doi.org/10.20870/oeno-one.2017.51.2.1870>
- Stranieri, S., Tedeschi, P., (2019). The Role of Quality in Wine Production and Market: European Rules, CAP and New Technology, in: A History of Wine in Europe, 19th to 20th Centuries, Volume II. Springer, pp. 255–274. [https://doi.org/10.1007/978-3-030-27794-9\\_11](https://doi.org/10.1007/978-3-030-27794-9_11)
- Strobl, C., Boulesteix, A.-L., Kneib, T., Augustin, T., Zeileis, A., (2008). Conditional variable importance for random forests. *BMC Bioinformatics* 9, 1–11. <https://doi.org/10.1186/1471-2105-9-307>
- Styc, Q., Lagacherie, P., (2021). Uncertainty assessment of soil available water capacity using error propagation: A test in Languedoc-Roussillon. *Geoderma* 391, 114968. <https://doi.org/10.1016/j.geoderma.2021.114968>
- Styc, Q., Lagacherie, P., (2019). What is the best inference trajectory for mapping soil functions: an example of mapping soil available water capacity over Languedoc Roussillon (France). *Soil Syst.* 3, 34. <https://doi.org/10.3390/soilsystems3020034>
- Tagliavini, M., Rombola, A.D., (2001). Iron deficiency and chlorosis in orchard and vineyard ecosystems. *Eur. J. Agron.* 15, 71–92. [https://doi.org/10.1016/S1161-0301\(01\)00125-3](https://doi.org/10.1016/S1161-0301(01)00125-3)
- Tintinger, M., (2020). Wine estates competitiveness and resilience in Languedoc-Roussillon in the context of climate change. Master Thesis submitted for the obtention of Agricultural Engineer degree at Bordeaux Sciences Agro.
- Tonietto, J., Carbonneau, A., (2004). A multicriteria climatic classification system for grape-growing regions worldwide. *Agric. For. Meteorol.* 124, 81–97. <https://doi.org/10.1016/j.agrformet.2003.06.001>
- Tonietto, J., Sotés Ruiz, V., Zanús, M.C., Montes, C., Uliarte, E., Bruno, L., Climaco, P., Pena, A., Guerra, C.C., Catania, C., (2014). The Effect of Viticultural Climate on Red and White Wine Typicity Characterization in Ibero-American grape-growing regions. *Embrapa Uva E Vinho-Artigo Em Periód.* <http://www.alice.cnptia.embrapa.br/alice/handle/doc/986819>
- Torregrosa, L., Bigard, A., Doligez, A., Lecourieux, D., Rienth, M., Luchaire, N., Pieri, P., Chatbanyong, R., Shahood, R., Farnós, M., (2017). Developmental, molecular and genetic studies on grapevine response to temperature open breeding strategies for adaptation to warming. *OENO One* 51, 155–165. <https://doi.org/10.20870/oeno-one.2017.51.2.1587>
- Touzard, J.-M., Aigrain, P., Bois, B., Duchêne, E., de Cortazar Atauri, I.G., Giraud-Heraud, E., Gautier, J., Hannin, H., Ollat, N., (2017). Lessons from a Prospective on the French Wine Industry Under Climate Change (2050). Presented at the 4. Global Science Conference On Climate Smart Agriculture, pp. 20-vues.

- Trought, M., Howell, G., Cherry, N.J., (1999). Practical considerations for reducing frost damage in vineyards. <https://hdl.handle.net/10182/4236>
- Van Ittersum, M.K., Cassman, K.G., Grassini, P., Wolf, J., Titttonell, P., Hochman, Z., (2013). Yield gap analysis with local to global relevance—a review. *Field Crops Res.* 143, 4–17. <https://doi.org/10.1016/j.fcr.2012.09.009>
- van Leeuwen, C., Bois, B., (2018). Update in unified terroir zoning methodologies, in: E3S Web of Conferences. *EDP Sciences*, Zaragoza, Spain, p. 01044. <https://doi.org/10.1051/e3sconf/20185001044>
- van Leeuwen, C., Roby, J.-P., de Rességuier, L., (2018). Soil-related terroir factors: a review. *OENO One* 52, 173–188. <https://doi.org/10.20870/oeno-one.2018.52.2.2208>
- van Leeuwen, C., Roby, J.-P., Pernet, D., Bois, B., (2010). Methodology of soil-based zoning for viticultural terroirs. *Bull. OIV* 83, 13.
- Van Wart, J., van Bussel, L.G., Wolf, J., Licker, R., Grassini, P., Nelson, A., Boogaard, H., Gerber, J., Mueller, N.D., Claessens, L., (2013). Use of agro-climatic zones to upscale simulated crop yield potential. *Field Crops Res.* 143, 44–55. <https://doi.org/10.1016/j.fcr.2012.11.023>
- Vaudour, E., Costantini, E., Jones, G.V., Mocali, S., (2015). An overview of the recent approaches to terroir functional modelling, footprinting and zoning. *SOIL* 1, 287–312. <https://doi.org/10.5194/soil-1287-2015>
- Vautard, R., van Oldenborgh, G.J., Bonnet, R., Li, S., Robin, Y., Kew, S., Philip, S., Soubeyroux, J.-M., Dubuisson, B., Viovy, N., Reichstein, M., Otto, F., Garcia de Cortazar-Atauri, I., (2022). Human influence on growing-period frosts like the early April 2021 in Central France. *Nat. Hazards Earth Syst. Sci. Discuss.* 1–25. <https://doi.org/10.5194/nhess-2022-41>
- Vaysse, K., Lagacherie, P., (2017). Using quantile regression forest to estimate uncertainty of digital soil mapping products. *Geoderma* 291, 55–64. <https://doi.org/10.1016/j.geoderma.2016.12.017>
- Vaysse, K., Lagacherie, P., (2015). Evaluating digital soil mapping approaches for mapping GlobalSoilMap soil properties from legacy data in Languedoc-Roussillon (France). *Geoderma Reg.* 4, 20–30. <https://doi.org/10.1016/j.geodrs.2014.11.003>
- Venios, X., Korkas, E., Nisiotou, A., Banilas, G., (2020). Grapevine Responses to Heat Stress and Global Warming. *Plants* 9, 1754. <https://doi.org/10.3390/plants9121754>
- Vicente-Serrano, S.M., Beguería, S., López-Moreno, J.I., (2010). A multiscalar drought index sensitive to global warming: the standardized precipitation evapotranspiration index. *J. Clim.* 23, 1696–1718. <https://doi.org/10.1175/2009JCLI2909.1>
- Vidal, J.-P., Martin, E., Franchistéguy, L., Baillon, M., Soubeyroux, J.-M., (2010). A 50-year high-resolution atmospheric reanalysis over France with the Safran system. *Int. J. Climatol.* 30, 1627–1644. <https://doi.org/10.1002/joc.2003>
- Villalobos-Soublett, E., Gutiérrez-Gamboa, G., Balbontín, C., Zurita-Silva, A., Ibacache, A., Verdugo Vázquez, N., (2021). Effect of Shading Nets on Yield, Leaf Biomass and Petiole Nutrients of a Muscat of Alexandria Vineyard Growing under Hyper-Arid Conditions. *Horticulturae* 7, 445. <https://doi.org/10.3390/horticulturae7110445>
- Vrignon-Brenas, S., Metay, A., Laporatti, R., Gharibi, S., Fraga, A., Dauzat, M., Rolland, G., Pellegrino, A., (2019). Gradual responses of grapevine yield components and carbon status to nitrogen supply. *OENO One* 53, 289–306. <https://doi.org/10.20870/oeno-one.2019.53.2.2431>
- Vrsic, S., Kocsis, L., Pulko, B., (2016). Influence of substrate pH on root growth, biomass and leaf mineral contents of grapevine rootstocks grown in pots. [https://www.sid.ir/en/VEWSSID/J\\_pdf/84820160216.pdf](https://www.sid.ir/en/VEWSSID/J_pdf/84820160216.pdf)
- Wei, T., and Simko, V., (2017). R package “corrplot”: Visualization of a Correlation Matrix (Version 0.84). Available from <https://github.com/taiyun/corrplot>
- White, R.E., (2020). The value of soil knowledge in understanding wine terroir. *Front. Environ. Sci.* 8, 12. <https://doi.org/10.3389/fenvs.2020.00012>
- Wickham, H., (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York, 2016. <https://cran.microsoft.com/snapshot/2015-01-06/web/packages/ggplot2/ggplot2.pdf>
- Wickham, H., Romain François, Lionel Henry, Kirill Müller, (2018). *dplyr: A Grammar of Data Manipulation*. R package version. <https://cran.r-project.org/package=dplyr>
- Wikialine. (2009). Languedoc-Roussillon region locator map. Creative Commons 3.3 license. Downloaded the 5th October 2019 on: [https://fr.wikipedia.org/wiki/Fichier:Languedoc-Roussillon\\_region\\_locator\\_map.svg](https://fr.wikipedia.org/wiki/Fichier:Languedoc-Roussillon_region_locator_map.svg)
- Williams, L.E., Levin, A.D., Fidelibus, M.W., (2022). Crop coefficients (Kc) developed from canopy shaded area in California vineyards. *Agric. Water Manag.* 271, 107771. <https://doi.org/10.1016/j.agwat.2022.107771>
- Xiao, F., Yang, Z., Lee, K., (2017). Photosynthetic and physiological responses to high temperature in grapevine (*Vitis vinifera* L.) leaves during the seedling stage. *J. Hortic. Sci. Biotechnol.* 92, 2–10. <https://doi.org/10.1080/14620316.2016.1211493>
- Zuur, A.F., Ieno, E.N., Walker, N., Saveliev, A.A., Smith, G.M., (2009). *Mixed effects models and extensions in ecology with R, Statistics for Biology and Health*. Springer New York. <https://doi.org/10.1007/978-0-387-87458-6>