

# In vitro antiprotozoal action of 7 spices, medicinal plants and essential oils against 3 stages of *Eimeria tenella*

C. GIRARD<sup>(1)</sup>, A. SILVESTRE<sup>(2)</sup>, T. CHABRILLAT<sup>(3)</sup>, S. KERROS<sup>(1)</sup>  
<sup>(1)</sup>PHYTOSYNTHESE, F-63200 MOZAC, FRANCE  
<sup>(2)</sup>INRAE, UNIVERSITÉ DE TOURS, ISP, F-37380 NOUZILLY, FRANCE


Growing societal concerns to replace coccidiostats in poultry farms have led to the use of alternatives such as plant bio-actives. To investigate the mode of action of natural molecules, we designed this *in vitro* study to screen anti-protozoal activity of 4 spices or medicinal plants described by non-volatile active molecules (MedP) and 3 essential oils (EO) against 3 different stages of *E. tenella*.

## MATERIAL AND METHODS

- Natural products tested:**
  - **MedP:** Fabacea powder, Urticaceae extract, Punicaceae extract, Acanthaceae extract
  - **EO:** Lilaceae EO, Lamiaceae EO, Lauraceae EO
- Negative control:** H<sub>2</sub>O, soybean hemicellulose, DMSO
- Protozoal strain:** *Eimeria tenella* recombined strain
- 3 experimental steps after cytotoxicity assay (colorimetric MTS test, 1.5·10<sup>4</sup> MDBK seed, incubation 5h at 37°C):
  - 1** Direct anti-protozoite activity test: incubation 41°C for 1h. Killed sporozoites were stained with Evans Blue Dye. Each sample tested in triplicate. **Viability rate with fluorescence microscopy as % viability = number of live parasite in tested sample / number of live parasite in negative control\*100**
  - 2** Invasion inhibition test: 96 microplates: 1.5·10<sup>4</sup> MDBK cells, 1.5·10<sup>4</sup> *E. tenella* sporozoites expressed enzymatic activity; substrate added for 2h at 37°C, 4 replicates per tested sample. **% invasion = absorbance (550 nm) tested sample / absorbance (550 nm) negative control\*100**
  - 3** Sporulation rate test: 5.10<sup>5</sup> non-sporulated oocysts (in 2% potassium dichromate solution), incubation for 72h at 26°C; each sample tested in triplicate. **% sporulation = number of sporulated oocysts in tested sample / number of sporulated oocyst in negative control\*100**
- Statistical test:** Mann-whitney test compared to negative control for each test (p<0.05).


## RESULTS AND DISCUSSION

The 4 MedP samples (Fabacea powder, Urticaceae extract, Punicaceae extract, Acanthaceae extract) were less cytotoxic (lethal dose 10% (LD10) than the 3 EO samples (Lilaceae EO, Lamiaceae EO, Lauraceae EO) (MedP diluted at 10<sup>-2</sup>, EO diluted at 10<sup>-4</sup>-10<sup>-6</sup>).




**Fig.1** DIRECT ANTI-PROTOZOAL ACTIVITY AGAINST *E. TENELLA* SPOOROZOITES (\*: p<0.05; \*\*: p<0.01; \*\*\*: p<0.001)

MedP samples showed direct anti-protozoal activity against *E. tenella* stages: sporozoite viability rates were between 76.2% and 27% (p=0.05). Lilaceae EO and Lauraceae EO exhibited 81% and 77% (p=0.05) sporozoite viability rates, respectively (Fig. 1).



**Fig.2** MDBK CELL INVASION BY *E. TENELLA* (\*: p<0.05; \*\*: p<0.01; \*\*\*: p<0.001)

Punicaceae extract showed a reduction of 32% of cell invasion.



**Fig.3** SPORULATION INHIBITION OF *E. TENELLA* (\*: p<0.05; \*\*: p<0.01; \*\*\*: p<0.001)

Acanthaceae extract and Urticaceae extract showed a sporulation inhibition of *E. tenella* oocysts (p=0.05) (Fig. 3). These medicinal plant extracts exhibited anti-protozoal activity against the different *E. tenella* stages contrary to the tested EOs which target only sporozoites. These results were in accordance with studies reporting several modes of action of medicinal plant extracts, both direct anti-protozoal activity and protective cell activity [1].

## CONCLUSION

Medicinal plant extracts exhibited an anti-protozoal activity against the different *E. tenella* stages. EOs only target sporozoites in this study. These findings help to understand how plant bio-actives could act to reduce *E. tenella* pathogenicity. Further studies could be carried out to test *in vitro* and/or *in vivo* potential synergies between different natural compounds.

## REFERENCE

<sup>1</sup> WUNDERLICH et al. (2014), « Towards Identifying Novel Anti-Eimeria Agents: Trace Elements, Vitamins, and Plant-Based Natural Products ». Parasitology Research 113, no 10: 3547-56