



HAL
open science

Genomic prediction for grain yield and micro-environmental sensitivity in winter wheat

Miguel Raffo, Beatriz C. D. Cuyabano, Renaud Rincent, Pernille Sarup, Laurence Moreau, Tristan Mary-Huard, Just Jensen

► **To cite this version:**

Miguel Raffo, Beatriz C. D. Cuyabano, Renaud Rincent, Pernille Sarup, Laurence Moreau, et al.. Genomic prediction for grain yield and micro-environmental sensitivity in winter wheat. *Frontiers in Plant Science*, 2023, 13, 10.3389/fpls.2022.1075077 . hal-04127792

HAL Id: hal-04127792

<https://hal.inrae.fr/hal-04127792>

Submitted on 13 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



OPEN ACCESS

EDITED BY

Yusheng Zhao,
Leibniz Institute of Plant Genetics
and Crop Plant Research (IPK), Germany

REVIEWED BY

Lars Rönnegård,
Dalarna University, Sweden
Marco Lopez-Cruz,
Michigan State University, United States

*CORRESPONDENCE

Miguel A. Raffo
✉ mraffo@qgg.au.dk

SPECIALTY SECTION

This article was submitted to
Plant Breeding,
a section of the journal
Frontiers in Plant Science

RECEIVED 20 October 2022

ACCEPTED 23 December 2022

PUBLISHED 01 February 2023

CITATION

Raffo MA, Cuyabano BCD, Rincenc R,
Sarup P, Moreau L, Mary-Huard T and
Jensen J (2023) Genomic prediction for
grain yield and micro-environmental
sensitivity in winter wheat.
Front. Plant Sci. 13:1075077.
doi: 10.3389/fpls.2022.1075077

COPYRIGHT

© 2023 Raffo, Cuyabano, Rincenc, Sarup,
Moreau, Mary-Huard and Jensen. This is an
open-access article distributed under the
terms of the [Creative Commons Attribution
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that
the original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution or
reproduction is permitted which does not
comply with these terms.

Genomic prediction for grain yield and micro-environmental sensitivity in winter wheat

Miguel A. Raffo^{1*}, Beatriz C. D. Cuyabano², Renaud Rincenc³,
Pernille Sarup⁴, Laurence Moreau³, Tristan Mary-Huard^{3,5}
and Just Jensen¹

¹Center for Quantitative Genetics and Genomics, Aarhus University, Aarhus, Denmark, ²Université Paris Saclay, INRAE, AgroParisTech, GABI, Domaine de Vilvert, Jouy-en-Josas, France, ³Génétique Quantitative et Evolution – Le Moulon, INRAE, CNRS, AgroParisTech, Université Paris-Saclay, Gif-sur-Yvette, France, ⁴Nordic Seed A/S, Odder, Denmark, ⁵Université Paris-Saclay, AgroParisTech, INRAE, UMR MIA-Paris Saclay, Palaiseau, France

Individuals within a common environment experience variations due to unique and non-identifiable micro-environmental factors. Genetic sensitivity to micro-environmental variation (i.e. micro-environmental sensitivity) can be identified in residuals, and genotypes with lower micro-environmental sensitivity can show greater resilience towards environmental perturbations. Micro-environmental sensitivity has been studied in animals; however, research on this topic is limited in plants and lacking in wheat. In this article, we aimed to (i) quantify the influence of genetic variation on residual dispersion and the genetic correlation between genetic effects on (expressed) phenotypes and residual dispersion for wheat grain yield using a double hierarchical generalized linear model (DHGLM); and (ii) evaluate the predictive performance of the proposed DHGLM for prediction of additive genetic effects on (expressed) phenotypes and its residual dispersion. Analyses were based on 2,456 advanced breeding lines tested in replicated trials within and across different environments in Denmark and genotyped with a 15K SNP-Illumina-BeadChip. We found that micro-environmental sensitivity for grain yield is heritable, and there is potential for its reduction. The genetic correlation between additive effects on (expressed) phenotypes and dispersion was investigated, and we observed an intermediate correlation. From these results, we concluded that breeding for reduced micro-environmental sensitivity is possible and can be included within breeding objectives without compromising selection for increased yield. The predictive ability and variance inflation for predictions of the DHGLM and a linear mixed model allowing heteroscedasticity of residual variance in different environments (LMM-HET) were evaluated using leave-one-line-out cross-validation. The LMM-HET and DHGLM showed good and similar performance for predicting additive effects on (expressed) phenotypes. In addition, the accuracy of predicting genetic effects on residual dispersion was sufficient to allow genetic selection for resilience. Such findings suggests that DHGLM may be a good choice to increase grain yield and reduce its micro-environmental sensitivity.

KEYWORDS

micro-environmental sensitivity, climatic resilience, genetic heterogeneity of residual variance, genomic selection, wheat

Introduction

Developing cultivars well adapted to the different environmental conditions where production is performed is one of the main goals of breeding programs. The performance of cultivars in different environments is affected by genotype-by-environment interactions (G×E). The G×E can result from environmental variation either due to differences in years and localities termed macro-environmental sensitivity, or from variability due to environmental disturbances within environments termed micro-environmental sensitivity. Selecting genotypes with lower macro and micro-environmental sensitivity is beneficial as they can present greater buffer capacity against external disturbances, thus contributing to a higher and more stable yield in a target population of environments. Given the global context of climate change and the expansion of agriculture to new areas, the development of climate-resilient varieties is also relevant to safeguarding the sustainability of wheat production (Tilman et al., 2011; Ray et al., 2013).

Micro-environmental sensitivity is defined as the genetic sensitivity to those environmental features that are specific to each individual (i.e. micro-environmental variation) and that cannot be identified or measured at each particular plot (Falconer and Mackay, 1996; Hill and Mulder, 2010; Walsh and Lynch, 2018). The micro-environmental sensitivity has been called in literature (confusingly) non-genetic variance, stochastic variation or general environmental variance, among other denominations (Falconer and Mackay, 1996; Dworkin, 2005; Morgante et al., 2015); nevertheless, several studies show that the micro-environmental sensitivity is partly under genetic control (Sorensen and Waagepetersen, 2003; Mulder et al., 2007; Ordas et al., 2008; Mulder et al., 2009; Vandenplas et al., 2013; Jung et al., 2017). The study of micro-environmental sensitivity has gained relevance in recent years for several animal species such as chickens (Wolc et al., 2009), pigs (Sorensen and Waagepetersen, 2003; Sell-Kubiak et al., 2015; Sell-Kubiak et al., 2022), sheep (SanCristobal-Gaudy et al., 1998; SanCristobal-Gaudy et al., 2001), cattle (Mulder et al., 2013a; Vandenplas et al., 2013), and aquaculture species (Sae-Lim et al., 2016). However, to the best of our knowledge, studies on micro-environmental sensitivity have barely been conducted in plants (e.g. Ordas et al., 2008 for maize), and have not been developed on commercial wheat varieties.

Before describing methods to estimate genetic variation in residual dispersion, it is useful to distinguish between biological (also seen as quantitative, Hill and Mulder, 2010) and statistical models used to infer genetic parameters. The 'biological' denotation refers to the theoretical genetic model specifying the genetic effects underlying the phenotype, while the 'statistical' sense refers to the model used to estimate genetic parameters related to these effects and is useful for predicting genetic values and estimating the response of selection (Falconer and Mackay, 1996). Several quantitative models explaining the influence of genetic effect on micro-environmental variance have been proposed in the literature (reviewed by Hill and Mulder, 2010); for example, the standard deviation model (Garcia et al., 2009), the additive model (Hill and Zhang, 2004; Mulder et al., 2007), and the exponential model (SanCristobal-Gaudy et al., 1998; SanCristobal-Gaudy et al., 2001; Sorensen and Waagepetersen, 2003). The exponential model, further described in the next section, has been

proposed as a good alternative since the genetic values in micro-environmental variance are additive on a log scale, and it avoids the problem of having negative variances (Hill and Mulder, 2010; Walsh and Lynch, 2018).

Studying the genetic component of micro-environmental sensitivity is challenging as it implies modelling genetic effects at the level of residual dispersion (as a new trait). Different statistical methods to estimate variance components (VCs) and breeding values in micro-environmental variance have been proposed in animal breeding (SanCristobal-Gaudy et al., 1998; Sorensen and Waagepetersen, 2003; Rönnegård et al., 2010; Felleki et al., 2012; Jung et al., 2017). As a general distinction, those methods can be classified into a frequentist (likelihood-based) or a Bayesian framework. Sorensen and Waagepetersen (2003) applied a Bayesian methodology to estimate the genetic effects on the (expressed) phenotype and residual dispersion, as well as the genetic correlation between them. Among the main likelihood-based methods, Rönnegård et al. (2010) used a double hierarchical generalized linear model (DHGLM) developed by Lee and Nelder (2006). This methodology fits an algorithm that iterates between a linear mixed model at the level of (expressed) phenotypes and a generalized linear model (GLM) with gamma-distributed residuals at the level of residual dispersion. Felleki et al. (2012) extended the DHGLM model proposed by Rönnegård et al. (2010) to include the correlation between random genetic effects for (expressed) phenotypes and residual dispersion. The models proposed by Rönnegård et al. (2010) and Felleki et al. (2012) have been successfully applied since then and have been further extended to include genomic information and macro-environmental sensitivity, among other applications (Mulder et al., 2013a; Mulder et al., 2013b; Jung et al., 2017; Ehsaninia et al., 2019; Madsen et al., 2021; Sell-Kubiak et al., 2022). In addition, the DHGLM methodology has proven to be a computationally efficient approach to estimating micro-environmental sensitivity (Rönnegård et al., 2010; Felleki et al., 2012). Despite the availability of methodologies, common linear mixed models used for genomic prediction (GP) in plant breeding assume homogeneous residual variance within environments and do not account for micro-environmental sensitivity. Hence, more sophisticated methods are required to study micro-environmental sensitivity in plant breeding.

Several genetic parameters can be estimated to understand the implications of micro-environmental sensitivity in breeding; among the most relevant are the Mulder-Hill heritability of residual variance (h_d^2), the genomic coefficient of variation of residual variance or evolvability (GCV_E), and the correlation between additive genetic effects in (expressed) phenotypes and residual dispersion (ρ_{g,g_d}). The h_d^2 is analogous to the narrow sense heritability in the classical sense (h^2 , Falconer and Mackay, 1996), and it is defined as the slope of the regression of the estimated additive values for dispersion on the squared phenotypes (P^2 , Mulder et al., 2007). The GCV_E indicates the proportion of micro-environmental variance that can be changed by selection, and the ρ_{g,g_d} allows to infer how the selection on additive genetic effects on (expressed) phenotypes will affect the additive genetic effects on dispersion. Estimates for these parameters can be obtained from the output of the DHGLM model and are useful references to develop more resilient cultivars exhibiting lower

micro-environmental sensitivity. In addition, the h_a^2 and GCV_E can be useful for comparison across traits and species.

Plant species where inbred lines, single-crosses between inbred lines, or clones can be developed may present a high potential for studying micro-environmental sensitivity. That occurs because the availability of replications on the same individual contributes to a more accurate estimation of genetic variation and breeding values in micro-environmental variance (Vandenplas et al., 2013; Jung et al., 2017; Madsen et al., 2021). In this sense, studies on micro-environmental sensitivity in wheat can represent an opportunity, as highly homozygous inbred lines can be developed (e.g. six generations of selfing, F_6), so that individuals originating from the same line can be considered genetically homogeneous. Another reason that makes wheat a valuable species for this study is that there has been broad literature reporting a considerable variation due to macro-environmental sensitivity (Bhatt and Derera, 1975; Cooper et al., 1995; Sial et al., 2000; Roozeboom et al., 2008; Lopez-Cruz et al., 2015; Crossa et al., 2017; Ly et al., 2017; Sukumaran et al., 2017; Ly et al., 2018; Crossa et al., 2021; Raffo et al., 2022a), and therefore, it encourages hypothesizing on a relevant genetic variation for micro-environmental sensitivity. In this study, we used a winter wheat breeding population phenotyped for grain yield, and we had two specific objectives:

1. To quantify the genetic variation on residual dispersion and, if it exists, the genetic correlation between genetic effects on (expressed) phenotype and residual dispersion for wheat grain yield using a double hierarchical generalized linear model (DHGLM).
2. To evaluate the predictive performance of the proposed double hierarchical generalized linear model for prediction of additive genetic effects on (expressed) phenotype and residual dispersion using cross-validation (CV) analysis.

To simplify reading, in the next sections we will refer to (expressed) phenotype as EGY (i.e. expressed grain yield) and to residual dispersion as DGY (i.e. residual dispersion of grain yield).

Materials and methods

Plant material

The data consisted of 2,456 sixth-generation (F_6) winter wheat lines (*T. aestivum* L.) and 21,894 plot observations developed by Nordic Seed A/S breeding company. The breeding lines originated from nine breeding cycles (BC) tested in the years 2013 to 2021 (cycle 1: 2013–2014, cycle 2: 2014–2015, cycle 3: 2015–2016, cycle 4: 2016–2017, cycle 5: 2017, cycle 6: 2018, cycle 7: 2019, cycle 8: 2020, cycle 9: 2021) in three locations in Denmark (DK): Skive (north-west DK), Odder (central DK), and Holeby (south DK). In total, 26-year-location subsets were successfully assessed, and one failed (Odder, 2018) due to operational issues. The number of lines, plot observations, and average line replications per BC and for the whole population are presented together with descriptive statistics for grain yield in Table 1. Each BC came from approximately 60 parental line-crosses followed by selfing until F_6 , including family-based selection until third-generation (F_3) and creating single seed descent (SSD) lines in the fourth-generation (F_4). The experimental trials for each year-location combination comprised 15 blocks of 46 sowing plots of 8.25 m², containing two replications of 21 F_6 lines and two checks assigned randomly within each block. All lines were phenotyped for grain yield measured as kg per plot (8.25 m²). Agronomic practices were standardized within and across year-location subsets (e.g. application of disease treatments, fertilization).

Genotyping

DNA were extracted based on a modified CTAB method (Rogers and Bendich, 1985). A 15K Illumina Infinium iSelect HD Custom Genotyping BeadChip technology (Wang et al., 2014) was used for genotyping. Quality control was performed by removing genotyped SNPs with a call rate lower than 0.90 and minor allele frequency (MAF) lower than 5%. Missing genotypes were imputed with the mean value after centering the genomic matrix (~3% of missing

TABLE 1 Descriptive statistics for the grain yield (kg grain/8.25m²) of F_6 wheat breeding lines.

Breeding Cycle	Year	No. of Lines	No. of Plots	Average no. of line reps.	Average yield (SD)	Min. - Max Values	Coefficient of variation (%)
1	2013	78	1,060	13.59	8.89 (0.60)	6.48 – 10.45	6.70
2	2014	324	3,512	10.84	8.70 (0.76)	3.84 – 11.05	8.84
3	2015	232	2,574	11.09	8.64 (1.13)	4.75 – 11.47	13.11
4	2016	338	3,598	10.64	8.31 (1.37)	5.03 – 11.80	13.69
5	2017	160	1,012	6.32	9.13 (0.96)	6.21 – 11.40	10.58
6	2018	359	1,721	4.79	8.59 (0.46)	7.06 – 10.25	5.33
7	2019	315	2,589	8.22	9.38 (1.07)	6.04 – 12.36	11.38
8	2020	354	3,040	8.59	9.37 (0.68)	6.23 – 11.54	7.27
9	2021	305	2,788	9.14	9.02 (0.68)	6.45 – 11.32	7.55
Total	–	2,456	21,894	8.91	9.15 (0.82)	6.04 – 12.36	9.01

No., number; Reps., replications; SD, standard deviation; Min, Minimum; Max, Maximum.

values imputed). A total of 12,893 SNPs remained after the quality control.

Corrected phenotypes

As a first step, a linear mixed model including all relevant effects for the population was utilized to analyze raw data. Such a model has been previously used in Raffo et al. (2022b) with a similar dataset from the same breeding program, and in brief, it was defined as: $y = Xb + Z_1l + Z_2g + Z_3f + \sum_{i=1}^9 Z_{i+3}s + e$, where Xb , Z_1l , Z_2g and Z_3f were defined as below in the LMM-HET (eq. 1), $\sum_{i=1}^9 Z_{i+3}s$ is the spatial effect where s is the vector of spatial effect with $s \sim N(0, I\sigma_s^2)$ and σ_s^2 is the spatial effect variance, and e is a vector of random residuals with $e \sim N(0, I\sigma_e^2)$. The spatial effect contains the X and Y coordinate of the plot of the observation (i.e., target plot) and the eight surrounding plots ($n=9$), and thus, it is the sum of the effects centered on those plots. A detailed description of the model and the spatial effect is given in Raffo et al. (2022b) under the heading “I +G_A-model”. Second, the raw phenotypes were corrected by subtracting the estimates of the spatial effect (s) in order to correct for spatial variability within the experimental fields. The corrected phenotypes (y_c) were defined as: $y_c = y - Zs$, where y represents the vector of raw phenotypes, and $Z = \sum_{i=1}^9 Z_{i+3}$. This allowed us to reduce the number of parameters to estimate in the DHGLM, facilitating convergence and decreasing computing time.

Statistical analysis

Linear mixed model (LMM-HET)

A linear mixed model allowing for heteroscedasticity of residual variance for the different environments (hereinafter LMM-HET) was utilized in order to have a reference point for assessing the DHGLM. The LMM-HET was defined as:

$$y_c = Xb + Z_1g + Z_2l + Z_3f + e_i \tag{1}$$

where y_c is a vector for the response variable EGY (i.e. corrected phenotypes) with 21,894 observations from the 2,456 lines, X and Z_n are the design matrices for fixed and random effects, respectively; b is a vector of fixed trial effect nested within year-location; g is a vector of additive genetic values with $g \sim N(0, G\sigma_g^2)$, σ_g^2 is the genomic additive variance, and G is the additive genomic relationship matrix (GRM) according to the first method of VanRaden (2008): $G = \frac{QQ'}{2\sum p_j(1-p_j)}$, where p_j is the allele frequency of the j^{th} SNP; Q is $M-P$, with M as the SNP matrix coded -1, 0, 1, and P the matrix with the allele frequency of SNP j calculated as $1(2(p_j-0.5))$ for column j ; l is a vector of line effect with $l \sim N(0, I\sigma_l^2)$, where I is an identity matrix and σ_l^2 is the variance due to uncorrelated line effects, which are common to all replicates of a line; f is a vector of line \times environment interaction effect (L \times E), with “E” defined for the different year-location subsets and $f \sim N(0, I\sigma_f^2)$, where σ_f^2 is the variance due to uncorrelated L \times E effects; and e_i is a vector of

random residuals with $e \sim NIID(0, I\sigma_{e_i}^2)$, where $\sigma_{e_i}^2$ is the residual heterogeneous variance fitted for the different year-location subsets ($i=1, 2, \dots, 26$).

The narrow (h^2) and broad-sense (H^2) heritabilities (Falconer and Mackay, 1996) at the plot level were estimated for LMM-HET for each year-location group (i) as: $h_i^2 = d(G)\hat{\sigma}_g^2 / \hat{\sigma}_{p_i}^2$, and $H_i^2 = (\hat{\sigma}_l^2 + d(G)\hat{\sigma}_g^2) / \hat{\sigma}_{p_i}^2$, where $d(G)$ is the mean diagonal value of G with $d(G)=1.84$, which can be interpreted as one plus the genomic inbreeding coefficient for the population (VanRaden, 2008); $\hat{\sigma}_g^2$ is the estimated genomic additive variance; $\hat{\sigma}_l^2$ is the estimated variance due to uncorrelated line effects; and $\hat{\sigma}_{p_i}^2$ is the estimated phenotypic plot variance for each year-location subset (subscript “ i ”) calculated as: $\hat{\sigma}_{p_i}^2 = \hat{\sigma}_l^2 + d(G)\hat{\sigma}_g^2 + \hat{\sigma}_f^2 + \hat{\sigma}_{e_i}^2$.

Double hierarchical generalized linear model

First, we start by briefly describing the exponential model (SanCristobal-Gaudy et al., 1998), which is the base quantitative model assumed in our study. The exponential model is an extension of the classical model ($P=\mu+A+E$), and it was defined as: $P = \mu + A_m + L + F + \exp[0.5 \ln(\sigma_{E,exp}^2) + 0.5A_{v,exp}] \epsilon$, where P is the phenotype, μ is the intercept, A_m is the breeding value for the main effect, L is the polygenic genetic effect, F is the genetic \times macro-environment interaction effect, \exp specifies that the exponential function is used at the level of environmental variance, $\ln(\sigma_{E,exp}^2)$ is the natural logarithm of the environmental variance, $A_{v,exp}$ is the breeding value for the environmental variance, and ϵ is a scaled environmental deviation with variance one. As previously commented, in the exponential model the genetic values in environmental variance are additive on a log scale, and thus, it avoids the problem of having negative variances (Hill and Mulder, 2010; Walsh and Lynch, 2018).

The estimation of VCs and breeding values was performed running an iterative Average Information Restricted Maximum Likelihood (AI-REML) algorithm in DMU software (Madsen and Jensen, 2013) by the implementation of a DHGLM procedure (Lee and Nelder, 2006; Rönnegård et al., 2010; Felleki et al., 2012; Mulder et al., 2013a). The DHGLM was used to estimate genomic breeding values and genetic parameters for EGY and DGY. The DHGLM iterates using bivariate analysis (Equation 2), fitting a linear mixed model at the level of corrected phenotypes (y_c), and a Gamma GLM at the level of the dispersion variable y_d (see Supplementary Material 1 for an additional formal definition of the DHGLM), where $y_{d_n} = \frac{\hat{e}_n^2}{1-\hat{A}_n}$, with \hat{e}_n^2 as the squared estimated residual for the y_{c_n} observation, and \hat{A}_n is the ‘leverage’ defined as the diagonal element of the hat-matrix H (Hoaglin and Welsh, 1978) corresponding to y_{c_n} . The hat-matrix H is also known as the projection matrix since it provides \hat{y}_c from y_c as $\hat{y}_c = H^* y_c$. The DHGLM is an algorithm fitting the exponential model and is given by Equation 2:

$$\begin{bmatrix} y_c \\ y_d \end{bmatrix} = \begin{bmatrix} X & 0 \\ 0 & X_d \end{bmatrix} \begin{bmatrix} b \\ b_d \end{bmatrix} + \begin{bmatrix} Z_1 & 0 \\ 0 & Z_2 \end{bmatrix} \begin{bmatrix} g \\ g_d \end{bmatrix} + \begin{bmatrix} Z_3 \\ 0 \end{bmatrix} [I] + \begin{bmatrix} Z_4 \\ 0 \end{bmatrix} [f] + \begin{bmatrix} e \\ e_d \end{bmatrix} \tag{2}$$

where y_c and y_d are the vectors of response variables for EGY and DGY, respectively; X , b , Z_n , l , and f were defined as in LMM-HET.

Note that \mathbf{l} and \mathbf{f} were included only for EGY since it was not possible for DGY due to problems of convergence in the REML algorithm. The convergence problems may be caused by a random effect with a variance close to 0, by between-trait correlations of -1 or 1, or by two different parameters in the model that are difficult to separate (i.e. high correlation between random effects for the same trait). \mathbf{X}_d is the design matrix for the fixed effects in DGY, \mathbf{b}_d is the vector of fixed trial effect nested within year-location in DGY; \mathbf{g} and \mathbf{g}_d are the vectors of additive genetic values for EGY and DGY, respectively, following the distribution

$$\begin{bmatrix} \mathbf{g} \\ \mathbf{g}_d \end{bmatrix} \sim N\left(0, \begin{bmatrix} \sigma_g^2 & \sigma_{g,g_d} \\ \sigma_{g,g_d} & \sigma_{g_d}^2 \end{bmatrix} \otimes \mathbf{G}\right)$$

where σ_g^2 and $\sigma_{g_d}^2$ are the genomic additive variance for EGY and DGY, respectively, and σ_{g,g_d} is the covariance between genetics effect in EGY and DGY, \otimes denotes the Kronecker product, and \mathbf{G} the GRM as previously defined for the LMM-HET model; \mathbf{e} and \mathbf{e}_d the vectors of random residual effects for EGY and DGY, respectively, following the distribution

$$\begin{bmatrix} \mathbf{e} \\ \mathbf{e}_d \end{bmatrix} \sim N\left(0, \begin{bmatrix} \mathbf{W}^{-1}\sigma_e^2 & \mathbf{0} \\ \mathbf{0} & \mathbf{W}_d^{-1}\sigma_{e_v}^2 \end{bmatrix}\right)$$

where $\mathbf{W} = \text{diag}(\hat{y}_d)^{-1}$ and $\mathbf{W}_d = \text{diag}(\frac{1-\Lambda}{2})$ with Λ being the leverage (diagonal element of the hat-matrix \mathbf{H}), and σ_e^2 and $\sigma_{e_v}^2$ the scaling for residual variances constrained to have an average of one. Given that results on each trait of the bivariate model depend on each other, an iterative procedure was required to update \mathbf{W} and \mathbf{W}_d until convergence of variance component estimates. Convergence was assumed when the relative difference in variance component estimates of two successive iterations was lower than 10^{-5} . The bivariate model in Equation 2 was fitted using a reweighted least square (IRWLS) algorithm as follows:

1. Run a linear mixed model (LMM) with the same effects as in eq. 1 but assuming homoscedasticity of residual variance on \mathbf{y}_c .
2. Calculate $\mathbf{y}_{d_n} = \hat{e}_n^2 / (1 - \Lambda_n)$, and $\mathbf{W}_{d_n} = \text{diag}(1 - \Lambda_n) / 2$
3. Run a weighted gamma generalized linear model (GLM) with a log-link function for response \mathbf{y}_d and weight \mathbf{W}_d
4. Calculate $\mathbf{W} = \text{diag}(\hat{y}_d)^{-1}$
5. Run bivariate model in Equation 2
6. Update \mathbf{y}_d , \mathbf{W} , and \mathbf{W}_d
7. Iterates steps 3-6 until convergence

The model in Equation 2 and the iterative estimation procedure were developed by Felleki et al. (2012) and are based on a combination of the DHGLM proposed by Rönnegård et al. (2010) and the method proposed by Mulder et al. (2009). Note that convergence in step 5 is required to reach global convergence of the whole algorithm. The h_i^2 , H_i^2 , and $\sigma_{e_i}^2$ for EGY were estimated for the DHGLM using the equations described in the “Linear mixed model (LMM-HET)” section, but using an approximation for $\sigma_{e_i}^2$ computed as $\hat{\sigma}_{e_i}^2 = \frac{\sum_{j=1}^n (x_{ij} - \bar{x}_i)^2}{n_i - 1 - npar_i}$, where x_{ij} represents the residual for the j

observation in the environment i , \bar{x}_i is the mean value of residuals in the environment i , n_i is the number of observations in the environment i , and $npar_i$ is the number of fixed effects estimated in the model for the environment i . Thus, the residual variance approximation is based on the variance of the final vector of residuals.

Genetic parameters associated with micro-environmental sensitivity

Three genetic parameters were computed in order to interpret the micro-environmental sensitivity in our population: i) Mulder-Hill heritability of residual variance (h_d^2), ii) genomic coefficient of variation of residual variance or evolvability (GCV_E), and iii) genetic correlation between additive effects on mean and dispersion (ρ_{g,g_d}). A description of the estimated parameters is provided next.

The h_d^2 can be defined (by analogy to h^2) as the regression of additive genetic values in dispersion on the squared phenotypes (p^2).

The h_d^2 was estimated as defined by Mulder et al. (2007): $h_d^2 =$

$$\frac{\hat{\sigma}_{g_d,add}^2}{2\hat{\sigma}_p^4 + 3\hat{\sigma}_{g_d,add}^2},$$

where $\hat{\sigma}_{g_d,add}^2$ is the rescaled genomic additive variance for DGY on the additive scale, and $\hat{\sigma}_p^4$ is the squared phenotypic variance. Note that for the estimation of h_d^2 , the additive genetic variance estimated for DGY required to be converted from the exponential scale ($\hat{\sigma}_{g_d}^2$) to the additive scale ($\hat{\sigma}_{g_d,add}^2$). This conversion was performed using equations in Mulder et al. (2007):

$$\hat{\sigma}_{g_d,add}^2 = \hat{\sigma}_{e,exp}^4 \exp(\hat{\sigma}_{g_d}^2) - \hat{\sigma}_{e,add}^2,$$

where $\hat{\sigma}_{e,exp}^2 = \frac{[(1/\mathbf{W})\hat{\sigma}_e^2]}{[\exp(0.5\hat{\sigma}_{g_d}^2)]}$, $\hat{\sigma}_{g_d}^2$ is the additive genetic variance estimated for the residual variable on the exponential scale, $\hat{\sigma}_{g_d,add}^2 = [(1/\mathbf{W})\hat{\sigma}_e^2]$, with $[(1/\mathbf{W})\hat{\sigma}_e^2]$ and $\hat{\sigma}_e^2$ the average of the reciprocal weights and the residual variance in the mean model, respectively (see Mulder et al., 2007 for complete derivations). The genetic coefficient of variation (GCV_E) allows to infer how much the micro-environmental variance could be changed by selection (Mulder et al., 2016), and it was computed as $GCV_E =$

$$\sqrt{\hat{\sigma}_{g_d}^2}.$$

The genetic correlation between additive effects on EGY and DGY (ρ_{g,g_d}) was estimated as: $\rho_{g,g_d} = \hat{\sigma}_{g,g_d} / \sqrt{\hat{\sigma}_g^2 \hat{\sigma}_{g_d}^2}$, with $\hat{\sigma}_{g,g_d}$, $\hat{\sigma}_g^2$, and $\hat{\sigma}_{g_d}^2$ as previously defined for the DHGLM.

The genetic correlation between additive effects on EGY and DGY (ρ_{g,g_d}) was estimated as: $\rho_{g,g_d} = \hat{\sigma}_{g,g_d} / \sqrt{\hat{\sigma}_g^2 \hat{\sigma}_{g_d}^2}$, with $\hat{\sigma}_{g,g_d}$, $\hat{\sigma}_g^2$, and $\hat{\sigma}_{g_d}^2$ as previously defined for the DHGLM.

The genetic correlation between additive effects on EGY and DGY (ρ_{g,g_d}) was estimated as: $\rho_{g,g_d} = \hat{\sigma}_{g,g_d} / \sqrt{\hat{\sigma}_g^2 \hat{\sigma}_{g_d}^2}$, with $\hat{\sigma}_{g,g_d}$, $\hat{\sigma}_g^2$, and $\hat{\sigma}_{g_d}^2$ as previously defined for the DHGLM.

The genetic correlation between additive effects on EGY and DGY (ρ_{g,g_d}) was estimated as: $\rho_{g,g_d} = \hat{\sigma}_{g,g_d} / \sqrt{\hat{\sigma}_g^2 \hat{\sigma}_{g_d}^2}$, with $\hat{\sigma}_{g,g_d}$, $\hat{\sigma}_g^2$, and $\hat{\sigma}_{g_d}^2$ as previously defined for the DHGLM.

The genetic correlation between additive effects on EGY and DGY (ρ_{g,g_d}) was estimated as: $\rho_{g,g_d} = \hat{\sigma}_{g,g_d} / \sqrt{\hat{\sigma}_g^2 \hat{\sigma}_{g_d}^2}$, with $\hat{\sigma}_{g,g_d}$, $\hat{\sigma}_g^2$, and $\hat{\sigma}_{g_d}^2$ as previously defined for the DHGLM.

The genetic correlation between additive effects on EGY and DGY (ρ_{g,g_d}) was estimated as: $\rho_{g,g_d} = \hat{\sigma}_{g,g_d} / \sqrt{\hat{\sigma}_g^2 \hat{\sigma}_{g_d}^2}$, with $\hat{\sigma}_{g,g_d}$, $\hat{\sigma}_g^2$, and $\hat{\sigma}_{g_d}^2$ as previously defined for the DHGLM.

The genetic correlation between additive effects on EGY and DGY (ρ_{g,g_d}) was estimated as: $\rho_{g,g_d} = \hat{\sigma}_{g,g_d} / \sqrt{\hat{\sigma}_g^2 \hat{\sigma}_{g_d}^2}$, with $\hat{\sigma}_{g,g_d}$, $\hat{\sigma}_g^2$, and $\hat{\sigma}_{g_d}^2$ as previously defined for the DHGLM.

The genetic correlation between additive effects on EGY and DGY (ρ_{g,g_d}) was estimated as: $\rho_{g,g_d} = \hat{\sigma}_{g,g_d} / \sqrt{\hat{\sigma}_g^2 \hat{\sigma}_{g_d}^2}$, with $\hat{\sigma}_{g,g_d}$, $\hat{\sigma}_g^2$, and $\hat{\sigma}_{g_d}^2$ as previously defined for the DHGLM.

The genetic correlation between additive effects on EGY and DGY (ρ_{g,g_d}) was estimated as: $\rho_{g,g_d} = \hat{\sigma}_{g,g_d} / \sqrt{\hat{\sigma}_g^2 \hat{\sigma}_{g_d}^2}$, with $\hat{\sigma}_{g,g_d}$, $\hat{\sigma}_g^2$, and $\hat{\sigma}_{g_d}^2$ as previously defined for the DHGLM.

The genetic correlation between additive effects on EGY and DGY (ρ_{g,g_d}) was estimated as: $\rho_{g,g_d} = \hat{\sigma}_{g,g_d} / \sqrt{\hat{\sigma}_g^2 \hat{\sigma}_{g_d}^2}$, with $\hat{\sigma}_{g,g_d}$, $\hat{\sigma}_g^2$, and $\hat{\sigma}_{g_d}^2$ as previously defined for the DHGLM.

The genetic correlation between additive effects on EGY and DGY (ρ_{g,g_d}) was estimated as: $\rho_{g,g_d} = \hat{\sigma}_{g,g_d} / \sqrt{\hat{\sigma}_g^2 \hat{\sigma}_{g_d}^2}$, with $\hat{\sigma}_{g,g_d}$, $\hat{\sigma}_g^2$, and $\hat{\sigma}_{g_d}^2$ as previously defined for the DHGLM.

The genetic correlation between additive effects on EGY and DGY (ρ_{g,g_d}) was estimated as: $\rho_{g,g_d} = \hat{\sigma}_{g,g_d} / \sqrt{\hat{\sigma}_g^2 \hat{\sigma}_{g_d}^2}$, with $\hat{\sigma}_{g,g_d}$, $\hat{\sigma}_g^2$, and $\hat{\sigma}_{g_d}^2$ as previously defined for the DHGLM.

The genetic correlation between additive effects on EGY and DGY (ρ_{g,g_d}) was estimated as: $\rho_{g,g_d} = \hat{\sigma}_{g,g_d} / \sqrt{\hat{\sigma}_g^2 \hat{\sigma}_{g_d}^2}$, with $\hat{\sigma}_{g,g_d}$, $\hat{\sigma}_g^2$, and $\hat{\sigma}_{g_d}^2$ as previously defined for the DHGLM.

The genetic correlation between additive effects on EGY and DGY (ρ_{g,g_d}) was estimated as: $\rho_{g,g_d} = \hat{\sigma}_{g,g_d} / \sqrt{\hat{\sigma}_g^2 \hat{\sigma}_{g_d}^2}$, with $\hat{\sigma}_{g,g_d}$, $\hat{\sigma}_g^2$, and $\hat{\sigma}_{g_d}^2$ as previously defined for the DHGLM.

The genetic correlation between additive effects on EGY and DGY (ρ_{g,g_d}) was estimated as: $\rho_{g,g_d} = \hat{\sigma}_{g,g_d} / \sqrt{\hat{\sigma}_g^2 \hat{\sigma}_{g_d}^2}$, with $\hat{\sigma}_{g,g_d}$, $\hat{\sigma}_g^2$, and $\hat{\sigma}_{g_d}^2$ as previously defined for the DHGLM.

The genetic correlation between additive effects on EGY and DGY (ρ_{g,g_d}) was estimated as: $\rho_{g,g_d} = \hat{\sigma}_{g,g_d} / \sqrt{\hat{\sigma}_g^2 \hat{\sigma}_{g_d}^2}$, with $\hat{\sigma}_{g,g_d}$, $\hat{\sigma}_g^2$, and $\hat{\sigma}_{g_d}^2$ as previously defined for the DHGLM.

and the vector of predicted additive genetic effects in EGY [$\rho(\bar{y}_e, \hat{g})$] and DGY [$\rho(\bar{y}_d, \hat{g}_d)$]. The \bar{y}_e and \bar{y}_d were obtained by subtracting the fixed effects \hat{b} and \hat{b}_d estimated with the full dataset from each corresponding plot observation for EGY and DGY, respectively, and then averaging the resulting values by line. ii) The correlation between predicted additive genetic values obtained with “whole” phenotypic information for all lines and with “partial” phenotypic information (predictions for all lines when their own phenotypes were masked in the CV) was computed for EGY and DGY ($r_{w,p}$, Legarra and Reverter, 2018). iii) A statistic for variance inflation in predicted genetic values ($b_{w,p}$) was estimated as the slope of the regression of predicted values obtained with whole phenotypic information on predicted values obtained with partial phenotypic information, $b_{w,p} = \frac{cov(\hat{g}_w, \hat{g}_p)}{var(\hat{g}_p)}$ (Legarra and Reverter, 2018). An ordinary non-parametric bootstrap with replacement based on full sample size $m = 2,456$, and 10,000 replicates was used to obtain the standard error of the PA, $r_{w,p}$, and $b_{w,p}$. A two-tailed paired t-test (critical P -value = 0.01) was used to compare PA, $r_{w,p}$ and $b_{w,p}$ for the LMM-HET and DHGLM. The maximum potential PA for g predictions was computed for each year-location subset as $\sqrt{kh_i^2/(1+(n-1)h_i^2)}$, where k was the average number of line repetitions and h_i^2 the heritability as previously defined, and then the average potential PA was calculated across the 26 year-location subsets. For g_d the maximum potential PA was computed using the same formula but replacing h_i^2 with h_d^2 (note that it is an approximation as the Gamma distribution is not taken into account).

Results

Phenotypic data

A total of 2,456 F_6 lines with an average number of replication of 8.91 were phenotyped for grain yield (kg grain/8.25m²), corresponding to 21,894 plot observations (see Table 1 for descriptive statistics per BC and for the whole population). Note that for the interpretation of results in tons of grain/ha instead of kg grain/8.25m², the presented values have to be multiplied by a conversion factor equal to 1.212. A considerable level of variation was observed for the different year-location subsets. The average grain yield varied from 6.90 to 10.45 kg grain/8.25m² and the coefficients of variation varied from 4.01 to 10.38% for the different year-location subsets. A boxplot describing grain yield per year-location subset is presented in Figure 1.

Variance components and heritability of additive genetic effect on mean

The proposed LMM-HET and DHGLM were utilized to estimate VCs on EGY and DGY, and the results are presented in Table 2.

A substantial amount of variance was observed in estimates for additive genetic variance ($\hat{\sigma}_g^2$), line effects ($\hat{\sigma}_l^2$) and line \times environment interaction ($\hat{\sigma}_{lj}^2$) for EGY with both models. The $\hat{\sigma}_g^2$ had the same value for both models ($\hat{\sigma}_g^2 = 0.054$). In general, small




FIGURE 1
Boxplot of grain yield (kg/8.25m²) per year-location subset. The x-axis labels “env#1” to “env#26” indicates the 26 year-location environments where lines were tested. The black dashed line represents the overall average grain yield for the complete population.

TABLE 2 Variance components and heritability estimates for the linear mixed model allowing heteroscedasticity of residual variance for the different environments (LMM-HET) and the double hierarchical generalized linear model (DHGLM).

Models	LMM-HET								DHGLM											Cor.
									Mean model						Dispersion					
	$\hat{\sigma}_g^2$	$\hat{\sigma}_l^2$	$\hat{\sigma}_l^2$	YL#	$\hat{\sigma}_{error}^2$	$\hat{\sigma}_P^2$	h^2	H^2	$\hat{\sigma}_g^2$	$\hat{\sigma}_l^2$	$\hat{\sigma}_l^2$	YL#	$\hat{\sigma}_{error}^2$	$\hat{\sigma}_P^2$	h^2	H^2		$\hat{\sigma}_{g_d}^2$	h_d^2	
Estimates	0.054 (0.008)*	0.060 (0.004)	0.118 (0.003)	env#1	0.129	0.361	0.150	0.315	0.054 (0.007)	0.062 (0.004)	0.108 (0.003)	env#1	0.120	0.344	0.156	0.337	0.004 (0.004)	0.033	0.502	
				env#2	0.080	0.311	0.173	0.365				env#2	0.105	0.329	0.163	0.352				
				env#3	0.038	0.269	0.201	0.423				env#3	0.114	0.338	0.158	0.343				
				env#4	0.079	0.310	0.174	0.366				env#4	0.110	0.334	0.160	0.347				
				env#5	0.114	0.346	0.156	0.329				env#5	0.105	0.329	0.163	0.352				
				env#6	0.068	0.299	0.181	0.380				env#6	0.110	0.335	0.160	0.346				
				env#7	0.106	0.337	0.160	0.337				env#7	0.072	0.297	0.181	0.391				
				env#8	0.348	0.579	0.093	0.196				env#8	0.083	0.307	0.174	0.377				
				env#9	0.054	0.285	0.189	0.399				env#9	0.104	0.329	0.163	0.353				
				env#10	0.036	0.267	0.202	0.425				env#10	0.124	0.348	0.154	0.333				
				env#11	0.099	0.330	0.164	0.345				env#11	0.107	0.331	0.162	0.350				
				env#12	0.072	0.304	0.178	0.374				env#12	0.102	0.327	0.164	0.355				
				env#13	0.064	0.296	0.183	0.385				env#13	0.132	0.356	0.150	0.325				
				env#14	0.077	0.308	0.175	0.369				env#14	0.104	0.328	0.163	0.353				
				env#15	0.067	0.298	0.181	0.381				env#15	0.107	0.331	0.162	0.350				
				env#16	0.063	0.294	0.183	0.386				env#16	0.050	0.274	0.195	0.422				
				env#17	0.078	0.309	0.175	0.368				env#17	0.048	0.272	0.197	0.426				
				env#18	0.064	0.296	0.183	0.385				env#18	0.054	0.278	0.193	0.417				
				env#19	0.088	0.319	0.169	0.357				env#19	0.057	0.281	0.190	0.412				
				env#20	0.081	0.312	0.173	0.365				env#20	0.053	0.277	0.193	0.418				
				env#21	0.046	0.278	0.195	0.410				env#21	0.055	0.279	0.192	0.416				
				env#22	0.335	0.566	0.095	0.201				env#22	0.053	0.277	0.193	0.418				
				env#23	0.048	0.280	0.193	0.407				env#23	0.050	0.274	0.195	0.423				
				env#24	0.090	0.321	0.168	0.354				env#24	0.066	0.290	0.185	0.399				
				env#25	0.087	0.318	0.170	0.357				env#25	0.070	0.294	0.182	0.394				
				env#26	0.082	0.313	0.173	0.363				env#26	0.063	0.287	0.187	0.404				
Average					0.096	0.327	0.171	0.359					0.085	0.310	0.175	0.377				
Min.					0.036	0.267	0.093	0.196					0.048	0.272	0.150	0.325				
Max.					0.348	0.579	0.202	0.425					0.132	0.356	0.197	0.426				

*Values in round brackets are the standard deviation (SD) of estimates; †for estimates $\hat{\sigma}_{error}^2$, $\hat{\sigma}_P^2$, h^2 , and H^2 values are presented for the 26 year-location (YL#) subsets (“env#1” to “env#26”); Cor.: genetic correlation; $\hat{\sigma}_{g_d}^2$ result is in exponential scale. Min., Minimum; Max., Maximum.

differences in estimates for $\hat{\sigma}_i^2$ and $\hat{\sigma}_f^2$ were observed between models, where the DHGLM had a slightly higher $\hat{\sigma}_i^2$ (0.062) compared to the LMM-HET (0.060), and lower $\hat{\sigma}_f^2$ (0.108) compared to the LMM-HET (0.118). The residual variance ($\hat{\sigma}_{e_i}^2$) was obtained for the 26 different year-location subsets with both models (see Table 2 for comparing specific year-location subsets). Generally, similar values for $\hat{\sigma}_{e_i}^2$ were observed between several of the 26-year-location subsets. The largest differences in $\hat{\sigma}_{e_i}^2$ were observed for environments #8 and #22 (env#8 and env#22 in Table 2), where the DHGLM had considerably lower $\hat{\sigma}_{e_i}^2$ (env#8: 0.083; env#22: 0.053) compared to the LMM-HET (env#8: 0.348 and env#22: 0.335).

The estimated phenotypic variance ($\hat{\sigma}_p^2$), narrow-sense (h_i^2) and broad-sense (H_i^2) heritabilities were obtained for the 26-year-location subsets with both models (Table 2). The $\hat{\sigma}_p^2$ ranged from 0.267 to 0.579 with an average value of 0.327 for the LMM-HET, and from 0.272 to 0.356 with an average of 0.310 for the DHGLM. The h_i^2 varied across year-locations subsets from 0.093 to 0.202 for the LMM-HET and from 0.150 to 0.197 for the DHGLM. The average h_i^2 were 0.175 for the DHGLM and 0.171 for the LMM-HET. The H_i^2 ranged across year-locations from 0.196 to 0.425 for the LMM-HET and from 0.325 to 0.377 for the DHGLM, and the average H_i^2 was 0.377 for the DHGLM and 0.359 for the LMM-HET. The highest estimates for heritabilities were obtained for environments #10 for the LMM-HET and environment #17 for the DHGLM, and the lowest were observed for environments #8 for the LMM-HET and #13 for the DHGLM.

The DHGLM was used for the estimation of fixed effects (\hat{b}_d) and additive genetic variance in DGY ($\hat{\sigma}_{g_d}^2$). The \hat{b}_d captured the variation due to trial effects nested within year-location at the level of DGY, and a substantial variation among \hat{b}_d was observed (Supplementary Material 2, Figure 1S). The $\hat{\sigma}_{g_d}^2$ had a value of 0.004 (estimated in exponential scale), and it is reported in Table 2. The interpretation of $\hat{\sigma}_{g_d}^2$, is linked to the estimation of genetic parameters associated with micro-environmental sensitivity, and thus it was later used for the estimation of the Mulder-Hill heritability of residual variance (h_d^2) and the genomic coefficient of variation of residual variance or evolvability (GCV_E , see next section: “Genetic parameters estimated for micro-environmental sensitivity”). The bivariate model in the DHGLM procedure allowed the estimation of the genetic correlation between additive genetic effects in EGY and DGY ($\rho_{g,g_d} = 0.502$). Further interpretation of ρ_{g,g_d} is presented in the following section. Differences in the predicted additive genetic values in DGY (\hat{g}_d) were reflected in differences of within line variability, where extreme cases for lines with lower and higher \hat{g}_d had low and high within line variation in DGY, respectively (Figure 2).

Genetic parameters estimated for micro-environmental sensitivity

The VCs presented in the previous section were utilized to estimate three genetic parameters associated with micro-environmental sensitivity (h_d^2 , GCV_E , and ρ_{g,g_d}):

- I. An estimate of Mulder-Hill heritability of residual variance (h_d^2 , Table 2) of 0.033 was found for grain yield in our population. Such a value is situated at an intermediate level according to the range previously reported for other species,

which varied from <0.01 to 0.10 (Wolc et al., 2009; Sonesson et al., 2013; Vandenplas et al., 2013; Sell-Kubiak et al., 2022).

- II. The genomic coefficient of variation of residual variance or evolvability (GCV_E) found for our population was 0.061. The GCV_E is useful to infer the selection response in reducing micro-environmental variance when lines with lower additive values in DGY are selected (Rönnegård et al., 2013; Lung et al., 2020). Therefore, a GCV_E of 0.061 indicates that reducing the average additive values in DGY in one unit of GCV_E will reduce the micro-environmental variance by 6.1%.
- III. The genetic correlation between additive effects in EGY and DGY (ρ_{g,g_d}) estimated for our population was 0.502 (Table 2). The positive correlation found suggests that selecting lines with higher additive values for \hat{g} will have an effect on increasing the additive values for \hat{g}_d and consequently will increase the micro-environmental sensitivity. This is confirmed by the strong linear relationship observed between \hat{g} and \hat{g}_d (Figure 3).

Comparative predictive performance of LMM-HET and DHGLM

The predictive ability (PA), calculated as the Pearson correlation (ρ) between the average value of lines after correcting for fixed effects and the vector of predictions of the genetic additive effects on EGY [$\rho(\bar{y}_c, \hat{g})$] and DGY [$\rho(\bar{y}_d, \hat{g}_d)$] obtained by LOO CV, is reported in Figure 4. The PA observed for predicted additive genetic values for EGY (predictions of g effect) in the DHGLM and the LMM-HET were 0.4292 and 0.4289, respectively, and no significant differences between models were observed in a bootstrap-based t-test (P -value > 0.01). The maximum potential PA for predictions of additive values for EGY (horizontal green lines in Figure 4) had similar values for both models, where the reported values were 0.807 for the DHGLM and 0.801 for the LMM-HET (average for the 26-year-location groups). The PA observed for predicted additive genetic values for DGY (predictions of g_d effect) in the DHGLM was 0.1003, with a maximum potential PA of 0.483. The PA for DGY was significantly lower in the t-test (P -value < 0.01) than the PAs reported for EGY with the DHGLM and the LMM-HET.

The correlation ($r_{w,p}$) between \hat{g} (and \hat{g}_d) obtained with “whole” phenotypic information for all lines and “partial” phenotypic information from the LOO CV analysis is presented in Figure 5. The $r_{w,p}$ reported for additive genetic values for EGY was significantly higher for the DHGLM (0.9517) than for the LMM-HET (0.9467) in the bootstrap-based t-test (P -value < 0.01). The $r_{w,p}$ estimated for the additive genetic values in DGY was 0.9049; this value was significantly lower (~4%), in the t-test (P -value < 0.01) than predictions for additive genetic values for EGY from both models.

The statistic for variance inflation of predicted additive genetic values ($b_{w,p}$) are presented in Figure 6. The expected $b_{w,p}$ when there is no variance inflation in predictions is 1 (black dashed horizontal line in Figure 6); lower and higher values than the unity represent over or under-dispersion, respectively. In our study, we did not observe variance inflation in predictions for additive genetic values in EGY for the DHGLM (0.9951) and the LMM-HET (0.9955), and no significant difference between models was found. Conversely, the




FIGURE 2

Illustration of variability between line replications in the dispersion variable (y_d) after algorithm convergence for a line with low additive genetic value (A, left plot) and a line with high additive genetic value (B, right plot).

predictions for additive genetic values in DGY for the DHGLM revealed a low over-dispersion with a $b_{w,p}$ of 0.9346 and a bootstrap-based confidence interval ranging from 0.9159 to 0.9532. The $b_{w,p}$ for predictions of the additive genetic effects in EGY and DGY were significantly different in the t-test (P -value < 0.01).

Discussion

In this study, we utilized a DHGLM to investigate the genetic variation in residual variance for grain yield (micro-environmental sensitivity) and to assess the predictability of additive genetic effects in EGY (y_e) and DGY (y_d) variables. As far as we know, studies on micro-environmental sensitivity have been limited in plants, and this is the first time that genetic parameters for micro-environmental sensitivity have been investigated for wheat grain yield. We found that the micro-environmental sensitivity for grain yield is heritable and that there is potential for reducing micro-environmental sensitivity




FIGURE 3

Regression of predicted additive genetic values for the dispersion variable (g_d) after algorithm convergence on predicted additive genetic values for corrected phenotypes (g). The blue line represents the fitted curve of the linear regression.

and increasing resilience of wheat breeding lines. In addition, we found that the DHGLM had similar performance for predictions of additive genetic effects in EGY compared to the LMM-HET, and had suitable performance for prediction of additive genetic effects in DGY.

Variance components and heritability estimates using LMM-HET and DHGLM

The LMM-HET and DHGLM revealed substantial variance for g , l , and f (Table 2). The $\hat{\sigma}_g^2$ had the same estimate for both models. Small differences between models were found for $\hat{\sigma}_l^2$ and $\hat{\sigma}_f^2$, where a slightly higher $\hat{\sigma}_l^2$ and lower $\hat{\sigma}_f^2$ was observed for the DHGLM. The $\hat{\sigma}_l^2$ is the variance due to common line effects (l), which specify the genetic effects that are not captured by markers in the genomic term, and hence mostly non-additive variance is expected in $\hat{\sigma}_l^2$. The residual variance ($\hat{\sigma}_{e_i}^2$) was obtained for each year-location subset with both models, and generally similar values were obtained for the different environments. The largest differences were observed for environments #2 and #8 (Table 2), where the DHGLM had considerably lower $\hat{\sigma}_{e_i}^2$. The lower residual variance is generally associated with a better fit of models, and in our case, it could be related to a more specific definition of heterogeneous residuals due to additive genetic effect in the DHGLM. The $\hat{\sigma}_p^2$, h_i^2 and H_i^2 estimated with the LMM-HET and the DHGLM were in general similar across environments, except for environments #2 and #8 where the DHGLM had a considerable higher estimate of h_i^2 and H_i^2 . Despite the differences across environments, the h_i^2 and H_i^2 were in the range of previous studies for grain yield using Nordic Seed A/S data (Guo et al., 2020; Raffo et al., 2022a; Raffo et al., 2022b).

An alternative DHGLM was in addition implemented for VCs estimation to account for G×E due to additive-by-environment interactions in EGY. In this alternative formulation, the G×E effect was modelled using a genomic relationship matrix to relate lines within each environment. However, the G×E due to additive genotype-by-environment interactions was excluded from the analysis due to

convergence problems in the REML algorithm. In this sense, the line \times environment interaction effect (L \times E) can be seen as a G \times E effect capturing a combination of additive and non-additive effects, and helps to specify the G \times E effects for our population.

The DHGLM was used to estimate fixed effects (\hat{b}_d) and additive genetic variance in dispersion ($\hat{\sigma}_{g_d}^2$). The \hat{b}_d revealed a considerable variation, which implies that the specification of fixed effects in DGY is justified in the developed model. The fixed effects estimate the effect of trials nested within year-location subsets; therefore, they can capture variation in DGY due to differences in trials within the experimental field and across year-location environments. The genetic variance in DGY was $\hat{\sigma}_{g_d}^2 = 0.004$. Several authors have reported that using replications of genetically similar individuals (e.g. inbred lines, clones) can contribute to an accurate estimation of genetic effect in DGY (Mulder et al., 2007; Hill and Mulder, 2010; Jung et al., 2020). In this aspect, the analyzed population is a good source to obtain accurate estimates since several repetitions of each line were available, and there is a high genetic similarity between replications of F₆ lines. The high genetic similarity among replications occurs due to the F₆ lines originated from selfing of F_{3.5} lines, where segregation is very low as product of the high homozygosity due to previous generations of selfing (expected ~96.9%).

Genetic parameters for micro-environmental sensitivity and implications for breeding activities

Understanding the genetics of micro-environmental sensitivity for grain yield in wheat can be useful to optimize breeding strategies and improve the adaptation of wheat cultivars to the environments

where production is performed. The DHGLM allowed estimating genetic parameters associated with micro-environmental sensitivity such as h_d^2 and GCV_E which are discussed in this section.

The Mulder-Hill heritability of residual variance (h_d^2) is defined as the regression coefficient of the additive genetic values for the dispersion variable on the squared phenotypic deviation (Mulder et al., 2007) as an analogy to the heritability in the classical sense (Falconer and Mackay, 1996). The h_d^2 can be interpreted as an indication of the proportion of the total variance that is explained by genetic variability in residual dispersion ($\hat{\sigma}_{g_d}^2$). In general, h_d^2 estimates are expected to be low, and based on empirical studies, it has been in the range of <0.01 to 0.10 for different species (reviewed by Hill and Mulder, 2010, and Jung et al., 2020). In our study, we found a h_d^2 of 0.033, which is in the intermediate range reported in previous studies and implies that reducing micro-environmental sensitivity by selection is possible. For the interpretation of results, it is also relevant to consider that other potential sources of variation in DGY could affect the estimation of genetic parameters with an impact on decreasing or increasing the amount of residual genetic variance captured. For example, genetic variation due to G \times E interactions could also be present at the level of DGY. To address this issue, we have initially included in the model a G \times E interaction effect modeled with a genomic relationship matrix within each environment, and a L \times E interaction effect (as used to model EGY), however, those effects were later excluded due to convergence problems in the REML algorithm. An additional source of variation on residuals that could affect genetic parameters is a scale effect for higher means inducing higher variances in residuals (Falconer and Mackay, 1996; Yang et al., 2011; Mulder et al., 2016). This issue is further discussed in the next section “Genetic correlation between additive effects in EGY and DGY”.




FIGURE 4

Predictive ability (PA) for the linear mixed model allowing heteroscedasticity of residual variance for the different environments (LMM-HET) and double hierarchical generalized linear model (DHGLM) in a leave-one-line-out (LOO) cross-validation. g : predicted additive genetic values for corrected phenotypes (y_c). g_d : predicted additive genetic values for dispersion variable (y_d). Black bars represent the 95% confidence interval (CI) computed for each estimate as the standard deviation of the estimate multiplied by 1.96. The letter above the bar denotes significant differences between models in a t-test (P -value < 0.01). The horizontal green lines represent the theoretical maximum PAs.




FIGURE 5

Correlation between predicted additive genetic values obtained with whole phenotypic information and additive values obtained with partial phenotypic information ($r_{w,p}$) in a leave-one-line-out (LOO) cross-validation. LMM-HET: linear mixed model allowing heteroscedasticity of residual variance for the different environments. DHGLM: double hierarchical generalized linear model. g : predicted additive genetic values for corrected phenotypes (y_c). g_d : predicted additive genetic values for dispersion variable (y_d). Black bars represent the 95% confidence interval (CI) computed for each estimate as the standard deviation of the estimate multiplied by 1.96. The letter above the bar denotes significant differences between models in a t-test (P -value < 0.01).

The genetic coefficient of variation (GCV_E) allows inferring the potential for reducing micro-environmental sensitivity through selection (Mulder et al., 2007; Rönnegård et al., 2013; Mulder et al., 2016; Jung et al., 2017). Previous studies revealed large variation in the GCV_E across traits and species ranging from <0.01 to 0.86 (as reviewed by Sae-Lim et al., 2016 and Jung et al., 2020). In our population, we found a GCV_E of 0.061, which implies that reducing the average additive genetic values in dispersion in one unit of GCV_E will reduce the residual variance by 6.1% (Mulder et al., 2009). The GCV_E , is in addition, informative since (as the classical coefficient of variation) it can be compared across traits and species (Sonesson et al., 2013).

As revealed by h_d^2 and GCV_E , selecting lines with lower micro-environmental sensitivity is possible, and thus, it could be included within breeding goals to develop more resilient cultivars. This could be particularly valuable under future climate conditions where more variable and extreme weather events are expected. On the contrary, if genotype sensitivity is not considered for selection, and wheat lines are selected by their superior performance in specific environments, the selection could induce an indirect increase in sensitivity product of increasing the G×E component of selected lines (Falconer and Mackay, 1996).

Genetic correlation between additive effects in EGY and DGY

The knowledge of the genetic correlation between additive effects in EGY and DGY (ρ_{g,g_d}) is relevant to infer possible directions of micro-environmental sensitivity in breeding programs and guide selection decisions. A positive-sign ρ_{g,g_d} represent an unfavorable genetic correlation since it means that selecting for lines with higher additive value in EGY will result in higher micro-environmental sensitivity. In our

study, we found an estimate of ρ_{g,g_d} of 0.502 (Table 2); however, several authors have argued that, to some extent, ρ_{g,g_d} can be influenced by scale effects, where higher variances in residuals are associated with higher means (Falconer and Mackay, 1996; Yang et al., 2011; Mulder et al., 2016). The scale effect may be present if the coefficient of variation of a trait remains constant as its mean change, then its variance must change as well (Walsh and Lynch, 2018). An alternative approach to evaluate ρ_{g,g_d} without the potential trend caused by scale effects is using the natural logarithm (ln) of the main response variable (YGD) to perform the analysis (Sonesson et al., 2013; Sae-Lim et al., 2015; Sae-Lim et al., 2017). The log-transformation reduces the dependency of variance on mean due to scale, and thus ρ_{g,g_d} is expected to be mainly influenced by genetic micro-environmental sensitivity and not by the scale effect.

To evaluate how the scale effect affects our population, we have performed an additional analysis using the natural logarithm (ln) of y_c . The log-transformed y_c variable was observed normally distributed, and therefore the same model assumptions were made as for y_c . After log-transformation, we found a ρ_{g,g_d} value of -0.698. Similar results in terms of switching ρ_{g,g_d} from a positive to negative correlation after data transformation has been observed in previous studies for other species (Yang et al., 2011; Sae-Lim et al., 2015; Sae-Lim et al., 2017). The analysis based on ln (y_c) is expected to yield a better estimate of ρ_{g,g_d} as it is free from distortions caused by the scale effect, and therefore, it is a more accurate measure of correlation due to genetic effects. The negative correlation found is favorable for reducing micro-environmental sensitivity in breeding programs since selection for higher additive values in EGY will result in lower additive values in DGY for log-transformed y_c . In addition, we have observed a correlation close to 1 (0.99) between predictions for additive values in y_c and log-tranformed y_c . Based on these results, we concluded that select on both the EGY and DGY is possible and could represent a

comprehensive strategy to increase grain yield and reduce micro-environmental sensitivity in winter wheat.

Genomic prediction for grain yield and its micro-environmental sensitivity

The PA for predictions of additive genetic effect in EGY and DGY were evaluated in a leave-one-line-out (LOO) CV (Figure 4). No significant differences (P -value > 0.01) in PA for additive genetic effects in EGY were observed between the LMM-HET and DHGLM, and a similar trend was observed for the maximum potential PA (horizontal green lines in Figure 4). A small but significant improvement (P -value < 0.01) of 0.5% in $r_{w,p}$ was conferred by the DHGLM compared to the LMM-HET (Figure 5), and no variance inflation ($b_{w,p}$) was observed for predictions of additive effect in EGY (predictions of g effect) for any of the proposed models ($b_{w,p}$ values close to 1, Figure 6). The observed results revealed a good performance for predictions of additive genetic effect in EGY with both developed models. The good predictive performance for additive genetic effect in EGY in the DHGLM has also been observed in previous studies, especially when genomic information was used (Mulder et al., 2013a; Sell-Kubiak et al., 2015; Sae-Lim et al., 2017). In addition, the similar performance for prediction of breeding values for EGY using the LMM-HET and the DHGLM suggests that accounting for heterogeneous residuals by genotype in the DHGLM and the use of genetic correlations between additive effect in EGY and DGY in the DHGLM have not provided significant benefits for prediction.

The PA for additive genetic effects in DGY (predictions of g_d effect) was significantly lower than for additive genetic effect in EGY (Figure 4). This can be expected because given the low h_d^2 found, the potential PA of predicted additive values in dispersion is low (~0.48). The relationship between the trait's heritability and prediction accuracy has been reported in the literature, where for lower values of heritability, lower accuracies are expected (Wimmer et al., 2013; Muranty et al., 2015). The $r_{w,p}$ estimate for prediction of additive effects in DGY was high (0.905); however, it was significantly lower (about 4%, P -value < 0.01) than for additive effects in EGY. The $b_{w,p}$ for predictions in DGY revealed a low over-dispersion for predicted values as observed in a $b_{w,p}$ value below one (0.935). A potential reason explaining over-dispersion could be related to the presence of other effects affecting DGY that could not be included in the model. For example, as earlier stated, genetic variation due to G×E interactions could be present at the level of DGY. However, those effects could not be included for our population due to problems of convergence with the REML algorithm.

Limitations and prospects

The presented work can be seen as a proof of concept for studying micro-environmental sensitivity in crop breeding programs. Breeding program datasets could be reanalyzed in order to identify and select more resilient lines exhibiting lower micro-environmental sensitivity. Selecting for reduced micro-environmental sensitivity is a way to deal with/handle some specific elements of G×E. The variability at macro-




FIGURE 6

Boxplot of bootstrap distribution for the slope of the regression of additive genetic values obtained with whole phenotypic information on additive values obtained with partial phenotypic information ($b_{w,p}$) in a leave-one-line-out (LOO) cross-validation. LMM-HET: linear mixed model allowing heteroscedasticity of residual variance for the different environments. DHGLM: double hierarchical generalized linear model. g : predicted additive genetic values for corrected phenotypes (y_c). g_d : predicted additive genetic values for dispersion variable (y_d). The letter above the bar denotes significant differences between models in a t-test (P -value < 0.01). The black dashed line represents a regression coefficient of one, where no under or over-dispersion is present.

environmental scale is another important source of G×E in crops. Future studies extending the proposed DHGLM to account for macro-environmental sensitivity could represent a comprehensive approach to reducing sensitivity at micro- and macro-environmental levels. Furthermore, this may provide a better understanding of the relationship between micro- and macro-environmental sensitivity, and hence more research in the area is justified.

In addition, other G×E related effects and spatial effects were initially attempted to be included in the DHGLM. However, we found limitations for algorithm convergence when increasing model complexity by adding these G×E effects and prohibitive computational times when modelling spatial effects. The prohibitive computational times could be due to the spatial effect and the additive genetic effect for DGY (g_a) partially competing for the same variation (i.e. confounding factors between the spatial effect at the observation plot and micro-environmental variation). Consequently, the spatial effects were accounted for in a two-step approach, first estimating them in a separated linear mixed model and second subtracting their estimates from the raw phenotypes. If confounding between spatial effect and micro-environmental sensitivity effectively occurs, either not correcting for the spatial effect could inflate the estimates for micro-environmental sensitivity, or correcting by them beforehand (i.e. as in our study) would lead to a conservative estimate of the variance of micro-environmental sensitivity. The conservative estimation in our study could imply a stronger micro-environmental sensitivity than we estimated, and therefore, it may be even more important for wheat breeding. Therefore, further work is useful in order to improve model inference.

Conclusions

In this work, we used a double hierarchical generalized linear model (DHGLM) to study the micro-environmental sensitivity for grain yield in wheat. As far as we know, studies on micro-environmental sensitivity have been scarcely conducted in plants, and this is the first time that genetic parameters for micro-environmental sensitivity have been investigated in wheat. We found that the micro-environmental sensitivity for grain yield is heritable and that there is potential for its reduction, according to a Mulder-Hill heritability (\hat{h}_d^2) of 0.033 and a genomic coefficient of variation (GCV_E) of 0.061, respectively. The genetic correlation between additive effects for (expressed) grain yield and its residual dispersion (ρ_{gg_e}) was estimated, and we observed a correlation of 0.502. Further analysis using log-transformation of (expressed) phenotypes was performed to study a possible scale effect of higher dispersion variances induced by higher phenotypic values, and a negative genetic correlation was observed after transformation (-0.698). The estimate of ρ_{gg_e} using the log-transformation is a more reliable estimate of genetic correlation as it is free from distortions caused by the scale effect. Based on these results, we concluded that breeding for reduced micro-environmental sensitivity is possible and can be included within breeding objectives without compromising the selection for increased yield. In addition, the double hierarchical generalized linear model had a good predictive performance for additive effects on residual dispersion, and showed similar performance for predicting additive genetic effects on (expressed) grain yield compared to a linear mixed model allowing for heteroscedasticity of residual variance in different environments (LMM-HET). Such findings showed that the double hierarchical generalized linear models could be a good choice to predict

additive genetic effects on (expressed) grain yield and its residual dispersion for wheat breeding.

Data availability statement

The original contributions presented in the study are publicly available. This data can be found here: <https://doi.org/10.7910/DVN/AB43I3>.

Author contributions

JJ and MR: conceptualization. MR and PS: data curation. MR: formal analysis. JJ, MR: funding acquisition. MR, JJ, BC, PS, RR, LM, TM-H: investigation, methodology, and project administration. JJ and PS: resources and supervision. MR, JJ: software, validation, visualization, and writing—original draft. JJ, BC, PS, RR, LM, TM-H: writing—review and editing. All authors contributed to the article and approved the submitted version.

Funding

This research received funding from the National Research and Innovation Agency (ANII) of Uruguay (POS_EXT_2018_1_154284).

Acknowledgments

We thank Aarhus University for financial support for authors MR, and JJ, and the National Institute for Agricultural, Food and Environmental Research (INRAE) of France for financial support of BC, RR, LM, and TM-H. We also thank Nordic Seed A/S breeding company for PS funding, and Jeppe R. Andersen, J. Orabi and A. Jahoor from Nordic Seed A/S for providing phenotypic and genotypic data. Financial support from the National Research and Innovation Agency (ANII) of Uruguay for the PhD scholarship of author MR (code POS_EXT_2018_1_154284) is greatly acknowledged.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpls.2022.1075077/full#supplementary-material>

References

- Bhatt, G., and Derera, N. (1975). Genotype x environment interactions for, heritabilities of, and correlations among quality traits in wheat. *Euphytica* 24 (3), 597–604. doi: 10.1007/BF00132896
- Cooper, M., Woodruff, D., Eisemann, R., Brennan, P., and DeLacy, I. (1995). A selection strategy to accommodate genotype-by-environment interaction for grain yield of wheat: managed-environments for selection among genotypes. *Theor. Appl. Genet.* 90 (3), 492–502. doi: 10.1007/BF00221995
- Crossa, J., Fritsche-Neto, R., Montesinos-Lopez, O. A., Costa-Neto, G., Dreisigacker, S., Montesinos-Lopez, A., et al. (2021). The modern plant breeding triangle: Optimizing the use of genomics, phenomics, and enviromics data. *Front. Plant Sci.* 12. doi: 10.3389/fpls.2021.651480
- Crossa, J., Perez-Rodriguez, P., Cuevas, J., Montesinos-Lopez, O., Jarquin, D., de Los Campos, G., et al. (2017). Genomic selection in plant breeding: Methods, models, and perspectives. *Trends Plant Sci.* 22 (11), 961–975. doi: 10.1016/j.tplants.2017.08.011
- Dworkin, I. (2005). A study of canalization and developmental stability in the sternopleural bristle system of *Drosophila melanogaster*. *Evolution* 59 (7), 1500–1509. doi: 10.1111/j.0014-3820.2005.tb01799.x
- Ehsaninia, J., Hossein-Zadeh, N. G., and Shadparvar, A. A. (2019). Estimation of genetic variation for macro-and micro-environmental sensitivities of milk yield and composition in Holstein cows using double hierarchical generalized linear models. *J. Dairy. Res.* 86 (2), 145–153. doi: 10.1017/S0022029919000293
- Falconer, D., and Mackay, T. (1996). Introduction to quantitative genetics. Essex. UK: Longman. Group.
- Felleki, M., Lee, D., Lee, Y., Gilmour, A. R., and Rönnegård, L. (2012). Estimation of breeding values for mean and dispersion, their variance and correlation using double hierarchical generalized linear models. *Genet. Res.* 94 (6), 307–317. doi: 10.1017/S0016672312000766
- Garcia, M., David, I., Garreau, H., Ibañez-Escriche, N., Mallard, J., Pommeret, D., et al. (2009). “Comparisons of three models for canalising selection or genetic robustness,” in *60. annual meeting of the European association for animal production* (Wageningen Academic Publishers).
- Guo, X., Svane, S. F., Füchtbauer, W. S., Andersen, J. R., Jensen, J., and Thorup-Kristensen, K. (2020). Genomic prediction of yield and root development in wheat under changing water availability. *Plant Methods* 16 (1), 1–15. doi: 10.1186/s13007-020-00634-0
- Hill, W. G., and Mulder, H. A. (2010). Genetic analysis of environmental variation. *Genet. Res.* 92 (5-6), 381–395. doi: 10.1017/S0016672310000546
- Hill, W. G., and Zhang, X.-s. (2004). Effects on phenotypic variability of directional selection arising through genetic differences in residual variability. *Genet. Res.* 83 (2), 121–132. doi: 10.1017/S0016672304006640
- Hoaglin, D. C., and Welsch, R. E. (1978). The hat matrix in regression and ANOVA. *Am. Statistician.* 32 (1), 17–22. doi: 10.1080/00031305.1978.10479237
- Iung, L., Carvalho, R., Neves, H., and Mulder, H. A. (2020). Genetics and genomics of uniformity and resilience in livestock and aquaculture species: A review. *J. Anim. Breed. Genet.* 137 (3), 263–280. doi: 10.1111/jbg.12454
- Iung, L., Neves, H., Mulder, H., and Carvalho, R. (2017). Genetic control of residual variance of yearling weight in nellore beef cattle. *J. Anim. Sci.* 95 (4), 1425–1433. doi: 10.2527/jas.2016.1326
- Lee, Y., and Nelder, J. A. (2006). Double hierarchical generalized linear models (with discussion). *J. R. Stat. Soc.: Ser. C. (Applied Statistics).* 55 (2), 139–185. doi: 10.1111/j.1467-9876.2006.00538.x
- Legarra, A., and Reverter, A. (2018). Semi-parametric estimates of population accuracy and bias of predictions of breeding values and future phenotypes using the LR method. *Genet. Sel. Evol.* 50 (1), 53. doi: 10.1186/s12711-018-0426-6
- Lopez-Cruz, M., Crossa, J., Bonnett, D., Dreisigacker, S., Poland, J., Jannink, J.-L., et al. (2015). Increased prediction accuracy in wheat breeding trials using a marker x environment interaction genomic selection model. *G3: Genes, Genomes, Genet.* 5 (4), 569–582. doi: 10.1534/g3.114.016097
- Ly, D., Chenu, K., Gauffreteau, A., Rincet, R., Huet, S., Gouache, D., et al. (2017). Nitrogen nutrition index predicted by a crop model improves the genomic prediction of grain number for a bread wheat core collection. *Field Crops Res.* 214, 331–340. doi: 10.1016/j.fcr.2017.09.024
- Ly, D., Huet, S., Gauffreteau, A., Rincet, R., Touzy, G., Mini, A., et al. (2018). Whole-genome prediction of reaction norms to environmental stress in bread wheat (*Triticum aestivum* L.) by genomic random regression. *Field Crops Res.* 216, 32–41. doi: 10.1016/j.fcr.2017.08.020
- Madsen, P., and Jensen, J. (2013). “An user’s guide to DMU, version 6, release 5.1. center for quantitative genetics and genomics,” in *Dept. Of molecular biology and genetics, university of Aarhus* (Denmark: Research Centre Foulum Tjele).
- Madsen, M., van der Werf, J., Börner, V., Mulder, H., and Clark, S. (2021). Estimation of macro-and micro-genetic environmental sensitivity in unbalanced datasets. *Animal* 15 (12), 100411. doi: 10.1016/j.animal.2021.100411
- Morgante, F., Sorensen, P., Sorensen, D. A., Maltecca, C., and Mackay, T. F. C. (2015). Genetic architecture of micro-environmental plasticity in *Drosophila melanogaster*. *Sci. Rep.* 5 (1), 9785. doi: 10.1038/srep09785
- Mulder, H., Bijma, P., and Hill, W. (2007). Prediction of breeding values and selection responses with genetic heterogeneity of environmental variance. *Genetics* 175 (4), 1895–1910. doi: 10.1534/genetics.106.063743
- Mulder, H., Crump, R., Calus, M., and Veerkamp, R. (2013a). Unraveling the genetic architecture of environmental variance of somatic cell score using high-density single nucleotide polymorphism and cow data from experimental farms. *J. Dairy. Sci.* 96 (11), 7306–7317. doi: 10.3168/jds.2013-6818
- Mulder, H. A., Gienapp, P., and Visser, M. E. (2016). Genetic variation in variability: Phenotypic variability of fledging weight and its evolution in a songbird population. *Evolution* 70 (9), 2004–2016. doi: 10.1111/evo.13008
- Mulder, H., Hill, W., Vereijken, A., and Veerkamp, R. (2009). Estimation of genetic variation in residual variance in female and male broiler chickens. *Animal* 3 (12), 1673–1680. doi: 10.1017/S1751731109990668
- Mulder, H. A., Rönnegård, L., Fikse, W. F., Veerkamp, R. F., and Strandberg, E. (2013b). Estimation of genetic variance for macro-and micro-environmental sensitivity using double hierarchical generalized linear models. *Genet. Selection. Evol.* 45 (1), 1–14. doi: 10.1186/1297-9686-45-23
- Muranty, H., Troggo, M., Sadok, I. B., Rifai, M. A., Auwerkerken, A., Banchi, E., et al. (2015). Accuracy and responses of genomic selection on key traits in apple breeding. *Horticult. Res.* 2. doi: 10.1038/hortres.2015.60
- Ordas, B., Malvar, R. A., and Hill, W. G. (2008). Genetic variation and quantitative trait loci associated with developmental stability and the environmental correlation between traits in maize. *Genet. Res.* 90 (5), 385–395. doi: 10.1017/S0016672308009762
- Raffo, M., Sarup, P., Andersen, J., Orabi, J., Jahoor, A., and Jensen, J. (2022a). Integrating a growth degree-days based reaction norm methodology and multi-trait modeling for genomic prediction in wheat. *Front. Plant Sci.* 13. doi: 10.3389/fpls.2022.939448
- Raffo, M. A., Sarup, P., Guo, X., Liu, H., Andersen, J. R., Orabi, J., et al. (2022b). Improvement of genomic prediction in advanced wheat breeding lines by including additive-by-additive epistasis. *Theor. Appl. Genet.*, 1–14. doi: 10.21203/rs.3.rs-424490/v1
- Ray, D. K., Mueller, N. D., West, P. C., and Foley, J. A. (2013). Yield trends are insufficient to double global crop production by 2050. *PLoS One* 8 (6), e66428. doi: 10.1371/journal.pone.0066428
- Rogers, S. O., and Bendich, A. J. (1985). Extraction of DNA from milligram amounts of fresh, herbarium and mummified plant tissues. *Plant Mol. Biol.* 5 (2), 69–76. doi: 10.1007/BF00020088
- Rönnegård, L., Felleki, M., Fikse, F., Mulder, H. A., and Strandberg, E. (2010). Genetic heterogeneity of residual variance-estimation of variance components using double hierarchical generalized linear models. *Genet. Selection. Evol.* 42 (1), 1–10. doi: 10.1186/1297-9686-42-8
- Rönnegård, L., Felleki, M., Fikse, W., Mulder, H., and Strandberg, E. (2013). Variance component and breeding value estimation for genetic heterogeneity of residual variance in Swedish Holstein dairy cattle. *J. Dairy. Sci.* 96 (4), 2627–2636. doi: 10.3168/jds.2012-6198
- Roozeboom, K. L., Schapaugh, W. T., Tuinstra, M. R., Vanderlip, R. L., and Milliken, G. A. (2008). Testing wheat in variable environments: genotype, environment, interaction effects, and grouping test locations. *Crop Sci.* 48 (1), 317–330. doi: 10.2135/cropsci2007.04.0209
- Sae-Lim, P., Gjerde, B., Nielsen, H. M., Mulder, H., and Kause, A. (2016). A review of genotype-by-environment interaction and micro-environmental sensitivity in aquaculture species. *Rev. Aquacult.* 8 (4), 369–393. doi: 10.1111/raq.12098
- Sae-Lim, P., Kause, A., Janhunen, M., Vehviläinen, H., Koskinen, H., Gjerde, B., et al. (2015). Genetic (co) variance of rainbow trout (*Oncorhynchus mykiss*) body weight and its uniformity across production environments. *Genet. Selection. Evol.* 47 (1), 1–10. doi: 10.1186/s12711-015-0122-8
- Sae-Lim, P., Kause, A., Lillehammer, M., and Mulder, H. A. (2017). Estimation of breeding values for uniformity of growth in Atlantic salmon (*Salmo salar*) using pedigree relationships or single-step genomic evaluation. *Genet. Selection. Evol.* 49 (1), 1–12. doi: 10.1186/s12711-017-0308-3
- SanCristobal-Gaudy, M., Bodin, L., Elsen, J.-M., and Chevalet, C. (2001). Genetic components of litter size variability in sheep. *Genet. Selection. Evol.* 33 (3), 249–271. doi: 10.1186/1297-9686-33-3-249
- SanCristobal-Gaudy, M., Elsen, J.-M., Bodin, L., and Chevalet, C. (1998). Prediction of the response to a selection for canalisation of a continuous trait in animal breeding. *Genet. Selection. Evol.* 30 (5), 423–451. doi: 10.1186/1297-9686-30-5-423
- Sell-Kubiak, E., Bijma, P., Knol, E., and Mulder, H. (2015). Comparison of methods to study uniformity of traits: application to birth weight in pigs. *J. Anim. Sci.* 93 (3), 900–911. doi: 10.2527/jas.2014-8313
- Sell-Kubiak, E., Knol, E. F., and Lopes, M. (2022). Evaluation of the phenotypic and genetic background of variability based on litter size of Large white pigs. *Genet. Selection. Evol.* 54 (1), 1–15. doi: 10.1186/s12711-021-00692-5
- Sial, M., Arain, M., and Ahmad, M. (2000). Genotypex environment interaction on bread wheat grown over multiple sites and years in Pakistan. *Pakistan J. Bot.* 32 (1), 85–91.
- Sonesson, A. K., Ødegård, J., and Rönnegård, L. (2013). Genetic heterogeneity of within-family variance of body weight in Atlantic salmon (*Salmo salar*). *Genet. Selection. Evol.* 45 (1), 1–8. doi: 10.1186/1297-9686-45-41

- Sorensen, D., and Waagepetersen, R. (2003). Normal linear models with genetically structured residual variance heterogeneity: a case study. *Genet. Res.* 82 (3), 207–222. doi: 10.1017/S0016672303006426
- Sukumaran, S., Crossa, J., Jarquín, D., and Reynolds, M. (2017). Pedigree-based prediction models with genotype × environment interaction in multi-environment trials of CIMMYT wheat. *Crop Sci.* 57 (4), 1865–1880. doi: 10.2135/cropsci2016.06.0558
- Tilman, D., Balzer, C., Hill, J., and Befort, B. L. (2011). Global food demand and the sustainable intensification of agriculture. *Proc. Natl. Acad. Sci.* 108 (50), 20260–20264. doi: 10.1073/pnas.1116437108
- Vandenplas, J., Bastin, C., Gengler, N., and Mulder, H. (2013). Genetic variance in micro-environmental sensitivity for milk and milk quality in Walloon Holstein cattle. *J. Dairy Sci.* 96 (9), 5977–5990. doi: 10.3168/jds.2012-6521
- VanRaden, P. M. (2008). Efficient methods to compute genomic predictions. *J. Dairy Sci.* 91 (11), 4414–4423. doi: 10.3168/jds.2007-0980
- Walsh, B., and Lynch, M. (2018). *Evolution and selection of quantitative traits* (Oxford University Press).
- Wang, S., Wong, D., Forrest, K., Allen, A., Chao, S., Huang, B. E., et al. (2014). Characterization of polyploid wheat genomic diversity using a high-density 90 000 single nucleotide polymorphism array. *Plant Biotechnol. J.* 12 (6), 787–796. doi: 10.1111/pbi.12183
- Wimmer, V., Lehermeier, C., Albrecht, T., Auinger, H.-J., Wang, Y., and Schön, C.-C. (2013). Genome-wide prediction of traits with different genetic architecture through efficient variable selection. *Genetics* 195 (2), 573–587. doi: 10.1534/genetics.113.150078
- Wolc, A., White, I., Avendano, S., and Hill, W. (2009). Genetic variability in residual variation of body weight and conformation scores in broiler chickens. *Poultry Sci.* 88 (6), 1156–1161. doi: 10.3382/ps.2008-00547
- Yang, Y., Christensen, O. F., and Sorensen, D. (2011). Analysis of a genetically structured variance heterogeneity model using the box-cox transformation. *Genet. Res.* 93 (1), 33–46. doi: 10.1017/S0016672310000418