



**HAL**  
open science

## A multi-approach improves the description of the entomopathogenic nematodes distribution in corn fields

Elisabeth Depuydt, Jean-Claude Ogier, Lisa Cabre, Nusrat Ali, Patrice Mahieu, Cécile Villenave, Eric Nguema-Ona, Sophie Gaudriault

### ► To cite this version:

Elisabeth Depuydt, Jean-Claude Ogier, Lisa Cabre, Nusrat Ali, Patrice Mahieu, et al.. A multi-approach improves the description of the entomopathogenic nematodes distribution in corn fields. 16. Meeting of the IOBC-WRPS Working Group Biological and Integrated Control of Plant Pathogens, Jun 2023, Wageningen (NL), Netherlands. hal-04129939

**HAL Id: hal-04129939**

**<https://hal.inrae.fr/hal-04129939v1>**

Submitted on 15 Jun 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# A multi-approach improves the description of the entomopathogenic nematodes distribution in corn fields

Elisabeth Depuydt<sup>1,2</sup>, Jean-Claude Ogier<sup>1</sup>, Lisa Cabre<sup>2</sup>, Nusrat Ali<sup>3</sup>, Patrice Mahieu<sup>4</sup>, Cécile Villenave<sup>5</sup>, Eric Nguema-Ona<sup>2</sup>, Sophie Gaudriault<sup>1</sup>

<sup>1</sup>DGIMI, Univ Montpellier, INRAE, Montpellier, France

<sup>2</sup>Agro Innovation International TIMAC AGRO, Plant Nutrition Laboratory, Crop Management under Biotic and Abiotic Stress Team, Saint Malo, France

<sup>3</sup>Agro Innovation International TIMAC AGRO, Phys-Chem and BioAnalytics Department, Saint Malo, France

<sup>4</sup>Chamber of Agriculture, Pau, France


<sup>5</sup>ELISOL Environnement, Congénies, France

## Introduction


Entomopathogenic nematodes (EPNs), *Steinernema* sp. and *Heterorhabditis* sp., are soil-dwelling insect parasites distributed in all continents except in Antarctica. Moreover, the detection method may bias the diagnosis of the presence of EPNs in soils. The classical method for the detection of EPNs is the baiting method by *Galleria mellonella*. **The objective of this study is to combine several methods to improve the detection of EPNs in cultivated soils. We assumed that a multi-approach which combines baiting with successive trapping cycles (Abd-Elgawad, 2020, Nematology) and molecular detection (Campos-Herrera et al., 2011, Annals of Applied Biology) would provide the most accurate picture of the presence and diversity of EPNs in the sampled soils.**

## Sampling map

Map of the 43 sampling plots cultivated in corn in South Western France




## Experimental procedure




## Results

EPNs detection by successive trapping cycles (28%)




Three species were isolated: *S. carpocapsae*, *S. feltiae* and *H. bacteriophora*. The multiplication of the baiting cycles allowed to increase the number of positive plots and the species diversity of EPNs isolated


**Steinernema sp. detection by qPCR (19%)**



**Steinernema sp. detection by ddPCR (33%)**



Common/different positive plots between the three detection methods



**By combining all methods we obtained 49% of positive plots for *Steinernema***

## Conclusion

- ❖ Mismatches are observed between the detection methods (e.g. molecular detection of EPNs in negative plots by baiting).
- ❖ Using of multi-approach improves the detection of native EPNs in soils.
- ❖ A good picture of the distribution of EPNs in agricultural soils is useful for their use in biological control by conservation.