

Building better knowledge on meat quality determination through integrated data mining and curation of proteomics studies

Mohammed Gagaoua

▶ To cite this version:

Mohammed Gagaoua. Building better knowledge on meat quality determination through integrated data mining and curation of proteomics studies. 25. Congress of the Animal Science and Production Association (ASPA)., Jun 2023, Monopoli, Italy. pp.15. hal-04132114

HAL Id: hal-04132114 https://hal.inrae.fr/hal-04132114

Submitted on 18 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





ASPA 25th Congress Monopoli (BARI - ITALY), June 13-16, 2023

#ASPA2023 ASPA 25th Congress Book of Abstract

The 25th congress of the Animal Science and Production Association

"Animal Production Science: Innovations and sustainability for future generation" is under patronage of Loghi patrocini

Monopoli (BARI – ITALY), June 13–16, 2023

Venue

Torre Cintola Natural Sea Emotions Località Capitolo – Monopoli (BARI – ITALY)



reducing ruminal NH₃-N production and nitrogen excretion in urine as well as slowing dietary protein breakdown in the rumen and improving amino acid bioavailability in the small intestine. The production of ruminants and feed digestion in rumen can both be favorably influenced by feed additives such as probiotics and products derived from plants. These additives come in a variety of forms and work in several ways. Despite their diversity, they all eventually have an impact on the metabolic pathways for fermentation and/or the microbial community in the digestive tract. The kind of feed used, and the animal's physiological condition have an impact on its effectiveness. A fully comprehend complicated interactions that find and take place between hosts, feeds, and additives. This knowledge will allow us to identify the potential applications of a particular feed additive, hence minimizing inconsistent field responses.

KEYWORDS: Feed additives; ruminants; secondary metabolites; mode of action

Building better knowledge on meat quality determination through integrated data mining and curation of proteomics studies

Mohammed Gagaoua
PEGASE, INRAE, Saint-Gilles, France

Meat quality is very important for consumers, processors, and producers. The traditional methods for assessing meat quality are subjective and time-consuming, and they do not provide a comprehensive understanding of the factors that affect meat quality. Proteomics, a high-throughput technology for the analysis of entire proteins, has emerged as a powerful tool for investigating the molecular mechanisms underlying the conversion of muscle into meat, meat quality variability, and the discovery of biomarkers. However, the large amount of data generated by meat research proteomics experiments in the last two decades can be overwhelming, and the integration of published meat proteomics datasets through data mining approaches and bioinformatics is a promising way to expand our scientific knowledge. Furthermore, mining such data in their biological context may allow the extraction of novel and more robust biomarkers of meat quality traits. Data reuse is a leading, active, and evolving field suggested and very recently applied in the meat research field to rediscover and reshape the public proteomics data for a better understanding of the unknowns related to meat quality determination and refinement of the list of biomarkers so far proposed. In this talk, the application of integrated data mining and curation approaches to improve our understanding of the molecular basis of meat quality will be presented. The focus will be on important meat quality traits and their defects such as tenderness, color and post-mortem processing treatments (e.g. electrical stimulation). The most updated databases, patterns, knowledge and relationships between different factors and meat quality gained thanks to the integromics studies will be presented. Overall, integrated data mining and proteomics studies curation allowed for providing a comprehensive view of the biochemical mechanisms and molecular signatures underlying meat quality, which all together allowed better insights of their affecting factors. These novel approaches further allowed proposing robust biomarkers for the development of more accurate and efficient methods/tools for assessing meat quality. Integrated data mining and curation of proteomics studies will have important implications for the meat industry, as they will provide new insights into the factors that affect meat quality.

Technologies applied in embryonated eggs

Katarzyna Stadnicka

Faculty of Health Sciences, Nicolaus Copernicus University in Toruń, Ludwik Rydygier Collegium Medicum in Bydgoszcz, Bydgoszcz, Poland

Implementation of precise solutions to develop natural and sustainable 'immunity' in the animal is an element of a systemic approach defined by World Health Organization, and referred to as the One Health. Stakeholders of poultry production seek solutions to eliminate and/or control the pathogens transmitted with food and animals. Intentional stimulation of embryonic development and determination of the future post-embryonic health of the organism is possible by in ovo application of natural antioxidants and prebiotics, gut stabilizers like probiotics and other immunological enhancements, including vaccines. In light of the above, advancements in biotechnology and robotization have led to a fast development of several break-through practices in poultry hatcheries: in ovo vaccination, in ovo feeding, stimulation of embryonic development using prebiotics, probiotics and synbiotics, thermal manipulations and embryo sex determination. The work flow of a modern hatchery is almost fully automatized, and as such serves as an excellent, controlled space for implementation of novel in ovo techniques with elimination of human error and potentially adverse environmental factors. A post-embryonic life of a broiler chicken is limited to about 42 days. Therefore, considering the complete period from egg fertilization to animal maturity, the embryonic development in egg (21 days) constitutes a one third of a total life span of the animal. This biological feature enables poultry production to be highly transformative and allows for a shift of some of certain treatments from post

