



HAL
open science

Association of two co-occurring mutations at the AvrMlp7 avirulence locus in poplar rust during the breakdown of RMlp7 resistance gene

Louet Clémentine, Saubin Méline, Andrieux Axelle, Persoons Antoine, Gorse Mathilde, Pétrowski Jérémy, Mita Stéphane De, Fabre Bénédicte, Duplessis Sébastien, Frey Pascal, et al.

► To cite this version:

Louet Clémentine, Saubin Méline, Andrieux Axelle, Persoons Antoine, Gorse Mathilde, et al.. Association of two co-occurring mutations at the AvrMlp7 avirulence locus in poplar rust during the breakdown of RMlp7 resistance gene. IS-MPMI Congress eSymposium, Pathogen-Host Coevolution - Combating Resistance Breaking in Agriculture, Sep 2021, Online, United States. , 2021, 2021 IS-MPMI Congress: eSymposia Series Abstracts. 10.1094/MPMI-34-9-S2.1 . hal-04137548

HAL Id: hal-04137548

<https://hal.inrae.fr/hal-04137548v1>

Submitted on 6 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Association of two co-occurring mutations at the *AvrMlp7* avirulence locus in poplar rust during the breakdown of *RMlp7* resistance gene

Clémentine Louet, Méline Saubin, Axelle Andrieux, Antoine Persoons, Mathilde Gorse*, Jérémy Pétrowski, Stéphane De Mita**, Bénédicte Fabre, Sébastien Duplessis, Pascal Frey and Fabien Halkett

Université de Lorraine, INRAE, IAM, F-54000 Nancy, France

*Present address: INRAE, AgroParisTech, Université Paris-Saclay, BIOGER, F-78850 Thiverval-Grignon, France

**Present address: INRAE, Cirad, IRD, Montpellier SupAgro, Université de Montpellier, PHIM, F-34000 Montpellier, France

ABSTRACT

The deployment of plant varieties carrying resistance (*R*) genes exerts strong selection pressures towards pathogen populations. Several studies reported the rapid evolution of avirulence (*Avr*) genes to escape *R*-mediated plant immunity and identified a variety of mechanisms leading to virulence. The poplar rust fungus *Melampsora larici-populina* is the most damaging pathogen of poplars. A major adaptive event occurred in 1994 with the breakdown of *RMlp7* resistance gene in poplar. Population genomics studies identified a locus in the genome of *M. larici-populina*, which likely corresponds to the *AvrMlp7* candidate avirulence gene. We used a population genetics approach combined with dedicated qPCR assays on a comprehensive set of 281 isolates, covering 27 years (encompassing the resistance breakdown event), to validate the candidate locus and to assess its polymorphism. We found two mechanisms, a point mutation and a deletion, that allowed the pathogen to escape *RMlp7*-mediated resistance. Six diploid genotypes were thus characterized at the candidate locus (three avirulent and three virulent). In addition, a temporal analysis revealed that the two virulence alleles pre-existed (harboured as avirulent heterozygous genotypes) since the early samplings and were found in association (as virulent genotypes) at the time of the resistance breakdown. Our study documented that, in a diploid pathogen, combining virulence determinisms is adaptive.

Keywords

Melampsora larici-populina, molecular evolution, gene-for-gene interaction, pathogen, fungi