

Beneficial propionibacteria: from traditional cheeses to next generation probiotics

Gwénaël Jan, Benoît Foligné, Valérie Gagnaire, Gaëlle Boudry, Vasco Azevedo, Eric Guédon

▶ To cite this version:

Gwénaël Jan, Benoît Foligné, Valérie Gagnaire, Gaëlle Boudry, Vasco Azevedo, et al.. Beneficial propionibacteria: from traditional cheeses to next generation probiotics. IPC2023 - International Scientific Conference on Probiotics, https://probiotic-conference.net/scientific-committee/, Jun 2023, Bratislava (Slovaquie), Slovakia. hal-04141553

HAL Id: hal-04141553 https://hal.inrae.fr/hal-04141553

Submitted on 26 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Beneficial propionibacteria: from traditional cheeses to next generation probiotics

Authors: <u>Gwénaël JAN</u>¹, Benoit **FOLIGNÉ**², Valérie **GAGNAIRE**¹, Gaëlle **BOUDRY**³, Vasco **AZEVEDO**⁴, Eric **GUÉDON**¹

Affiliations:

- 1 INRAE, STLO, Institut Agro, Agrocampus Ouest, Rennes, France.
- 2 Univ. Lille, Inserm, CHU Lille, U1286-INFINITE-Institute for Translational Research in Inflammation, France.
- 3 Institut Numecan, INRAE, INSERM, Univ Rennes, Saint-Gilles-Rennes, France
- 4 Institute of Biological Sciences, Federal University of Minas Gerais (UFMG), Belo Horizonte, Brazil.

Swiss cheeses have been produced for centuries in a traditional way and their typical opening indicates the presence of propionibacteria. These last are indeed present in raw milk, their growth is favored by the specific technology of Swiss-type cheeses, and their high population in the ripened cheese is responsible for its characteristic sensory properties, including the opening and the aroma. In addition to this long history of safe use and consumption, they constitute a source of nutraceuticals. The specific propionic fermentation leads to the production of short chain fatty acids as main final obligatory products. These exert beneficial effects including nourishment of digestive epithelial cells, as well as promotion of differentiation and of apoptotic depletion of colon cancer cells. Propionibacteria also produce vitamins of the B group, which enhances the nutritional value of the food products they ferment. Some selected strains were moreover shown to produce immunomodulins, proteins able to modify the secretion of cytokines in human immune cells. These last were identified as proteinaceous components of propionibacteria surface layer. They also produce extracellular vesicles which modulate in vitro inflammation and the NFkB pathway in human intestinal epithelial cells. Accordingly, such strains modulate the immune response and mitigate inflammation. In mice, they prevent both mucositis and colitis, via such immunomodulins limiting acute inflammatory response and protecting the gut epithelial barrier. In pigs, their consumption also modulates the inflammatory response and the lymphocyte populations. Finally, clinical studies dealing with consumption of propionibacteria whey cultures suggest a beneficial role in the context of IBD. Propionibacteria strains were recently identified in the gut microbiota of healthy newborns, yet not in that of children with necrotizing enterocolitis. Indeed, propionibacteria were detected in human fecal samples, which means that propionibacteria may be traditional cheese starters or next generation probiotics. The growing knowledge of these peculiar actinobacteria, including their great strain-diversity, opens new avenues for the development of functional foods and of food supplements, in a general context of increasing incidence of ailments that include an inflammatory component. Moreover, their ability to ferment both dairy and vegetable matrices is presently investigated and the first promising advances open new perspectives for the development of functional fermented vegetable products.