



HAL
open science

In vitro digestion of two age-tailored dairy products in the aging gastrointestinal tract

Anaïs Lavoisier, Martine Morzel, Cindy Sounouvou, Bérénice Houinsou-Houssou, Chantal Septier, Carole Tournier, Gilles Feron, Didier Dupont

► To cite this version:

Anaïs Lavoisier, Martine Morzel, Cindy Sounouvou, Bérénice Houinsou-Houssou, Chantal Septier, et al.. In vitro digestion of two age-tailored dairy products in the aging gastrointestinal tract. <https://www.icef14.com/en/committees/6>. The 14. International Congress on Engineering and Food (ICEF 2023), Jun 2023, Nantes, France. hal-04142497

HAL Id: hal-04142497

<https://hal.inrae.fr/hal-04142497>

Submitted on 27 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NoDerivatives 4.0 International License

In vitro digestion of age-tailored dairy products in the ageing gastrointestinal tract

A. Lavoisier^a, M. Morzel^a, C. Sounouvou^b, B. Houinsou-Houssou^b, C. Septier^b, C. Tournier^b, G. Feron^b, D. Dupont^a

^a UMR STLO, INRAE, Institut Agro, Rennes, France
^b UMR CSGA, INRAE, Dijon, France

Introduction

Context

- For older adults (> 65 y.) insufficient protein intake can lead to sarcopenia, characterized by the loss of muscle mass, strength, and function.
- To avoid this condition healthy older adults need to increase the amount of high-quality proteins in their diet (at least 1g protein/kg body weight/day).
- Milk proteins are interesting to promote muscle health, and particularly whey proteins which are rich in leucine.
- Ageing leads to changes in the functionality of the digestive tract but the impact of ageing on the intake, digestion, and absorption of nutrients is still unclear.

Objective

Investigate the digestion of 2 high-protein (10 % w/w) dairy products, in different *in vitro* conditions, to compare the kinetics of proteolysis in the gastric and intestinal phase between "young" and older adults.

High-protein products

- WBD** = fermented dairy product formulated with a ratio of whey proteins to caseins of 80 to 20% (as opposed to milk),
- Skyr** = fermented dairy product containing mainly caseins.

Static *in vitro* digestion¹

Oral phase

1: 1 food: SSF dilution according to dry matter
pH = 7.0, no chewing, no amylase

Gastric phase


	Young adults	Older adults
pH:	3.0	3.7
Duration:	2 h	3 h
Pepsin:	2000 U ml ⁻¹	1200 U ml ⁻¹
Gastric lipase:	60 U ml ⁻¹	36 U ml ⁻¹

Intestinal phase

	Young adults	Older adults
[Ca ²⁺]:	0.6 mM	1 mM
pH:	7.0	7.0
Duration:	2 h	2 h
Pancreatin:	100 U ml ⁻¹	80 U ml ⁻¹
Bile salts:	10 mM	6.7 mM


Results

Rheological properties




WBD had rheological properties comparable to a stirred yogurt and was considered suitable for older adults in terms of texture.

Sensory analysis²



No relationships were observed between oral physiology (posterior functional units, salivary flow, and saliva viscosity) and comfort.

Young vs. older adults' digestion of the dairy products



Gastric phase:

- Decrease in pepsin concentration (~30%), and increase in pH (from 3 to 3.7) between models resulted in a reduction in protein hydrolysis
- Caseins were rapidly digested by pepsin even at pH 3.7
- Proteolysis depended on the composition & structure of the product

Intestinal phase: no significant differences in proteolysis were measured at the end of the digestion between models.

Conclusions

- The whey protein-based dairy product developed in this study is suitable for older adults.
- The digestion conditions used (young vs. older adult) influenced significantly the kinetics and extent of proteolysis in the gastric phase but not in the intestinal phase.
- A clinical study comparing the effect of WBD or Skyr on older adults' postprandial muscle synthesis is underway.

Acknowledgments: This work was funded by the EAT4AGE project that received funding from the French National Research Agency (ANR), subvention number: ANR-20-HDHL-0002-01, under the umbrella of the European Joint Programming Initiative "A Healthy Diet for a Healthy Life" (JPI HDHL) and of the ERA-NET Cofund ERA-HDHL (GA N° 696295 of the EU Horizon 2020 Research and Innovation Programme).