

# Ecophysiological modelling of plant-nematode interactions to understand plant tolerance

Joseph Penlap<sup>1,2</sup>, Suzanne Touzeau<sup>1,2</sup>, Frédéric Gognard<sup>1</sup> and Valentina Baldazzi<sup>1,2</sup>

<sup>1</sup>Université Côte d'Azur, INRIA, INRAE, CNRS, Sorbonne Université, BIOCORE, Sophia Antipolis, France

<sup>2</sup>Université Côte d'Azur, INRAE, CNRS, ISA, Sophia Antipolis, France | \*joseph.penlap@inria.fr

## 1. Context

**Root-knot nematodes (RKN), *Meloidogyne* spp.**

- small soil worms, obligate root endoparasites
  - clonal reproduction
  - ubiquitous polyphagous pest
  - 14% of global crop losses worldwide [1]
- [1] Djian-Caporalino, *EPPO Bulletin*, 2012




**Symptoms**

- wilting and root deformation (galls)
- stunted growth
- reduced water and nutrient uptake
- hijacking of plant resources (carbon)

## 2. Research question

**Strong variability** in plant response to RKN parasitism among species & cultivars





**Which mechanisms underlie plant tolerance?**

**Approach**

- experimental data (tomato, cucurbit, pepper) with and without RKN
- model coupling plant ecophysiology and pest population dynamics

## 4. Model calibration

**Experimental tomato data (ongoing)**



## 3. Integrated plant-pest model



**Pest**

- RKN stages:
  - eggs
  - free-living larvae J2
  - within-root larvae
  - mature females
- RKN demography


**Plant**

- Plant compartments: fruit, shoot, root
- Vegetative and reproductive phases
- Plant resources: carbon and water
- Resource uptake and transport

**Plant-RKN interactions**



**Plant model**



$$\begin{cases}
 \text{Root} & \begin{cases} \frac{dW_r}{dt} = \underbrace{G_r(C_r)W_r}_{\text{Growth}} - \underbrace{\gamma_r W_r}_{\text{Mortality}} - \underbrace{\epsilon\beta J_2 W_r}_{\text{Infected roots}} \\ \frac{dC_r}{dt} = \underbrace{T_r(W_r)}_{\text{Transport}} - \underbrace{G_r(C_r)W_r}_{\text{Growth}} - \underbrace{r_m W_r}_{\text{Respiration}} - \underbrace{C_{rh} C_r}_{\text{Rhizodeposition}} - \underbrace{\gamma F}_{\text{RKN feeding}} - \underbrace{k\epsilon\beta J_2 W_r}_{\text{Gall formation}} \end{cases} \\
 \text{RKN} & \begin{cases} \frac{dj}{dt} = \underbrace{\Omega(C_r)\beta J_2 W_r}_{\text{RKN entry}} - \underbrace{\eta J}_{\text{Maturation}} - \underbrace{(\mu_j + \mu_r)J}_{\text{Mortality}} \\ \frac{dF}{dt} = \underbrace{\theta(C_r)\eta J}_{\text{Maturation}} - \underbrace{(\mu_F + \mu_r)F}_{\text{Mortality}} \end{cases}
 \end{cases}$$

## 5. Perspectives

- Identify **key physiological and architectural traits** underlying plant tolerance to guide the selection of new tolerant cultivars
- **Long-term dynamics:** effect of plant tolerance, cultural practices (rotations, etc.) and abiotic conditions on soil infestation and crop damages [2]

[2] Nilusmas et al., *Evolutionary Applications*, 2020