



HAL
open science

Homeobox Transcription Factor HbxA Influences Expression of over One Thousand Genes in the Model Fungus *Aspergillus nidulans*

S.S. Pandit, J. Zheng, Y. Yi, Sophie Lorber, Olivier Puel, S. Dhingra, E.A. Espeso, A.M Calvo

► **To cite this version:**

S.S. Pandit, J. Zheng, Y. Yi, Sophie Lorber, Olivier Puel, et al.. Homeobox Transcription Factor HbxA Influences Expression of over One Thousand Genes in the Model Fungus *Aspergillus nidulans*. 2023. hal-04148836

HAL Id: hal-04148836

<https://hal.inrae.fr/hal-04148836>

Preprint submitted on 3 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Homeobox Transcription Factor HbxA Influences Expression of over One Thousand Genes**
2 **in the Model Fungus *Aspergillus nidulans***

3 **Pandit S.S.¹, Zheng J.², Yi Y.², Lorber S.³, Puel O.³, Dhingra S.⁴, Espeso E.A.⁵, Calvo A.M.¹.**

4

5 ¹ Department of Biological Sciences, Northern Illinois University, DeKalb, Illinois, USA

6 ² Nebraska Food for Health Center, Department of Food Science and Technology, University of
7 Nebraska, Lincoln, Nebraska, USA.

8 ³ Toxalim (Research Centre in Food Toxicology), Université de Toulouse, INRAE, ENVT, INP-
9 Purpan, UPS, Toulouse, France

10 ⁴ Department of Biological Sciences and Eukaryotic Pathogen Innovation Center, Clemson
11 University, Clemson, South Carolina, USA

12 ⁵ Department of Cellular and Molecular Biology, Centro de Investigaciones Biológicas Margarita
13 Salas (C.S.I.C.), Madrid, Spain

14

15 **ABSTRACT**

16 In fungi, conserved homeobox-domain (HD) proteins are transcriptional regulators governing
17 development. In *Aspergillus* species, several HD transcription factor genes have been identified,
18 among them, *hbxA/hbxI*. For instance, in the opportunistic human pathogen *Aspergillus*
19 *fumigatus*, *hbxA* is involved in conidial production and germination, as well as virulence and

20 secondary metabolism (SM), including production of fumigaclavines, fumiquinazolines, and
21 chaetominine. In the agriculturally important fungus *Aspergillus flavus*, disruption of *hbx1*
22 results in fluffy aconidial colonies unable to produce sclerotia. *hbx1* also regulates production of
23 aflatoxins, cyclopiazonic acid and aflatrem. Furthermore, transcriptome studies revealed that
24 *hbx1* has a broad effect on the *A. flavus* genome, including numerous genes involved in SM.
25 These studies underline the importance of the HbxA/Hbx1 regulator, not only in developmental
26 processes but also in the biosynthesis of a broad number of fungal natural products, including
27 potential medical drugs and mycotoxins. To gain further insight into the regulatory scope of
28 HbxA in *Aspergilli*, we studied its role in the model fungus *Aspergillus nidulans*. Our present
29 study of the *A. nidulans hbxA*-dependent transcriptome revealed that more than one thousand
30 genes are differentially expressed when this regulator was not transcribed at wild-type levels,
31 among them numerous transcription factors, including those involved in development as well as
32 in SM regulation. Furthermore, our metabolomics analyses revealed that production of several
33 secondary metabolites, some of them associated with *A. nidulans hbxA*-dependent gene clusters,
34 was also altered in deletion and overexpression *hbxA* strains compared to the wild type,
35 including synthesis of nidulanins A, B and D, versicolorin A, sterigmatocystin, austinol,
36 dehydroaustinol, and three unknown novel compounds.

37 **INTRODUCTION**

38 Developmental studies of the model filamentous fungus *Aspergillus nidulans* have provided
39 broad valuable insight into the genetic regulatory mechanisms of morphogenesis in fungi (1–3).
40 *A. nidulans* efficiently disseminates by asexual reproduction, forming specialized structures
41 called conidiophores, which bares large numbers of air-borne conidia. Activation of
42 conidiogenesis is mediated by several transcription factor genes, including *flb* genes, such as

43 *flbB*, *flbC*, *flbD*, and *flbE*, which activate the central regulatory pathway comprised of *brlA*, *abaA*
44 and *wetA* (4). This model organism is also able to reproduce sexually by producing cleistothecia,
45 fruiting bodies containing meiospores called ascospores. Cleistothecia form by aggregation of
46 vegetative mycelia, surrounded by nursing Hülle cells. This results in the formation of
47 cleistothecial primordia, which later mature into melanized cleistothecia (5,6). Several genes are
48 involved in the regulation of these processes, including *nsdD*, *medA*, *phoA*, *stuA*, *lsdA* and *tubB*
49 (7–12).

50 Other developmental regulators include Homeobox-domain transcription factors (HD-TFs).
51 These are global regulators governing developmental processes in many eukaryotic organisms
52 (13–15). The HD contains approximately 66 conserved amino acid that bind to the promoter of
53 genes governing development and other cellular processes in fungi, plants, and animals. In
54 general, fungi possess 6–12 HD-TF genes in their genome (13,14,16,17). The first reported HD-
55 TF gene is *pah1* in *Podospora anserina*, where it controls microconidiation as well as mycelial
56 branching (18). Loss-of-function of seven HD-TF genes in this fungus revealed their role in
57 sexual development (19). Another study showed that several HD-TFs in the rice
58 pathogen *Magnaporthe oryzae* are necessary for proper hyphal growth, asexual development,
59 and appressorium formation (20,21). In three species of *Fusarium*, loss-of-function of
60 the *htf1* homeobox gene leads to alteration of phialides during conidiophore formation,
61 accompanied by a drastic reduction in conidial production (22). In the fungus *Botrytis cinerea*,
62 the BcHOX8 gene has been shown to regulate growth, conidiation, and virulence in different
63 host plants (16). Also, lack of the *GRF10* HD-TF gene in the human pathogen *Candida albicans*
64 resulted in a decrease in growth, defects in chlamydospore morphology, alterations in biofilm
65 production, and a reduction of virulence (23).

66 HD-TFs are also key regulators in species of the genus *Aspergillus*. In the agriculturally relevant
67 fungus *Aspergillus flavus*, deletion of eight HD-FT genes revealed that *hbx1* in particular, was
68 required for normal vegetative growth and production of conidia and sclerotia. The regulation of
69 morphological development as well as regulation of SM, are often genetically linked (24–26).
70 Interestingly, in this case also, the production of secondary metabolites, including mycotoxins
71 (aflatoxins, cyclopiazonic acid and aflatrem), was under the regulation of *hbx1* (17).
72 Furthermore, study of the *hbx1*-dependent transcriptome indicated its importance in
73 morphological development and in regulation of secondary metabolite production (27).
74 Remarkably, the gene category corresponding to SM was the most affected by *hbx1*.
75 Additionally, in our previous study of the *hbx1* homolog in the opportunistic human pathogen
76 *Aspergillus fumigatus*, *hbxA*, showed that this gene is necessary for proper spore formation,
77 regulating expression of *brlA*, *flbB*, *flbD* and *fluG* (28). The *hbxA* gene also influenced
78 germination rate and virulence in a neutropenic mouse model. Interestingly, as in the case of *A.*
79 *flavus*, *A. fumigatus hbxA* affected production of various secondary metabolites, including
80 fumigaclavines, fumiquinazolines, compounds that accumulate in asexual structures, whose
81 production is linked to *brlA* expression (29–32), and chaetominine, an alkaloid compound that is
82 being tested to combat leukemia cells(20). Both *A. flavus* and *A. fumigatus* studies indicate that
83 HbxA/Hbx1 is a global regulator of SM in these fungi, in addition to its role in morphogenesis.
84 HbxA also affects *A. nidulans* conidiation (33,34) in a similar manner as that in *A. flavus* and *A.*
85 *fumigatus*(17,27,28). To gain further inside into the regulatory scope of *hbxA* in the genus
86 *Aspergillus*, in the present study, we characterized its role in the model fungus *A. nidulans* by
87 transcriptome and metabolomics approaches. Our findings indicate that more than one thousand
88 genes were differentially expressed in the absence of this regulator or when it was over-

89 expressed, as compared to the wild type. These include several transcription factor genes,
90 including those involved in development and SM production. Our study revealed that numerous
91 secondary metabolites gene clusters are *hbxA*-dependent in *A. nidulans*. Furthermore, our
92 analyses also indicated that *A. nidulans* metabolome is affected by *hbxA*, including production of
93 some unknown novel compounds.

94 **MATERIALS and METHODS**

95 *Phylogenetic Analysis*

96 Deduced amino acid sequences of HbxA homologs were obtained from FUNGIDB
97 (<https://fungidb.org/fungidb/>) website. BLASTp was performed against the protein sequence
98 database (pdb). Percentage (%) similarity was found using Pairwise sequence alignment using
99 EMBOSS Needle (ebi.ac.uk/Tools/psa/emboss_needle/). The phylogenetic tree was constructed using
100 MEGA v6.0 and the Maximum Likelihood model with bootstrap value of 1000.

101 *Strains used and culture conditions*

102 The *A. nidulans* strains used in this study are listed in Table 1. Strains were grown on glucose
103 minimal medium (GMM) (35) with appropriate supplements for their respective auxotrophic
104 markers (35). For solid medium, agar (15 g/L) was added. Strains were stored as 30% glycerol
105 stocks at -80°C.

106 **Table 1: Strains used in this study**

Strain Name	Pertinent Genotype	Source
RMJP1.49	<i>pyrG89; argB2; Δnku::argB; pyroA4</i>	(36)

TSSP38.1	<i>pyrG89; argB2; Δnku::argB; pyroA4; ΔhbxA::pyrG; pyroA</i>	This Study
TSSP40.1	<i>pyrG89; argB2; ΔnkuA::argB; pyroA4; ΔhbxA::pyrG, hbxA::pyroA</i>	This Study
TSSP34.1	<i>pyrG89; argB2; ΔnkuA::argB; pyroA4; gpdA(p)::ΔhbxA::trpC(t)::pyrG; pyroA</i>	This Study
TRV50.2	Wild type	(37)

107

108 ***Generation of the *hbxA* deletion strain ($\Delta hbxA$)***

109 The DNA cassette employed to obtain the deletion *hbxA* strain (TSSP38.1) was generated by
110 fusion polymerase chain reaction (PCR) through a previously described method (38). All
111 primers used in this study are listed in Table 2. The 1.5 kb 5` UTR region of the *hbxA* locus was
112 PCR amplified using P#2154/SD3 and P#2155 primers from genomic DNA of the *A. nidulans*
113 FGSC4 wild-type strain. Similarly, the 1.1 kb 3` UTR of *hbxA* was amplified using P#2156 and
114 P#2157 primers also from genomic DNA. The 1.9 kb *A. fumigatus pyrG* selectable marker was
115 amplified from plasmid p1439 (39) using P#2158 and P#2159 primers. The 5` and 3` UTR
116 fragments were fused to the selectable *pyrG* marker using P#2160/SD9 and P#2161 primers. The
117 resultant fusion product was transformed into RMJP1.49 strain using a polyethylene glycol
118 mediated protocol as described previously (38). Transformants were confirmed by diagnostic
119 PCR using P#2154 and P#963 primers. The selected deletion *hbxA* strain was then transformed
120 with a DNA fragment containing the *A. nidulans pyroA* gene, PCR amplified with primers
121 P#1042 and P#1045 from genomic DNA, resulting in strain TSSP38.1.

122

123 **Table 2: Primers used in this study**

Name	Sequence (5' → 3')
P#2154/SD3	CCCGCTGATGTATGGTGAGGC
P#2155	TGGTGTAGGATGCGATGCGG
P#2156	CATCTCCTCCTTCAACACCAGGG
P#2157	GGTCTGAGGTCTTGCCGTTTCC
P#2158	CCGCATCGCATCCTACACCAACCGGTCGCCTCAAACAATGCTCT
P#2159	CCCTGGTGTGAAGGAGGAGATGGTCTGAGAGGAGGCACTGATGCG
P#2160/SD9	CGCTCCCTTGAAACTCCGAGAG
P#2161	CACAGTAGGCACGAATGGCGTT
SD1	ACCGGTCGCCTCAAACAATGCTCT
SD2	GTCTGAGAGGAGGCACTGATGCG
SD4	GCGTTTTATTCTTGTTGACATGGGGTCCCTTAGCCGAAATTGGTGGG
SD5	CCCATGTCAACAAGAATAAAACGC
SD6	CCGAGTGGAGATGTGGAGT
SD7	ACTCCACATCTCCACTCGGGCCCATATCTTCCGTAGCAGTC
SD8	AGAGCATTGTTTGAGGCGACCGGTGACGGAGAGCTGAGAGTCCTAG
SD10	CAGAGCACCGCCGTGGTATTG
P#2962	GTCTCGTAGGTCTCTTGACGACCG
P#2238	AAAAAAGGCGCGCCATGAATTATATCCATCATCCATACCCTTTCGCTG

P#2239 AAAAAAAAAAGCGGCCGCTTAGCCGAAATTGGTGGGGGTC
P#1042 GCCGAAAAGGACCACGAATACCCGC
P#1045 CACCGCCAACGGAGACAATCAAGCC
P#963 GAGCAGCGTAGATGCCTCGAC
P#2093 GACCTAATACAGCCCCTACAACGACC
P#2218 GCGGCCGCTTAGCCGAAATTGGTGGGGGTC

124

125

126 ***Generation of the *hbxA* complementation strain (*hbxA-com*)***

127 The complementation strain (TSSP40.1) was generated by re-introducing the wild-
128 type *hbxA* allele into the $\Delta hbxA$ strain at the same locus. The complementation cassette was
129 generated as follows: first, a DNA fragment containing the *hbxA* coding region and a 3.7 kb
130 5' UTR was PCR amplified using P#2154/SD3 and SD4, and the *trpC* terminator fragment was
131 amplified with primers SD5 and SD6 using *A. nidulans* genomic DNA as a template. The *A.*
132 *fumigatus pyroA* gene (Afub_055620) was amplified from genomic DNA using primers SD7 and
133 SD8. *A. fumigatus pyrG* was amplified from plasmid p1439 (38) using primers SD1 and SD2.
134 All four PCR fragments were fused together using primers P#2160/SD9 and SD10 in a single
135 reaction using Prime Star DNA polymerase (Clontech, USA). The resulting fusion product was
136 then transformed into the *hbxA* deletion strain (TSSP38.1) using methods previously described
137 (38). Fungal transformants were confirmed using diagnostic PCR with primers P#2154 and
138 P#2962.

139 ***Generation of the *hbxA* overexpression strain (OE*hbxA*)***

140 To generate the over-expression *hbxA* strain (TSSP34.1), the coding region of *hbxA* was first
141 amplified from *A. nidulans* genomic DNA using P#2238 and P#2239 primers. The resulting PCR
142 product was digested with *AscI* and *NotI* and ligated to pTRS2 plasmid, previously digested with
143 the same enzymes. pTRS2 contains the *gpdA* promoter, *gpdA*_(p), and *trpC* terminator, *trpC*_(t).
144 The resulting plasmid, pSSP34.1, was transformed into the *A. nidulans* RJMP1.49 strain, and
145 transformants were screened by PCR using P#2093 and P#2218 primers. The selected
146 overexpression *hbxA* strain was then transformed with a DNA fragment containing the *A.*
147 *nidulans pyroA* gene, PCR amplified with primers P#1042 and P#1045 from genomic DNA,
148 resulting in strain TSSP34.1.

149 ***Transcriptome analysis***

150 *RNA purification and sequencing*

151 Plates containing 25 mL of solid GMM with the appropriate supplements were top-agar
152 inoculated with 5 mL of medium containing $\sim 5 \times 10^6$ spores/mL of wild-type (WT) control,
153 $\Delta hbxA$, *hbxA*-com or OE*hbxA* (Table 1). Cultures were incubated in the dark at 37°C. After 72 h
154 of incubation, mycelia were collected, frozen in liquid nitrogen, and lyophilized. Total RNA was
155 extracted from mycelia using an RNeasy Plant Mini Kit (Qiagen, Germantown, Maryland, USA)
156 following the manufacturer's protocol. RNA was further purified using Dynabeads mRNA
157 Purification Kit (Thermo Fisher Scientific Inc., Massachusetts, USA). RNA quality was assessed
158 using an Agilent Bioanalyzer. Sequencing was performed as a HiSeq 2000 single read 1x100bp
159 lane. The experiment was carried out with 3 biological replicates.

160 *Read mapping, decontamination and Read count*

161 The RNA reads were trimmed by trim_galore (40) with the default parameter. Kraken2 (41) was
162 run on trimmed reads to check the contamination. Then, reads were mapped to reference genome
163 downloaded from FungiDB (*Aspergillus nidulans* FGSC4)(42) . Unmapped reads were removed
164 to get clean reads. The clean reads were then repaired to pair-end reads with BBTools (43).
165 These final clean pair-end reads were remapped to reference genome again using hisat2 ((44).
166 Mapped reads in SAM format were sorted by coordinates with samtools (45) to obtain the BAM
167 format mapped reads. Then read count and TPM (Transcripts Per Kilobase Million) were
168 calculated by running StringTie (46) and python script. The parameters were set not to infer new
169 transcripts with the reference gene annotation file (also downloaded from FungiDB).

170 *Differentially expressed coding genes (DEGs)*

171 The read counts table was used as input for DEseq2 (47). This package was used to determine
172 DEGs by comparing read counts between two strains. Significant up regulated genes were
173 determined with $-\log_{10}$ q-value ≤ 2 and \log_2 fold change ≥ 2 , while significant down
174 regulated genes were defined with $-\log_{10}$ q-value ≤ 2 and \log_2 fold change ≤ -2 . Control vs.
175 OE*hbxA* and Control vs. Δ *hbxA*. python script was developed to convert gene id between
176 FungiDB and FungiFun2 so that the webserver of FungiFun2 can be used to perform FunCat
177 term annotation and enrichment of DEGs for Control vs. OE*hbxA* and Control vs. Δ *hbxA*(48).
178 Heat maps of TPM (transcript per million) values of DEGs of secondary metabolism clusters
179 were calculated by averaging all TPM values of all replicates.

180 Evaluation of differentially expressed ortholog genes in *A. nidulans* and *A. flavus* was carried out
181 by using the MCL algorithm in combination with all-versus-all protein BLAST search, similar to
182 a method previously described (49). Proteins with BLAST hits were filtered with the following

183 parameters: 1, query and subject coverage is greater than 60%. 2, e-value is less than 1^{-5} . 3, the
184 percent of identity is greater than 60%. And then, the filtered hits were fed into OrthoMCL with
185 an inflation parameter of 2 to generate orthogroups between these two species.

186 To analyze changes in the expression of genes in secondary metabolite biosynthetic gene clusters
187 (SMGs), 67 SMGs were extracted (50). SMGs expression related figures were plotted with
188 python seaborn package. In addition, expression of 521 transcript factors (TFs) was also
189 analyzed. The list of TFs and their function annotations were derived from a previous report
190 (51).

191 *Metabolomics*

192 *Thin- Layer Chromatography*

193 Wild-type control, $\Delta hbxA$, $hbxA$ -com, and OE $hbxA$ were top-agar inoculated with 5 mL of
194 medium containing $\sim 5 \times 10^6$ spores/mL on solid GMM and grown at 37°C for 3 days. Three 16-
195 mm diameter cores per plate were collected and extracted with chloroform. Overnight dried
196 extracts were resuspended in 200 μ L chloroform. Sample were separated using thin-layer
197 chromatography (TLC) as previously described (28,52) on silica gel plates using benzene and
198 glacial acetic acid [95:5(v/v)] as solvent system. Aluminum chloride (15% in ethanol) was then
199 sprayed, and plates were baked for 10 min at 80 °C. Bands were visualized under UV light (375
200 nm). Sterigmatocystin (ST) standard was purchased from Sigma-Aldrich (St. Louis, MO, USA).

201 *Analysis of secondary metabolites by liquid chromatography combined with mass spectrometry*
202 *(LC-MS)*

203 Chloroform extracted samples were also analyzed by LC-MS. Samples were resuspended in 500
204 μL of acetonitrile/water (50:50, v/v), shaken vigorously for 30 s and then treated with a sonicator
205 (Bransonic 221 Ultrasonic bath, Roucaire, Les Ulis, France) for 2 h. A volume of 250 μL of pure
206 ACN was added to each sample, followed by vigorous shaking (30s) and centrifugation (pulse).
207 Secondary metabolites analysis was performed using Acquity ArcSystem HPLC (Waters, Saint-
208 Quentin-en-Yvelines, France) combined with an LTQ Orbitrap XL high-resolution mass
209 spectrometer (Thermo Fisher Scientific, Les Ulis, France). A volume of 10 μL of the suspension
210 was injected into a reversed-phase 150 mm \times 2.0 mm, Luna[®] 5 μm C18 column (Phenomenex,
211 Torrance, CA, U.S.A.). Water acidified with 0.1% formic acid was used as phase A and 100%
212 acetonitrile was used as phase B with the following elution gradient: 0 min 20% B, 30 min 50%
213 B, from 35 to 45 min 90% B, from 50 to 60 min 20% B at 30 °C at a flow rate of 0.2 mL min⁻¹.
214 HRMS acquisitions were achieved with electrospray ionization (ESI) in positive and negative
215 modes, as previously reported (28). MS/MS spectra were obtained with CID mode at low
216 resolution and collision energy of 35%.

217 *Statistical analysis*

218 Statistical analysis was applied to analyze all quantitative data in this study utilizing analysis of
219 variance (ANOVA) in conjunction with a Tukey multiple-comparison test using a *p* value of
220 <0.05 for samples that are determined to be significantly different.

221

222 **RESULTS**

223 *HbxA is conserved in numerous fungal species*

224 Our phylogenetic analysis confirmed that the *hbxA* deduced amino acid sequence corresponds to
225 a transcription factor containing a homeodomain. HbxA homologs are present in other
226 *Aspergillus* species, including *A. flavus* (17,27), *A. fumigatus* (28), *Aspergillus niger* and
227 *Aspergillus terreus* (Fig 1, Table 3), as well as in species of other fungal genera, such as
228 *Alternaria alternata*, *Arthrotrrys flagrans*, *Ascospaera apis*, *Blastomyces dermatitidis*,
229 *Histoplasma capsulatum*, *Microsporium canis*, *Penicillioopsis zonata*, *Penicillium rubens*,
230 *Talaromyces marneffeii* and *Trichophyton tonsurans* (Fig 1, Table 3). Of the sequences analyzed,
231 *A. niger* HbxA was the closest homolog to *A. nidulans* HbxA, with 56.40% identity and 68.4%
232 sequence similarity.

233 **Table 3: Phylogenetic analysis of *A. nidulans* HbxA and homologs in other fungal species. HbxA**
234 **homologs were retrieved from FUNGIDB website and BLASTp was performed against protein**
235 **sequence database. % similarity was found utilizing Pairwise sequence alignment using *A. nidulans***
236 **HbxA as search query against each protein of interest using EMBOSS Needle.**

Species (sorted)	Identity% (Needle)	Similarity% global Pairwise alignment
<i>Aspergillus niger</i>	56.4	68.4
<i>Aspergillus flavus</i>	52.9	65.2
<i>Penicillioopsis zonata</i>	44.5	54.3
<i>Penicillium rubens</i>	43.7	56.9
<i>Talaromyces marneffeii</i>	37.1	49.3
<i>Blastomyces dermatitidis</i>	36.9	49.9
<i>Histoplasma capsulatum</i>	36.9	49.5
<i>Aspergillus terreus</i>	35.5	46.4
<i>Microsporium canis</i>	34.0	43.7

<i>Trichophyton tonsurans</i>	32.3	44.9
<i>Ascosphaera apis</i>	30.3	41.8
<i>Arthrobotrys flagrans</i>	21.5	34.3
<i>Alternaria alternata</i>	20	31.5

237

238 **Fig 1: Phylogenetic analysis of *Aspergillus nidulans* HbxA.** The phylogenetic tree was
239 constructed using MEGA v6.0 and the Maximum Likelihood model with bootstrap value
240 of 1000 (<http://megasoftware.net/>).

241

242

243 **Fig 2: Multiple sequence alignment of *A. nidulans* HbxA with other fungal**
244 **homologs.**

245 The HbxA deduced amino acid sequences were aligned using
246 clustalOmega(<https://www.ebi.ac.uk/Tools/msa/clustalo/>). Data was visualized with
247 boxshade using ENDScript server (<https://escript.ibcp.fr/ESPrpt/cgi-bin/ESPrpt.cgi>
248)(53)<https://doi.org/10.1093/nar/gku316>

249

250 ***hbxA* is required for normal development in *A. nidulans***

251 To determine the regulatory scope of *hbxA* in *A. nidulans*, three strains were generated, a
252 deletion strain, $\Delta hbxA$, a complementation strain, *hbxA-com*, and an over-expression strain,
253 OE*hbxA* (Fig 3). Deletion, complementation and overexpression strains were confirmed by
254 diagnostic PCR, yielding the expected 3.01 kb PCR product for $\Delta hbxA$, a 3.96 kb DNA fragment

255 for *hbxA-com* and a 3.16 kb DNA fragment for OE*hbxA*. Our results confirmed that absence of
256 *hbxA* results in a drastic reduction of conidiation (Fig 4), as previously shown(33,34).

257 **Fig 3: Generation of *A. nidulans hbxA* deletion, complementation and**
258 **overexpression strains.** Confirmation of the deletion ($\Delta hbxA$), complementation (*hbxA-*
259 *com*) and overexpression (OE*hbxA*) by diagnostic PCR. (A) The diagram shows
260 replacement of *hbxA* with the marker gene *pyrG* by a double cross-over event. Primers
261 P#2154/SD3 and P#963 were used for the diagnostic PCR, obtaining the predicted 3.01
262 kb product. (B) Schematic representation showing reintroduction of the wild-type *hbxA*
263 allele at the *hbxA* locus in the deletion strain TSSP38.1. PCR with primers P#2154/SD3
264 and P#2962 confirmed the reintroduction of *hbxA* in the selected deletion strain; the
265 expected 3.96 kb product was obtained. (C) Linear diagram of *hbxA* overexpression
266 plasmid pSSP34.1. The overexpression transformant was confirmed by PCR with primers
267 2093 and 2218, which yielded the predicted 3.16 kb product.

268

269 **Fig 4. *hbxA* is required for normal conidiation in *A. nidulans*.** Cultures of wild type,
270 deletion, complementation and overexpression *hbxA* strains, top-agar inoculated on
271 GMM and incubated for 7 days in the dark at 37°C.

272

273 *hbxA* regulates secondary metabolism

274 Our TLC analysis indicated that deletion of *hbxA* reduces sterigmatocystin (ST) production in *A.*
275 *nidulans* by approximately 50 % when compared with levels in the wild-type strain (Fig 5).

276 Importantly, overexpression of *hbxA* completely blocked ST production. Additionally, synthesis

277 of other metabolites was also affected by deletion or forced overexpression of *hbxA* compared to
278 the control strain. The absence of metabolites was particularly notable in the OE*hbxA* strain
279 extracts. These results suggested that the regulatory role of *hbxA* is broader than originally
280 expected, controlling not only developmental processes but also acting as a global regulator of
281 secondary metabolism.

282 **Fig 5. Effect of *hbxA* on the production of ST and other secondary metabolites in *A.***
283 ***nidulans*.** Wild type, deletion, complementation and overexpression *hbxA* strains were
284 top-agar inoculated on glucose minimum medium (GMM) and incubated for 3 days in the
285 dark. (A) Extracts were analyzed by TLC. Black arrows indicate ST standard. The
286 experiment was carried out with three replicates. (B) Densitometry of TLC analysis of ST
287 levels. The densitometry was performed using the
288 <http://biochemlabsolutions.com/GelQuantNET.html> website. Error bars represent the standard
289 error. Columns of different letters represent values that are statistically different *p* value
290 of <0.05

291

292 ***hbxA*-dependent transcriptome in *A. nidulans***

293 *More than one thousand genes are regulated by hbxA in A. nidulans*

294 RNA-sequencing analysis revealed that of the predicted 11286 genes present in *A. nidulans*
295 genome (54), 552 were downregulated, and 195 were upregulated in the Δ *hbxA* strain compared
296 with the wild-type control strain (Table 4, Fig 6). Over-expression of *hbxA* resulted in an even
297 more pronounced effect on the *A. nidulans* transcriptome, where 1044 genes were
298 downregulated, and 424 genes were upregulated in the OE*hbxA* strain in comparison to the wild

299 type. In strong contrast, the comparison of the complementation strain and wild type showed that
300 the two strains present very similar expression patterns. Expression of 618 genes in the *A.*
301 *nidulans* genome was altered by either deletion or overexpression of *hbxA*, many of them
302 presenting the same expression pattern of upregulation or downregulation when *hbxA* was either
303 deleted or overexpressed (Fig 6).

304 **Fig 6. Number of DEGs in *A. nidulans* when expression of *hbxA* is altered by *hbxA***
305 **deletion or overexpression. (A)** Number of significantly upregulated (purple) and
306 significantly downregulated (orange) DEGs estimated by DeSeq2. **(B)** Volcano plot of
307 log₂ fold change vs. -log₁₀ q-value of all the genes in $\Delta hbxA$, and OE*hbxA* vs. control.
308 Significantly upregulated genes are shown as red dots, significant down regulated genes
309 are shown as blue dot and other genes are shown as black. The x-axis represents the log₂
310 of the fold change determined by DeSeq2. The y-axis is the log₁₀ of the adjusted p-value
311 from DeSeq2. The cut off log₁₀ fold change value to determine the upregulated
312 expression is greater than 2 while -2 is for down regulated expression. The -log₁₀ q-value
313 cutoff was set to 2 to determine the significant expression or not. **(C-D)** Venn Diagrams
314 showing the overlap of DEGs in $\Delta hbxA$ and OE*hbxA* **(C)**, and the overlap of upregulated
315 **(D)** and downregulated DEGs **(E)** in $\Delta hbxA$ and OE*hbxA*. Venn Diagrams were
316 constructed using [https://bioinformatics.psb.ugent.be/cgi-](https://bioinformatics.psb.ugent.be/cgi-bin/liste/Venn/calculate_venn.html)
317 [bin/liste/Venn/calculate_venn.html](https://bioinformatics.psb.ugent.be/cgi-bin/liste/Venn/calculate_venn.html) website.

318 *Comparison of hbxA/hbx1 DEGs in A. nidulans and A. flavus*

319 The comparison of the current *A. nidulans hbxA*-dependent transcriptome study with the
320 previous *A. flavus hbx1* results (27) is shown in Fig 7. Only a small percentage of homologs were
321 differentially expressed in the absence of *hbxA* and *hbx1* in *A. nidulans* and *A. flavus*,

322 respectively, with respect to the corresponding wild types. Most of the DEGs in *A. nidulans* are
323 not DEGs in the *A. flavus* study.

324 **Fig 7: Comparison of orthologous genes affected by deletion of *hbxA* in *A. nidulans***
325 **and *A. flavus*.** Both upregulated orthologous genes were colored in red. Both
326 downregulated orthologous genes were colored in blue. No expression changed
327 orthologous genes are colored in grey. Two orthologous genes having different regulation
328 status are colored in purple. The significantly regulated genes were defined as $|\log_2 \text{fold}$
329 $\text{change}| \leq 2$ and $q\text{-value} \leq 0.05$.

330

331 *Expression of numerous TF genes is hbxA-dependent in A. nidulans*

332 Based on our analysis, 74 out of 521 TFs genes in *A. nidulans* (51) were regulated by *hbxA* under
333 the culture conditions assayed (Table 5). Some of these differentially expressed TF genes also
334 presented the same expression pattern of upregulation or downregulation when *hbxA* was either
335 deleted or overexpressed (Fig 8).

336 **Fig 8. Number of transcription factor (TF) genes controlled by *hbxA* in *A. nidulans*.**

337 **(A)** Venn Diagram showing the overlap of differentially expressed TF genes in $\Delta hbxA$
338 and OE*hbxA*. **(B-C)** Venn Diagrams showing the overlap of upregulated **(B)** or
339 downregulated **(C)** TF genes in $\Delta hbxA$ and OE*hbxA*. Venn Diagrams were constructed
340 using https://bioinformatics.psb.ugent.be/cgi-bin/liste/Venn/calculate_venn.html website.

341

342 Our results indicated that overexpression of *hbxA* caused upregulation of developmental
343 regulators, including genes of the central developmental pathway, *brlA*, *abaA* (55–58), *fluffy*
344 genes *flbC* and *flbD* (59), and another HD-TF gene, *hbxB*, that regulates asexual and sexual
345 development in *A. nidulans*(34). In addition, the developmental regulatory gene *zcfA* (60) was
346 also upregulated by *hbxA* overexpression. Some of the upregulated TFs genes in OE*hbxA* are
347 involved in both governing development as well as SM, such as the master transcription factor
348 *mtfA* (37,61,62), *urda*, (63), *sclB* (64), *osaA* (65), and *velB* (66). Other upregulated *hbxA*-
349 dependent TF-DEGs annotated to be putatively involved in SM regulation include AN8391 and
350 AN6788. Other upregulated TF genes have an important role in primary metabolism, such as
351 *glcD*, which has a putative role in protein dimerization and activation of *areB*, (67), *galR*, which
352 is known to regulate the D-galactose catabolic pathway (68) and *creA* repressor of carbon
353 catabolite (69). Other upregulated TF genes were *rfeC*, whose ortholog in *Saccharomyces*
354 *cerevisiae* promotes *FLO11* expression (70), the *mcnB* fork-head like transcription factor (71), as
355 well as expression of some other uncharacterized putative transcription factors genes (Table 5).

356 Overexpression of *hbxA* in *A. nidulans* caused downregulation of other developmental genes
357 such as *flpA*, with a role in sexual development (72), *mat1*, involved in activation of the alpha-
358 domain mating-type protein (73). Overexpression of *hbxA* also caused downregulation of *metZ*,
359 a transcription factor involved in the regulation of sulfur metabolism (74). TF genes AN8377,
360 AN8645, AN3385 and AN8918 predicted to be involved in SM, are also downregulated in this
361 strain (Table S2).

362 Interestingly, deletion of *hbxA* also resulted in an increase in the expression of *brlA*, *abaA*, and
363 *urda*, as in the case of OE*hbxA*. It also increased the expression of *tah-3*, which is involved in
364 conidiophore development and tolerance for harsh plasma environment (75) (Table 5). Deletion

365 of *hbxA* also upregulated *veA* (Table 4). The *veA* gene product, VeA, which contains a NF- κ -B
366 like DNA-binding domain (76), is well known as a global regulator that interacts with at least nine
367 other proteins, LlmF, VapA, VipA, VipC, VelB, MpkB, FphA, LreB and LaeA (77), governing
368 several signaling pathways and consequently multiple cellular processes, including development
369 and SM (25).

370 Absence of *hbxA* in *A. nidulans* downregulated the expression of various transcription factors,
371 including the gene encoding the alpha-domain mating-type protein, *mat1*(73), as in
372 overexpression of *hbxA*. Deletion of *hbxA* also showed downregulation of *metZ*, involved in
373 methionine biosynthesis (78) the nitrogen-dependent *mdpE*, which regulates production of a
374 secondary metabolite called monodictyphenone (79). The putative SM TF gene AN4933 is
375 downregulated, and AN3385, AN8645 and AN8918 are also downregulated in deletion *hbxA*, as
376 in OE*hbxA*.

377

378 *hbxA* affects the expression of genes in SM gene clusters and biosynthesis of natural products
379 in *A. nidulans*

380 Our TLC analysis revealed that both deletion and overexpression of *hbxA* negatively affect ST
381 production (Fig 5) as well as the production of other secondary metabolites. Furthermore,
382 FunCat enrichment analysis revealed that differentially regulated genes in the Δ *hbxA* versus wild
383 type and OE*veA* versus wild type comparisons have significant functional overlap (Fig 9). DEGs
384 genes are dramatically enriched for secondary metabolism-related processes for both; most of
385 those genes are downregulated when *hbxA* is either deleted or overexpressed, particularly in the

386 latter. Other categories showing enrichment include disease, virulence, and defense; virulence
387 disease factors; C-compound and carbohydrate metabolism; and detoxification.

388 **Fig 9. FunCat enrichment of significant DEGs found in (A) $\Delta hbxA$ and (B) OE*hbxA***
389 **vs. control.** The $-\log_{10}$ of the q-value of DEGs in each term is proportional to the length
390 of the bars. FunCat annotations and q-value is determined by FungiFun2 webserver.
391 Downregulated genes are to the left of the origin and up regulated genes to the right.

392

393 To gain further understanding of the effect of *hbxA* on SM in *A. nidulans*, as part of our
394 transcriptome analysis, we identified DEGs in SM gene clusters and analyzed concomitant
395 production of secondary metabolites by a metabolomics approach. Our study revealed that
396 production of nidulaninA, nidulanin B and nidulanin D are *hbxA*- dependent (Fig 10A-C). Both,
397 deletion and overexpression of *hbxA*, completely inhibited the production of these compounds. In
398 addition, the Heatmap shown in Fig 10D indicates downregulation of some of the genes in the
399 nidulanin cluster (80), including the NRPS coding gene, *nlsA*, in both $\Delta hbxA$ and OE*hbxA*. This
400 reduction in *nlsA* expression was particularly notable in the latter.

401 **Fig 10: *hbxA* regulates the production of nidulanins in *A. nidulans*.** Wild-type (WT),
402 deletion ($\Delta hbxA$), complementation (*hbxA-com*) and overexpression (OE*hbxA*) strains
403 were top-agar inoculated on solid glucose minimum medium (GMM) at 37°C for 72 h,
404 when samples were collected, extracted with chloroform and analyzed by LC-HRMS in
405 positive mode (A-C) Quantification of nidulanin A (m/z 604.34943), B (m/z 620.34404),
406 D (m/z 536.28659) respectively. (D) Heat map of TPM values of nidulanin cluster
407 (DEGs) expression in *A. nidulans* $\Delta hbxA$ and OE*hbxA* with respect to wild type strain on

408 a log scale found in Inglis et al.(50). The TPM value of each gene was calculated by
409 averaging all the TPM values of all replicates.

410

411 LC-MS analysis of ST confirmed the TLC results, indicating that production of this mycotoxin
412 was reduced in $\Delta hbxA$ and absent in the overexpression strains (Fig 11). Unexpectedly, the
413 Heatmap in Fig 11B shows that most of the ST genes were not downregulated in the deletion
414 strain with respect to the wild type. However, most of the genes in this cluster were
415 downregulated in the overexpression strain, excluding the structural genes *stcK*, *stcJ*, *stcF* and
416 *stcC*, and the regulator, *aflR*.

417 **Fig 11. *hbxA* regulates the production of ST in *A. nidulans*.** Wild type (WT), deletion
418 ($\Delta hbxA$), complementation (*hbxA-com*) and overexpression (OE*hbxA*) strains were top-
419 agar inoculated on solid glucose minimum medium (GMM) at 37°C for 72 h, when
420 samples were collected, extracted with chloroform and analyzed by LC-HRMS in
421 positive mode. (A) Quantification of ST (*m/z* 325.07014). (B) Heat map of TPM values
422 of ST cluster (DEGs) expression in *A. nidulans* $\Delta hbxA$ and OE*hbxA* with respect to wild
423 type strain on a log scale found in Inglis et al. (50). The TPM value of each gene was
424 calculated by averaging all the TPM values of all replicates.

425

426 In addition, both $\Delta hbxA$ and OE*hbxA* strains were unable to synthesize the meroterpenoids
427 austinol and dehydroaustinol under conditions conducive to their production in the wild type (Fig
428 12). The genes involved in the synthesis of these two compounds are grouped in two clusters, A
429 and B (81). Our transcriptome analysis revealed that most of the genes in these two clusters are

430 downregulated in the *hbxA* deletion and also in the overexpression strains compared to the
431 control (Fig 12C and D). For example, genes *ausA-D* are down regulated in both $\Delta hbxA$ and
432 OE*hbxA* in gene cluster A. In cluster B, genes *ausE-G* and *ausM* are also downregulated in
433 $\Delta hbxA$ and OE*hbxA*. Additionally, expression of *ausH*, *ausL* and *ausN* is reduced in OE*hbxA*
434 with respect to the wild type.

435 **Fig 12. *hbxA* regulates the production of austinol and dehydroaustinol in *A.***
436 ***nidulans*.** Wild type (WT), deletion ($\Delta hbxA$), complementation (*hbxA-com*), and
437 overexpression (OE*hbxA*) strains were top-agar inoculated on solid glucose minimum
438 medium (GMM) at 37°C for 72 h, when samples were collected, extracted with
439 chloroform and analyzed by LC-HRMS in positive mode. Quantification of (A) austinol
440 (*m/z* 459.20059) and (B) dehydroaustinol (*m/z* 457.18524) compounds by full MS spectra
441 resolution of 60,000 with a range of mass-to-charge ratio (*m/z*) set to 50 to 800. (C &
442 D) Heatmap of TPM values of austinol cluster (DEGs) expression in *A. nidulans* $\Delta hbxA$
443 and OE*hbxA* with respect to wild type strain on a log scale found in Inglis et al. (50). The
444 TPM value of each gene was calculated by averaging all the TPM values of all replicates.

445

446 Our metabolomics study also indicated that the production of three novel, unknown secondary
447 metabolites was altered when *hbxA* was not expressed at wild-type levels. Two of these
448 compounds (*m/z* 423 and *m/z* 518 observed in negative mode) were absent in the *hbxA* deletion
449 strain and also in the overexpression strain (Fig 13). The third novel compound (*m/z* 489 in
450 negative mode) was produced at remarkably high levels in the *hbxA* deletion strain compared to
451 those in the wild type (Fig 13B).

452 **Fig 13: *hbxA* regulates the production of novel uncharacterized metabolites in *A.***
453 ***nidulans*.** Wild type (WT), deletion ($\Delta hbxA$), complementation (*hbxA-com*), and
454 overexpression (*OEhbxA*) strains were top-agar inoculated on solid glucose minimum
455 medium (GMM) at 37 °C for 72 h, when samples were collected, extracted with
456 chloroform and analyzed by LC-HRMS in negative mode. (A-C) Quantification of novel
457 uncharacterized metabolites with *m/z* of 423.18012, 489.18082, and 518.16482,
458 respectively.

459

460 DISCUSSION

461

462 HD-TFs have been shown to govern development in eukaryotes (13–15), including fungi
463 (13,14,16–21). Previous reports, together with the present study, indicate that these regulators are
464 conserved across different fungal genera. (Zheng et al., 2012; Ghosh et al., 2015). In *A. flavus*,
465 *hbxA*, an ortholog of *hbxA*, is also required for developmental processes, regulating genes in the
466 conidiation central pathway, such as *brlA* and *wetA* (17,27) as well as *flbA*, *flbC*, *flbD*, *flbE*, *fluG*
467 and *mat1-1*(27). In *A. fumigatus*, *hbxA* promotes *brlA*, *abaA* and *wetA*, as well as *flbB*, *flbD* and
468 *fluG* expression(28). Similarly, *hbxA* regulates conidiation in *A. nidulans*(33,34); our
469 transcriptome study showed that *hbxA* not only regulates *brlA*, as shown in (34), but also *abaA*,
470 *flbC* and *flbD*. These studies support that the *hbxA*-dependent regulatory mechanism of
471 conidiation is at least in part conserved in these three *Aspergillus* species and possibly in other
472 species of this genus.

473 Interestingly, our results revealed a broader regulatory scope for *hbxA* in *A. nidulans*, with more
474 than one thousand DEGs when *hbxA* was deleted or overexpressed in this model organism,
475 including numerous transcription factor genes. This was also the case for *A. flavus hbx1*(27).
476 However, most of the DEGs in *A. nidulans* are not DEGs in *A. flavus*; only a small percentage of
477 homologs were DEGs in the *hbxA* and *hbx1* mutants with respect to the controls. This suggests
478 that although the conservation of some of the regulatory mechanisms controlling conidiation
479 appears conserved, a great part of its regulatory input is specialized in different fungal species.

480 Some of the TF genes involved in governing development that were found *hbxA*-dependent also
481 control secondary metabolism in *A. nidulans*, for example, *mtfA* (37,61,62), *urdA* (63), *sclB* (64),
482 *osaA* (65) and *velB* (66). Furthermore, FunCat functional enrichment analysis showed that the
483 category of secondary metabolism-related processes was, by far, the most enriched in *A.*
484 *nidulans*. Our study showed that in *A. nidulans*, numerous genes in SM gene clusters were
485 regulated by *hbxA*. The secondary metabolism category was also enriched in *A. flavus*
486 (27)However, the wide variation of biosynthetic gene clusters across fungal species, even in
487 those phylogenetically close (82) could explain that although the major functional category is the
488 same in both species, namely SM, the percentage of differentially expressed homologs is low.

489 For example, *A. flavus hbx1* regulates genes in the aflatoxin, cyclopiazonic acid, aflatrem,
490 asparasone, piperazine, and aflavarin gene clusters(27), while in *A. nidulans*, our study shows
491 that *hbxA* controls genes in other gene clusters such as those responsible for the synthesis of
492 nidulanins A, B and D, austinol and dehydroaustinol. *A. nidulans* HbxA also control genes in the
493 ST gene cluster, which is partially conserved with that of aflatoxin in *A. flavus*. The regulatory
494 pattern was similar; absence of both *hbxA* and *hbx1* resulted in a reduction of toxin production
495 (17,27). In *A. flavus* deletion of *hbx1* downregulated *aflR* and other genes in the aflatoxin gene

496 cluster. However, this was not the case in *A. nidulans*, suggesting that the lower levels of ST in
497 the deletion strain, verified by both TLC and LC-MS, could be due to other factor(s). Our study
498 showed *veA* expression is *hbxA*-dependent. VeA is a global regulator that orchestrates numerous
499 biological processes in fungi (25,26), such as development and SM. VeA has been shown to
500 regulate the production of aflatoxins in *A. parasiticus* (83). It is possible that *hbxA*, in a *veA*-
501 dependent manner, could also influence compartmentalization of ST production in *A. nidulans*.
502 This reduction in ST in the deletion strain, contrast with a previous report (34) where an increase
503 in ST was described. It is possible that different experimental conditions in both studies could
504 have resulted in different outcomes. Nevertheless, the most striking result is the effect of *hbxA*
505 overexpression on ST biosynthesis as well as on the production of other metabolites. The
506 complete elimination of ST production by *hbxA* overexpression was, in this case, accompanied
507 by the downregulation of genes in the ST gene cluster. However, this downregulation of ST
508 genes was, as in the case of the deletion strain, not mediated by changes in *aflR* expression.

509 Our study revealed that *hbxA* regulates key genes in the nidulanin gene cluster and,
510 consequently, affects the production of the cyclic tetrapeptides nidulinins A, B and D. These
511 compound are found in *Aspergillus* and *Penicillium* species. The function of nidulanins is not yet
512 known. As in the case of ST, both deletion or overexpression of *hbxA* resulted in reduction or
513 elimination of nidulinins A, B and D production, suggesting that, as in the case of VeA, certain
514 balanced stoichiometry with respect to other regulatory factors could be needed for proper
515 function, perhaps also interacting with other regulatory proteins. One of the genes downregulated
516 in both deletion *hbxA* and overexpression *hbxA* strains is *nlsA*, encoding a non-ribosomal peptide
517 synthase necessary for the synthesis of nidulanin. This enzyme has been shown to also be

518 involved in the synthesis of fungisporin (84), which presents antibacterial activity (85), however
519 fungisporin was not detected in our study under the conditions tested.

520 LC-MS indicated that *hbxA* also controls austinol and dehydroaustinol production. These are two
521 meroterpenoids produced from polyketide and terpenoid precursors. Both austinol and
522 dehydroaustinol have been shown to inhibit the neuraminidase enzyme, suggesting a potential
523 for the development of new antiviral drugs (86). Austinol also showed antibacterial activity (87).
524 Alteration of wild-type *hbxA* transcription by deletion or forced overexpression also resulted in a
525 lack of production of these compounds, further supporting the possibility of a necessary
526 stoichiometry with other regulatory partners. Two separate gene clusters, A and B (81,88), are
527 required for the synthesis of these compounds. Both deletion and overexpression of *hbxA* showed
528 profound changes in the expression profile of both gene clusters, with numerous downregulated
529 structural genes, including the polyketide synthase gene *ausA*. The prenyltransferase gene *ausN*
530 was also downregulated in the overexpression strain.

531 In addition, our metabolomics analysis indicated that *A. nidulans hbxA* also controls the
532 production of three unknown novel compounds. Synthesis of two of these metabolites (m/z &423
533 and m/z 528) did not occur in the absence of *hbxA* or when this gene was overexpressed, while
534 the third novel compound (m/z 489) was produced at strikingly high levels in the *hbxA* deletion
535 strain. The identity, the association with MS gene clusters, or bioactive properties of these
536 compounds are still known and will be the subject of future studies.

537 Regarding additional roles of *hbxA* in *A. nidulans*, besides those in development and SM, our
538 FunCat functional enrichment analysis also indicated a possible role in primary metabolism, with
539 enrichment in the carbon-compound and carbohydrate metabolism category, particularly in the

540 *hbxA* overexpression strain. Upregulation of the carbon catabolite repressor TF gene *creA* (69)
541 was observed in this strain. *creA* is also under *hbx1* regulation in *A. flavus*(27). Other *A. nidulans*
542 *hbxA*-dependent regulatory genes involved in primary metabolism were, for example, *galR*,
543 which regulates the D-galactose catabolic pathway (68), and *glcD*, which has a putative role in
544 protein dimerization with and activation of *areB*, involved in nitrogen metabolism (67). Other
545 enriched categories were detoxification, virulence and disease factors and defense, suggesting its
546 possible involvement in pathogenesis. This agrees with the fact that the *hbxA* homolog in *A.*
547 *fumigatus* was shown to affect virulence in *A. fumigatus* (28)

548 In conclusion, we have shown that the regulatory TF gene *hbxA* governs the expression of
549 hundreds of genes in *A. nidulans*, modulating not only developmental genes, but also multiple
550 regulatory pathways. Consequently, *hbxA* governs different important aspects of this fungus'
551 biology, including a remarkable role in SM, regulating expression of several SM gene clusters
552 and natural product biosynthesis, including some novel compounds. Additionally, genes
553 associated with other cellular processes such as primary metabolisms, as well as defense and
554 virulence, are also influenced by *hbxA*. Interestingly, a functional conservation exists between
555 *hbxA* homologs in other *Aspergillus* species and possibly in other fungi.

556

557 **ACKNOWLEDGEMENTS**

558 This study was funded by Northern Illinois University

559

560 **REFERENCES**

- 561 1. Caesar LK, Kelleher NL, Keller NP. In the fungus where it happens: History and future propelling
562 *Aspergillus nidulans* as the archetype of natural products research. *Fungal Genet Biol.* 2020 Nov
563 1;144:103477.
- 564 2. Oiartzabal-Arano E, Perez-de-Nanclares-Arregi E, Espeso EA, Etxebeste O. Apical control of
565 conidiation in *Aspergillus nidulans*. *Curr Genet.* 2016 May 1;62(2):371–7.
- 566 3. Yu JH. Regulation of Development in *Aspergillus nidulans* and *Aspergillus fumigatus*. *Mycobiology.*
567 2010 Dec 31;38(4):229–37.
- 568 4. Etxebeste O, Garzia A, Espeso EA, Ugalde U. *Aspergillus nidulans* asexual development: making the
569 most of cellular modules. *Trends Microbiol.* 2010 Dec;18(12):569–76.
- 570 5. Aramayo R, Timberlake W e. The *Aspergillus nidulans* *yA* gene is regulated by *abaA*. *EMBO J.* 1993
571 May;12(5):2039–48.
- 572 6. Hermann TE, Kurtz MB, Champe SP. Laccase localized in hulle cells and cleistothecial primordia of
573 *Aspergillus nidulans*. *J Bacteriol.* 1983 May;154(2):955–64.
- 574 7. Bussink HJ, Osmani SA. A cyclin-dependent kinase family member (PHOA) is required to link
575 developmental fate to environmental conditions in *Aspergillus nidulans* | *The EMBO Journal* . 1998.
576 Available from: <https://www.embopress.org/doi/full/10.1093/emboj/17.14.3990>
- 577 8. Clutterbuck AJ. A mutational analysis of conidial development in *Aspergillus nidulans*. *Genetics.*
578 1969 Oct 1;63(2):317–27.
- 579 9. Han KH, Han KY, Yu JH, Chae KS, Jahng KY, Han DM. The *nsdD* gene encodes a putative GATA-type
580 transcription factor necessary for sexual development of *Aspergillus nidulans*. *Mol Microbiol.* 2001
581 Jul 1;41(2):299–309.

- 582 10. Kirk KE, Morris NR. The tubB alpha-tubulin gene is essential for sexual development in *Aspergillus*
583 *nidulans*. *Genes Dev.* 1991 Nov 1;5(11):2014–23.
- 584 11. Lee, Kim, S., Kim, S. J., Han, D. M., Jahng, K. Y., Chae, K. S. The *IsdA* gene is necessary for sexual
585 development inhibition by a salt in *Aspergillus nidulans* | SpringerLink. 2001. Available from:
586 <https://link.springer.com/article/10.1007/s002940100206>
- 587 12. Miller KY, Toennis TM, Adams TH, Miller BL. Isolation and transcriptional characterization of a
588 morphological modifier: the *Aspergillus nidulans* stunted (*stuA*) gene. *Mol Gen Genet MGG.* 1991
589 Jun 1;227(2):285–92.
- 590 13. Holland PWH. Evolution of homeobox genes. *Wiley Interdiscip Rev Dev Biol.* 2013 Feb;2(1):31–45.
- 591 14. Mukherjee K, Brocchieri L, Bürglin TR. A comprehensive classification and evolutionary analysis of
592 plant homeobox genes. *Mol Biol Evol.* 2009 Dec;26(12):2775–94.
- 593 15. Svingen T, Tonissen KF. Hox transcription factors and their elusive mammalian gene targets.
594 *Heredity.* 2006 Aug;97(2):88–96.
- 595 16. Antal Z, Rasclé C, Cimerman A, Viaud M, Billon-Grand G, Choquer M, et al. The Homeobox *BcHOX8*
596 Gene in *Botrytis Cinerea* Regulates Vegetative Growth and Morphology. *PLoS ONE* [Internet]. 2012
597 Oct 25 [cited 2018 Nov 29];7(10). Available from:
598 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3485016/>
- 599 17. Cary, Harris-Coward P, Scharfenstein L, Mack B, Chang PK, Wei Q, et al. The *Aspergillus flavus*
600 Homeobox Gene, *hbx1*, Is Required for Development and Aflatoxin Production. *Toxins* [Internet].
601 2017 Oct 12;9(10). Available from: <http://www.mdpi.com/2072-6651/9/10/315>

- 602 18. Arnaise S, Zickler D, Poisier C, Debuchy R. *pah1*: a homeobox gene involved in hyphal morphology
603 and microconidiogenesis in the filamentous ascomycete *Podospora anserina*. *Mol Microbiol*. 2001
604 Jan;39(1):54–64.
- 605 19. Coppin E, Berteaux-Lecellier V, Bidard F, Brun S, Ruprich-Robert G, Espagne E, et al. Systematic
606 deletion of homeobox genes in *Podospora anserina* uncovers their roles in shaping the fruiting
607 body. *PLoS One*. 2012;7(5):e37488.
- 608 20. Kim S, Park SY, Kim KS, Rho HS, Chi MH, Choi J, et al. Homeobox Transcription Factors Are Required
609 for Conidiation and Appressorium Development in the Rice Blast Fungus *Magnaporthe oryzae*.
610 Copenhagen GP, editor. *PLoS Genet*. 2009 Dec 4;5(12):e1000757.
- 611 21. Liu W, Xie S, Zhao X, Chen X, Zheng W, Lu G, et al. A homeobox gene is essential for conidiogenesis
612 of the rice blast fungus *Magnaporthe oryzae*. *Mol Plant-Microbe Interact MPMI*. 2010
613 Apr;23(4):366–75.
- 614 22. Zheng W, Zhao X, Xie Q, Huang Q, Zhang C, Zhai H, et al. A Conserved Homeobox Transcription
615 Factor *Htf1* Is Required for Phialide Development and Conidiogenesis in *Fusarium* Species. Yu JH,
616 editor. *PLoS ONE*. 2012 Sep 21;7(9):e45432.
- 617 23. Ghosh AK, Wangsanut T, Fonzi WA, Rolfes RJ. The *GRF10* homeobox gene regulates filamentous
618 growth in the human fungal pathogen *Candida albicans*. *FEMS Yeast Res*. 2015 Dec;15(8):fov093.
- 619 24. Calvo AM, Wilson RA, Bok JW, Keller NP. Relationship between Secondary Metabolism and Fungal
620 Development. *Microbiol Mol Biol Rev*. 2002 Sep 1;66(3):447–59.
- 621 25. Calvo AM, Lohmar JM, Ibarra B, Satterlee T. 18 Velvet Regulation of Fungal Development. In:
622 Wendland J, editor. *Growth, Differentiation and Sexuality* [Internet]. Cham: Springer International

- 623 Publishing; 2016. p. 475–97. Available from: [http://link.springer.com/10.1007/978-3-319-25844-](http://link.springer.com/10.1007/978-3-319-25844-7_18)
624 [7_18](http://link.springer.com/10.1007/978-3-319-25844-7_18)
- 625 26. Calvo AM, Cary JW. Association of fungal secondary metabolism and sclerotial biology. *Front*
626 *Microbiol.* 2015;6:62.
- 627 27. Cary, Entwistle S, Satterlee T, Mack BM, Gilbert MK, Chang PK, et al. The Transcriptional Regulator
628 Hbx1 Affects the Expression of Thousands of Genes in the Aflatoxin-Producing Fungus *Aspergillus*
629 *flavus*. *G3 Bethesda Md.* 2019;9(1):167–78.
- 630 28. Satterlee T, Nepal B, Lorber S, Puel O, Calvo AM. The Transcriptional Regulator HbxA Governs
631 Development, Secondary Metabolism, and Virulence in *Aspergillus fumigatus*. *Appl Environ*
632 *Microbiol.* 2020 Jan 21;86(3). Available from: <https://aem.asm.org/content/86/3/e01779-19>
- 633 29. Calvo AM. The VeA regulatory system and its role in morphological and chemical development in
634 fungi. *Fungal Genet Biol.* 2008 Jul;45(7):1053–61.
- 635 30. Cary, John E. Linz, Deepak Bhatnagar. *Microbial Foodborne Diseases | Mechanisms of Pathogenesis*
636 *and Toxin Sy.* 2014. Available from:
637 [https://www.taylorfrancis.com/books/mono/10.1201/9781482278873/microbial-foodborne-](https://www.taylorfrancis.com/books/mono/10.1201/9781482278873/microbial-foodborne-diseases-jeffrey-cary-john-linz-deepak-bhatnagar)
638 [diseases-jeffrey-cary-john-linz-deepak-bhatnagar](https://www.taylorfrancis.com/books/mono/10.1201/9781482278873/microbial-foodborne-diseases-jeffrey-cary-john-linz-deepak-bhatnagar)
- 639 31. Han KH, Lee DB, Kim JH, Kim MS, Han KY, Kim WS, et al. Environmental factors affecting
640 development of *Aspergillus nidulans*. *J Microbiol.* 2003;41(1):34–40.
- 641 32. Kosalková K, García-Estrada C, Ullán RV, Godio RP, Feltrer R, Teijeira F, et al. The global regulator
642 LaeA controls penicillin biosynthesis, pigmentation and sporulation, but not roquefortine C
643 synthesis in *Penicillium chrysogenum*. *Biochimie.* 2009 Feb;91(2):214–25.

- 644 33. Pandit, S. S., Satterlee, T., Nepal, B., Lorber, S., Puel, O., Espeso, E. A., et al. The transcriptional
645 regulatory HbxA governs development and secondary metabolism in *Aspergillus nidulans* and
646 *Aspergillus fumigatus*. 30th Fungal Genetics Conference, March 12-17, 2019, at Asilomar
647 Conference Grounds in Pacific Grove, CA; 2019.
- 648 34. Son SH, Son YE, Cho HJ, Chen W, Lee MK, Kim LH, et al. Homeobox proteins are essential for fungal
649 differentiation and secondary metabolism in *Aspergillus nidulans*. *Sci Rep*. 2020 Dec;10(1):6094.
- 650 35. Käfer E. Meiotic and Mitotic Recombination in *Aspergillus* and Its Chromosomal Aberrations. In:
651 Caspari EW, editor. *Advances in Genetics*. Academic Press; 1977. p. 33–131. Available from:
652 <http://www.sciencedirect.com/science/article/pii/S006526600860245X>
- 653 36. Feng X, Ramamoorthy V, Pandit SS, Prieto A, Espeso EA, Calvo AM. *cpsA* regulates mycotoxin
654 production, morphogenesis and cell wall biosynthesis in the fungus *Aspergillus nidulans*. *Mol*
655 *Microbiol*. 2017 Jul 1;105(1):1–24.
- 656 37. Ramamoorthy V, Dhingra S, Kincaid A, Shantappa S, Feng X, Calvo AM. The Putative C2H2
657 Transcription Factor MtfA Is a Novel Regulator of Secondary Metabolism and Morphogenesis in
658 *Aspergillus nidulans*. *PLOS ONE*. 2013 Sep 16;8(9):e74122.
- 659 38. Szewczyk E, Nayak T, Oakley CE, Edgerton H, Xiong Y, Taheri-Talesh N, et al. Fusion PCR and gene
660 targeting in *Aspergillus nidulans*. *Nat Protoc*. 2006 Dec;1(6):3111–20.
- 661 39. Stinnett SM, Espeso EA, Cobeño L, Araújo-Bazán L, Calvo AM. *Aspergillus nidulans* VeA subcellular
662 localization is dependent on the importin β carrier and on light. *Mol Microbiol*. 2007 Jan;63(1):242–
663 55.
- 664 40. Krueger F. Trim Galore. 2022. Available from: <https://github.com/FelixKrueger/TrimGalore>

- 665 41. Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2. *Genome Biol.* 2019 Nov
666 28;20(1):257.
- 667 42. Basenko EY, Pulman JA, Shanmugasundram A, Harb OS, Crouch K, Starns D, et al. FungiDB: An
668 Integrated Bioinformatic Resource for Fungi and Oomycetes. *J Fungi Basel Switz.* 2018 Mar
669 20;4(1):39.
- 670 43. Bushnell B, Rood J, Singer E. BBMerge - Accurate paired shotgun read merging via overlap. *PloS One.*
671 2017;12(10):e0185056.
- 672 44. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment and genotyping
673 with HISAT2 and HISAT-genotype. *Nat Biotechnol.* 2019 Aug;37(8):907–15.
- 674 45. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map
675 format and SAMtools. *Bioinforma Oxf Engl.* 2009 Aug 15;25(16):2078–9.
- 676 46. Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL. StringTie enables improved
677 reconstruction of a transcriptome from RNA-seq reads. *Nat Biotechnol.* 2015 Mar;33(3):290–5.
- 678 47. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data
679 with DESeq2. *Genome Biol.* 2014;15(12):550.
- 680 48. Priebe S, Kreisel C, Horn F, Guthke R, Linde J. FungiFun2: a comprehensive online resource for
681 systematic analysis of gene lists from fungal species. *Bioinforma Oxf Engl.* 2015 Feb 1;31(3):445–6.
- 682 49. Lind AL, Wisecaver JH, Smith TD, Feng X, Calvo AM, Rokas A. Examining the Evolution of the
683 Regulatory Circuit Controlling Secondary Metabolism and Development in the Fungal Genus
684 *Aspergillus*. *PLOS Genet.* 2015 Mar 18;11(3):e1005096.

- 685 50. Inglis DO, Binkley J, Skrzypek MS, Arnaud MB, Cerqueira GC, Shah P, et al. Comprehensive
686 annotation of secondary metabolite biosynthetic genes and gene clusters of *Aspergillus nidulans*, *A.*
687 *fumigatus*, *A. niger* and *A. oryzae*. *BMC Microbiol.* 2013 Apr 26;13(1):91.
- 688 51. Etxebeste O. Transcription Factors in the Fungus *Aspergillus nidulans*: Markers of Genetic
689 Innovation, Network Rewiring and Conflict between Genomics and Transcriptomics. *J Fungi.* 2021
690 Jul 25;7(8):600.
- 691 52. Cleveland DW. Peptide mapping in one dimension by limited proteolysis of sodium dodecyl sulfate-
692 solubilized proteins. In: *Methods in Enzymology*. Academic Press; 1983. p. 222–9. (Biomembranes
693 Part J: Membrane Biogenesis: Assembly and Targeting (General Methods, Eukaryotes); vol. 96).
694 Available from: <https://www.sciencedirect.com/science/article/pii/S0076687983960202>
- 695 53. Robert X, Gouet P. Deciphering key features in protein structures with the new ENDscript server.
696 *Nucleic Acids Res.* 2014 Jul 1;42(W1):W320–4.
- 697 54. Arnaud MB, Cerqueira GC, Inglis DO, Skrzypek MS, Binkley J, Chibucos MC, et al. The *Aspergillus*
698 Genome Database (AspGD): recent developments in comprehensive multispecies curation,
699 comparative genomics and community resources. *Nucleic Acids Res.* 2012 Jan;40(Database
700 issue):D653–9.
- 701 55. Adams TH, Boylan MT, Timberlake WE. *brlA* is necessary and sufficient to direct conidiophore
702 development in *aspergillus nidulans*. *Cell.* 1988 Jul 29;54(3):353–62.
- 703 56. Andrianopoulos A, Timberlake WE. The *Aspergillus nidulans* *abaA* gene encodes a transcriptional
704 activator that acts as a genetic switch to control development. *Mol Cell Biol.* 1994 Apr;14(4):2503–
705 15.


- 706 57. Chang YC, Timberlake WE. Identification of *Aspergillus* brlA response elements (BREs) by genetic
707 selection in yeast. *Genetics*. 1993 Jan 1;133(1):29–38.
- 708 58. Prade RA, Timberlake WE. The *Aspergillus nidulans* brlA regulatory locus consists of overlapping
709 transcription units that are individually required for conidiophore development. *EMBO J*. 1993
710 Jun;12(6):2439–47.
- 711 59. Etxebeste O, Ni M, Garzia A, Kwon NJ, Fischer R, Yu JH, et al. Basic-Zipper-Type Transcription Factor
712 FlbB Controls Asexual Development in *Aspergillus nidulans*. *Eukaryot Cell*. 2008 Jan;7(1):38–48.
- 713 60. Son YE, Cho HJ, Lee MK, Park HS. Characterizing the role of Zn cluster family transcription factor ZcfA
714 in governing development in two *Aspergillus* species. *PLoS ONE*. 2020 Feb 4;15(2):e0228643.
- 715 61. Smith TD, Calvo AM. The mtfA transcription factor gene controls morphogenesis, gliotoxin
716 production, and virulence in the opportunistic human pathogen *Aspergillus fumigatus*. *Eukaryot*
717 *Cell*. 2014 Jun;13(6):766–75.
- 718 62. Zhuang Z, Lohmar JM, Satterlee T, Cary JW, Calvo AM. The Master Transcription Factor mtfA
719 Governs Aflatoxin Production, Morphological Development and Pathogenicity in the Fungus
720 *Aspergillus flavus*. *Toxins*. 2016 Jan 20;8(1):29.
- 721 63. Pandit SS, Lohmar JM, Ahmed S, Etxebeste O, Espeso EA, Calvo AM. UrdA Controls Secondary
722 Metabolite Production and the Balance between Asexual and Sexual Development in *Aspergillus*
723 *nidulans*. *Genes*. 2018 Nov 23;9(12). Available from:
724 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6316066/>
- 725 64. Thieme KG, Gerke J, Sasse C, Valerius O, Thieme S, Karimi R, et al. Velvet domain protein VosA
726 represses the zinc cluster transcription factor SclB regulatory network for *Aspergillus nidulans*

- 727 asexual development, oxidative stress response and secondary metabolism. PLOS Genet. 2018 Jul
728 25;14(7):e1007511.
- 729 65. Alkahyyat F, Ni M, Kim SC, Yu JH. The WOPR Domain Protein OsaA Orchestrates Development in
730 *Aspergillus nidulans*. PLOS ONE. 2015 Sep 11;10(9):e0137554.
- 731 66. Bayram O, Krappmann S, Ni M, Bok JW, Helmstaedt K, Valerius O, et al. VelB/VeA/LaeA complex
732 coordinates light signal with fungal development and secondary metabolism. Science. 2008 Jun
733 13;320(5882):1504–6.
- 734 67. Arst HN, Hondmann DHA, Visser J. A translocation activating the cryptic nitrogen regulation gene
735 *areB* inactivates a previously unidentified gene involved in glycerol utilisation in *Aspergillus nidulans*
736 | SpringerLink. 1990. Available from: <https://link.springer.com/article/10.1007/BF00315805>
- 737 68. Kowalczyk JE, Gruben BS, Battaglia E, Wiebenga A, Majoor E, de Vries RP. Genetic Interaction of
738 *Aspergillus nidulans galR*, *xlnR* and *araR* in Regulating D-Galactose and L-Arabinose Release and
739 Catabolism Gene Expression. PLoS ONE. 2015 Nov 18;10(11):e0143200.
- 740 69. Strauss J, Horvath HK, Abdallah BM, Kindermann J, Mach RL, Kubicek CP. The function of CreA, the
741 carbon catabolite repressor of *Aspergillus nidulans*, is regulated at the transcriptional and post-
742 transcriptional level. Mol Microbiol. 1999 Apr;32(1):169–78.
- 743 70. Askenazi M, Driggers EM, Holtzman DA, Norman TC, Iverson S, Zimmer DP, et al. Integrating
744 transcriptional and metabolite profiles to direct the engineering of lovastatin-producing fungal
745 strains. Nat Biotechnol. 2003 Feb;21(2):150–6.

- 746 71. Ukil L, Varadaraj A, Govindaraghavan M, Liu HL, Osmani SA. Copy Number Suppressors of the
747 *Aspergillus nidulans* nimA1 Mitotic Kinase Display Distinctive and Highly Dynamic Cell Cycle-
748 Regulated Locations. *Eukaryot Cell*. 2008 Dec;7(12):2087–99.
- 749 72. Dyer PS, O’Gorman CM. Sexual development and cryptic sexuality in fungi: insights from *Aspergillus*
750 species. *FEMS Microbiol Rev*. 2012 Jan 1;36(1):165–92.
- 751 73. Pyrzak W, Miller KY, Miller BL. Mating Type Protein Mat1-2 from Asexual *Aspergillus fumigatus*
752 Drives Sexual Reproduction in Fertile *Aspergillus nidulans*. *Eukaryot Cell*. 2008 Jun;7(6):1029–40.
- 753 74. Piłsyk S, Natorff R, Sieńko M, Skoneczny M, Paszewski A, Brzywczy J. The *Aspergillus nidulans* metZ
754 gene encodes a transcription factor involved in regulation of sulfur metabolism in this fungus and
755 other Eurotiales. *Curr Genet*. 2015 May;61(2):115–25.
- 756 75. Xiong Y, Wu VW, Lubbe A, Qin L, Deng S, Kennedy M, et al. A fungal transcription factor essential for
757 starch degradation affects integration of carbon and nitrogen metabolism. *PLOS Genet*. 2017 May
758 3;13(5):e1006737.
- 759 76. Ahmed YL, Gerke J, Park HS, Bayram Ö, Neumann P, Ni M, et al. The Velvet Family of Fungal
760 Regulators Contains a DNA-Binding Domain Structurally Similar to NF-κB. Stock AM, editor. *PLoS*
761 *Biol*. 2013 Dec 31;11(12):e1001750.
- 762 77. Röhrig J, Yu Z, Chae KS, Kim JH, Han KH, Fischer R. The *Aspergillus nidulans* Velvet-interacting
763 protein, VipA, is involved in light-stimulated heme biosynthesis. *Mol Microbiol*. 2017;105(6):825–38.
- 764 78. Alaminos, Ramos. The methionine biosynthetic pathway from homoserine in *Pseudomonas putida*
765 involves the metW, metX, metZ, metH and metE gene products - PubMed. 2001. Available from:
766 <https://pubmed.ncbi.nlm.nih.gov/11479715/>

- 767 79. Chiang YM, Szewczyk E, Davidson AD, Entwistle R, Keller NP, Wang CCC, et al. Characterization of
768 the *Aspergillus nidulans* Monodictyphenone Gene Cluster. *Appl Env Microbiol.* 2010 Apr
769 1;76(7):2067–74.
- 770 80. Andersen MR, Nielsen JB, Klitgaard A, Petersen LM, Zachariasen M, Hansen TJ, et al. Accurate
771 prediction of secondary metabolite gene clusters in filamentous fungi. *Proc Natl Acad Sci U S A.*
772 2013 Jan 2;110(1):E99–107.
- 773 81. Lo HC, Entwistle R, Guo CJ, Ahuja M, Szewczyk E, Hung JH, et al. Two separate gene clusters encode
774 the biosynthetic pathway for the meroterpenoids austinol and dehydroaustinol in *Aspergillus*
775 *nidulans*. *J Am Chem Soc.* 2012 Mar 14;134(10):4709–20.
- 776 82. Rokas A, Mead ME, Steenwyk JL, Raja HA, Oberlies NH. Biosynthetic gene clusters and the evolution
777 of fungal chemodiversity. *Nat Prod Rep.* 2020 Jul 1;37(7):868–78.
- 778 83. Chanda A, Roze LV, Kang S, Artymovich KA, Hicks GR, Raikhel NV, et al. A key role for vesicles in
779 fungal secondary metabolism. *Proc Natl Acad Sci.* 2009 Nov 17;106(46):19533–8.
- 780 84. Ali, Marco I Ries, Peter P Lankhorst, Rob A M van der Hoeven, Olaf L Schouten, Marek Noga, et al. A
781 non-canonical NRPS is involved in the synthesis of fungisporin and related hydrophobic cyclic
782 tetrapeptides in *Penicillium chrysogenum* - PubMed. 2014. Available from:
783 <https://pubmed.ncbi.nlm.nih.gov/24887561/>
- 784 85. Himaja, M., Elizabeth, L., Moonjit, D., Asif, K. Synthesis and antimicrobial evaluation of N-
785 methylated analog of fungisporin. *Univ J Pharm.* 2015;4:10–4.

- 786 86. Gozari M, Alborz M, El-Seedi HR, Jassbi AR. Chemistry, biosynthesis and biological activity of
787 terpenoids and meroterpenoids in bacteria and fungi isolated from different marine habitats. *Eur J*
788 *Med Chem.* 2021 Jan 15;210:112957.
- 789 87. Fuloria NK, Raheja RK, Shah KH, Oza, M.J. Biological activities of meroterpenoids isolated from
790 different sources. *Frontiers Media SA*; 2022. 174 p.
- 791 88. Matsuda Y, Awakawa T, Mori T, Abe I. Unusual chemistries in fungal meroterpenoid biosynthesis.
792 *Curr Opin Chem Biol.* 2016 Apr 1;31:1–7.
- 793



0.20

	1	10	20	30	40
<i>Aspergillus nidulans</i>	MNYIHP	YFAGHPS	VPMEQH	LAVDT
<i>Aspergillus niger</i>	MNYLHP	YAFAGHAA	VPMEQP	VAFDPTMAHPSM
<i>Aspergillus flavus</i>	MNYLHP	YAYAGHAA	VPMEQP	IAYDPTMAHPSM
<i>Aspergillus oryzae</i>	MNYLHP	YAYAGHAA	VPMEQP	IAYDPTMAHPSM
<i>Penicillium zonata</i>	MSYLHS	YYPSSHPA	VPDQ	LVEFPAMAHSSM
<i>Penicillium rubens</i>	MSYIHP	WNYQGHQG	IPMDQH	MAYDPSMVP
<i>Talaromyces marneffei</i>	MNYLHP	YYPGVHAG	IHLDP	GLVHP.....AIA
<i>Blastomyces dermatitidis</i>	MSYLHP	YFPGGHA	IPVDQP	VGYGVP
<i>Histoplasma capsulatum</i>	MSYLHP	YFPGGHA	IPVDQP	FDYRVP
<i>Microsporium canis</i>	MNYLQAP	YQY	SPMDQP	MGYGMP
<i>Trichophyton tonsurans</i>	MNYLQTP	FQY	AMPDQP	MGYGMP
<i>Ascosphaera apis</i>	MNYLHP	YGYGAPT	LPIDQP	MPYGVP
<i>Arthrotrichum flagrans</i>	MEV	LSYDNKGS	MPSONP	NVPR	SDVHVTQGPS
<i>Alternaria alternata</i>

	50	60	70	80	90	100
<i>Aspergillus nidulans</i>	YAQP	PFD	DMVD	YYHQ	..P	MDYEEYA
<i>Aspergillus niger</i>	YPHP	PFD	DMVD	FYHQ	..P	MDYEEYA
<i>Aspergillus flavus</i>	YPHP	PFD	DMVD	FYHQ	..P	MDYEEYA
<i>Aspergillus oryzae</i>	YPHP	PFD	DMVD	FYHQ	..P	MDYEEYA
<i>Penicillium zonata</i>	YPHP	PFD	EMVD	FYPP	..P	MDYEEYA
<i>Penicillium rubens</i>	YPHP	PFD	EMVD	FYPP	..P	MDYEEYA
<i>Talaromyces marneffei</i>	LTRPAY	ELAD	YYTHMP	LM	DEEY	ENLSRPRLTKE
<i>Blastomyces dermatitidis</i>	LPNAP	IDLA	YYHQ	AAALE	DFFEY	ENLSRPRLTKE
<i>Histoplasma capsulatum</i>	LPNAP	IDLA	YYHQ	AAALE	DFFEY	ENLSRPRLTKE
<i>Microsporium canis</i>	VPYHQ	NDIHGF	YTT	GAI	EYEEY	ENLSRPRLTKE
<i>Trichophyton tonsurans</i>	IPYHQ	NDIHGF	YTT	GAI	EYEEY	ENLSRPRLTKE
<i>Ascosphaera apis</i>	GHHV	PVELANS	FYQAVT	LD	ED	ESLSRPRLTKE
<i>Arthrotrichum flagrans</i>	GNHF	PV	IPRFT	TP	PVSGP	NEESSVK
<i>Alternaria alternata</i>	MDSS	GGST	TD	AP	SAPKQ

	110	120	130	140	150
<i>Aspergillus nidulans</i>	QOTHLSLPRVAN	WFQNRRAKAK	QOKRQEE	YERMQ	KAKAEAE
<i>Aspergillus niger</i>	AQTNLSLPRVAN	WFQNRRAKAK	QOKRQEE	FEKMQ	KAKAEAE
<i>Aspergillus flavus</i>	AQTNLSLPRVAN	WFQNRRAKAK	QOKRQEE	FERMQ	KAKTEAAE
<i>Aspergillus oryzae</i>	AQTNLSLPRVAN	WFQNRRAKAK	QOKRQEE	FERMQ	KAKTEAAE
<i>Penicillium zonata</i>	AQTNLSLPRVAN	WFQNRRAKAK	QOKRQEE	FERMQ	KAKAQAE
<i>Penicillium rubens</i>	AQTNLSLPRVAN	WFQNRRAKAK	QOKRQEE	FEKMT	KAKAEAE
<i>Talaromyces marneffei</i>	VQTNLSLPRVAN	WFQNRRAKAK	QOKRQEE	FERMQ	REAKEKED
<i>Blastomyces dermatitidis</i>	AQTNLTLP	RVANWFQNRRAKAK	QOKRQEE	FERMQ	ASEKDEQW
<i>Histoplasma capsulatum</i>	AQTNLTLP	RVANWFQNRRAKAK	QOKRQEE	FERMQ	ASNGEQW
<i>Microsporium canis</i>	LQTSLTLP	RVANWFQNRRAKAK	QOKRQEE	FEKMQ	AKEKAMAAE
<i>Trichophyton tonsurans</i>	LQTSLTLP	RVANWFQNRRAKAK	QOKRQEE	FEKMQ	AKEKAMAAE
<i>Ascosphaera apis</i>	MQTNLTLP	RVANWFQNRRAKAK	QOKRQAE	YEKKL	AEEKAEKQNGK
<i>Arthrotrichum flagrans</i>	HAINLSPT	RVNIWFQNRRAKAK	KHKEIQE	AKMAQ	ILETAGRERMAKIALGTPMAL
<i>Alternaria alternata</i>	TRLGVPLDK	INWFQNRRAKVK	QDRKKLM	NQY	NMTMSLPPFGHSHVPM

	160	170	180	190
<i>Aspergillus nidulans</i>	SDSQRSAAEKDEKKQD	..DSK	APT	PKP
<i>Aspergillus niger</i>	SK	..EDAKDSKDET	..DKD	TP
<i>Aspergillus flavus</i>	SN	..P	..DVKEET	..DKE
<i>Aspergillus oryzae</i>	SN	..P	..DVKEET	..DKE
<i>Penicillium zonata</i>	SGSTVKAET	PDSSNA	..S	LAP
<i>Penicillium rubens</i>	GSRKGS	..ANE	ESE	..KSA
<i>Talaromyces marneffei</i>	GCDQKSP	IKHKDDNSHGTTKSP	TP	TQAS
<i>Blastomyces dermatitidis</i>	GES	..P	..KQSELPESST	TP
<i>Histoplasma capsulatum</i>	GAS	..P	..KQSERLESSA	TP
<i>Microsporium canis</i>	SSDEQKQKSE	QDKNSI	..LN	TN
<i>Trichophyton tonsurans</i>	SSDEQQRSE	QDKSSM	..S	TP
<i>Ascosphaera apis</i>	GESKEKQES	QPQESQ	NE	AEATQAA
<i>Arthrotrichum flagrans</i>	ES	..P	..S	M
<i>Alternaria alternata</i>	NHY	..A	..HPQE	QQHPHMLM

	200	210	220
<i>Aspergillus nidulans</i>	QQTRS	ENSN
<i>Aspergillus niger</i>	QKTRS	ESAREATFAS
<i>Aspergillus flavus</i>	HKTGS	ESAREATFAS
<i>Aspergillus oryzae</i>	HKTGS	ESAREATFAS
<i>Penicillium zonata</i>	QKTPS	ESAREATFAS
<i>Penicillium rubens</i>	HKTGS	ESAREATFAS
<i>Talaromyces marneffei</i>	QKTGS	DLAQEKTYAS
<i>Blastomyces dermatitidis</i>	QQASD	FTTKTGCPES
<i>Histoplasma capsulatum</i>	QQASN	SISQFGLPEP
<i>Microsporium canis</i>	LQTPA	EEKPEPRFDA
<i>Trichophyton tonsurans</i>	LQTPA	DEEPEPKFEV
<i>Ascosphaera apis</i>	TSPPY	PSK
<i>Arthrotrichum flagrans</i>	TSPPY	PSK
<i>Alternaria alternata</i>	QQHQHQ	QQQFDMQHG	LHSV

230
Aspergillus_nidulans G...GQGT...
Aspergillus_niger DRYGQDGENSQ...
Aspergillus_flavus EHYSPDEQGP...
Aspergillus_oryzae EHYSPDEQGP...
Penicillium_zonata DQFGRSGVEGEAVA...
Penicillium_rubens DRFGRRINPRSKNE...
Talaromyces_marneffei DQYTGPSDDHISVG...
Blastomyces_dermatitidis AQFNQPVENNGPDR...
Histoplasma_capsulatum AQFSQPGEDNVP...
Microsporium_canis ...EVPNPEPKLI...
Trichophyton_tonsurans ...EIPSSPEAKMI...
Ascosphaera_apis TRSEQPVLPRPSNNAQNG...
Arthrobotrys_flagrans SSYELEGVSRQGYHP...
Alternaria_alternata EFYDTTGLSNAYSDDLAFSVPAPMPNDLAPSHPEFDNFADFLDYSALAAATNTSNSASA

240 250
Aspergillus_nidulans ...SMGSSG...SVSP...D
Aspergillus_niger ...SPSMDG...SVSP...QN
Aspergillus_flavus ...ATIHEG...SVSP...ET
Aspergillus_oryzae ...ATIHEG...SVSP...ET
Penicillium_zonata PPP...ADALEG...SMP...ET
Penicillium_rubens VPEED...EEEEAVSP...SMP...EE
Talaromyces_marneffei TDR...AFDVRQTNIPVPSANNTPQTSANS...DS
Blastomyces_dermatitidis ...E...SSAITYHLSQ...SS...DN
Histoplasma_capsulatum ...G...PSDIKYHFPQ...SS...DN
Microsporium_canis SPPMK...EDQEMSG...NVVD...TAQPPFSQP...WG
Trichophyton_tonsurans SSPFK...DSQEMSG...TMTT...TAQPSF...WA
Ascosphaera_apis SQPQSTGNVNQQNF...GATSVSPAISV...NVVPDQIH...YH
Arthrobotrys_flagrans ...AAPYDG...LPPP...SDW...NS
Alternaria_alternata EAQQSTGSISSDASP...YNTQ...SGTTQSP...NGPTPPSIASLN...T

260
Aspergillus_nidulans SAVWSSVNST...NGE...LSVPG...
Aspergillus_niger AISTWTSQ...SS...QGALGYVT...GESL...HDAGHD...SMQ...NV...
Aspergillus_flavus AMSWSASQSP...QEHLGYS...AAESL...QNVQHS...D...TL...
Aspergillus_oryzae AMSWSASQSP...QEHLGYS...AAESL...QNVQHS...D...TL...
Penicillium_zonata SLTWTPSQSP...EDTFYGNL...NAATSFTSMVESLSVAEMETPQOS...FRR...
Penicillium_rubens PASWGSAE...H...NDNVGYST...TEQTSYSTSCHQAM...ADISNPSHS...MQQ...NLDVYS...
Talaromyces_marneffei SIATWTPSQSP...EEGYEFGS...LNNVPFASEVA...QMDNSPNP...DVS...TSHGFG...
Blastomyces_dermatitidis HAITWSPQGA...EDSDFPGH...LNRHHDNPLEIRV...Q...GSPHFP...
Histoplasma_capsulatum HAITWSSSQDI...EDHIVFPH...LNKPHGNPLEICV...S...ESQQFP...
Microsporium_canis ...PTPVP...QIEVNVGS...ADHPVATSAP...IM...ELDSNGYWSPDSSGIVNQPT...
Trichophyton_tonsurans ...STGVP...QIGVNVGS...ADAHTVSTAAP...VM...EIDSNGYWSPDSSGLVSOQY...
Ascosphaera_apis DAGQ...PT...
Arthrobotrys_flagrans SVSYTPTPAGQREDP...FDFNM...SALNS...PYHNAPL...SG...VPPV...QDD
Alternaria_alternata SQMAHPKQVEEPEDQFAPYS...LAQASA...SEQTLFFW...P...

270 280 290
Aspergillus_nidulans ...LENSQSFSDY...RSASDA...GAS...YN
Aspergillus_niger ...QF...HPSQNE...W...LQTSKLSGY...RSASDA...EVS...YN
Aspergillus_flavus ...QF...HSSQNE...EWSG...VQGT...KSFPGY...HSSNDA...EAS...YS
Aspergillus_oryzae ...QF...HSSQNE...EWSG...VQGT...KSFPGY...HSSNDA...EAS...YS
Penicillium_zonata ...PSDGWSHHHHQH...HHHHHHHHM...QEHVSKHHHP...SASAMEV...
Penicillium_rubens ...NH...LPPHTEWSED...RESRHD...NLV...YS
Talaromyces_marneffei ...SI...TSRSQ...MNP...QFQ...RAE...IHA...SDPM...YG
Blastomyces_dermatitidis ...SIQFSHEPD...G...W...GYSMP...HLVG...EQRHNSQDS...GDG...FY
Histoplasma_capsulatum ...CIQFNHEP...GE...W...E...CHSMP...HLMG...KQRNSSQDL...GEG...FH
Microsporium_canis ...DLLCNGSSPEEL...PC...V...Q...PT...T...P...FS...QS...FM
Trichophyton_tonsurans ...ELLCNNSSPED...LSC...Q...V...PT...T...AFT...QS...FM
Ascosphaera_apis ...ELPFRD...T...W...T...G...S...F...AG...NG...S...TE...MV...SG
Arthrobotrys_flagrans YSAFAAAVQ...MNSM...PK...MM...NL...K...SR...V...PS...GLV...LK
Alternaria_alternata ...QDGS...SQ...MYP...Q...SNFY...Q...S...NT...S...A...H...I...L...S...T...P...E...Q...A...R...K...L...S...A...P...S

300 310
Aspergillus_nidulans SMQFALQ...ADAANA...RRAS...SR...SLETGQTARPG...
Aspergillus_niger GVOYPLQ...QDLSL...RRGSS...DELAD...TL...EGI...GIN...THPSS...
Aspergillus_flavus AAQYTLH...PESSL...RRGSS...DLAD...SL...EGI...GIH...AA...
Aspergillus_oryzae AAQYTLH...PESSL...RRGSS...DLAD...SL...EGI...GIH...AA...
Penicillium_zonata SFTYPS...AASMEFT...RRGSS...DLAD...SL...EHI...GID...ATDSV...
Penicillium_rubens NMQYMPM...QAPDIVT...RRES...SALT...S...LEGI...GIC...TTGQS...
Talaromyces_marneffei SLSYSSLQ...PPSATSSR...RPSAS...EELAD...S...IGI...GIN...TAALA...
Blastomyces_dermatitidis QIPFHALQ...SPLYPEP...RRDSY...SSE...QSE...LAD...TL...F...TE...IN...NVI...L...SP
Histoplasma_capsulatum PIPFHALQ...SPLYPEP...RRGSS...SSE...QSE...LAD...TL...F...TE...IN...SMI...ET...SP
Microsporium_canis QSDMPLYS...SQNLVNQ...RHV...SL...PME...QG...DI...SPRS...GPN...SDSEL...TH
Trichophyton_tonsurans PSELPLYS...SQSIVNV...RHS...SL...PME...QG...DI...SPSS...GPN...SDSE...TH
Ascosphaera_apis VITPMPCT...PLMEHE...H...GDMV...SP...M...VNT...PV...ST...MET...PAT...SV...ISP
Arthrobotrys_flagrans NSDRPAF...S...RVAT...C...PD...L...SNV...ES...ISM...Q...THR...PM...HT...SS...ET...SS...P...V...LEN...G...PK...QL
Alternaria_alternata DL...DIP...L...HF...RED...AF...A...RRNS...S...SS...NL...AN...NM...DAI...H

```
Aspergillus_nidulans      . . . . . KKPARSLSLPAETDR.GLPRVGTSRSTSMLS.TS..TMS
Aspergillus_niger        . . . . . QLVN...EGD...RSSWKEPSKELDLAARRKRPRPAAIGTSRSSSMLT.GSS..TMS
Aspergillus_flavus       . . . . . GLPI...RTD...RSSWKEAGKELDLAARRKRPRPAAIGTSRSSSMLA.GSAA.SMS
Aspergillus_oryzae       . . . . . GLPI...RTD...RSSWKEAGKELDLAARRKRPRPAAIGTSRSSSMLA.GSAA.SMS
Penicillium_zonata        . . . . . HLNP...QRVDPAAWKEPGKELDLAARRKRPRPAAIGTSASGRSSLAAGTVSMS
Penicillium_rubens       . . . . . GLSQ...PVDREAVWKEPGKELDLAARRKRPRPAAIGTSGRPLANSTMSLS
Talaromyces_marneffei    . . . . . G...SMDSSMWRPEKELDLAARRKRPRPAAIGTAHHRLS...TNPMSVMS
Blastomyces_dermatitidis QNTPHLNLAQLHHQAEPSTNWRYPEKELVDIARRKRPRPAAIGTSMSRS...YGPSMS
Histoplasma_capsulatum  RNTPHLNMAQLRHQVDPSTNWRYPEKELVDIARRKRPRPAAIGTPAMRS...YGPSMS
Microsporium_canis       SE..G...RQLPRLHISTSDNAIGLAARRKRPRPAAIGTSLRSAL..GGPMS
Trichophyton_tonsurans  PE..G...RQPPRLHISTSDNAIGLAARRKRPRPAAIGTSGFGRTV..GGPMS
Ascosphaera_apis        MAP...PTVSDMDTWRQFKKELVDIARRKRPRPAAIGTATLGRSF..TGPSMS
Arthrotrrys_flagrans    .AAPGM...ERSQYTERPSLQEIALLRRLRPMPPSLGNARTQRIFHG.KI.NGS
Alternaria_alternata    . . . . . IRNSTPDGFPDPPDQSSIAARRK.RPVNLLSSAMRSASYSAA..PMS
```

```
Aspergillus_nidulans      PTTRG..QNY..GT...VKQSKSAQNLG...SRYAGVRKPS.AQRSPNLSTFAEAGVL
Aspergillus_niger        PSTRL..PSY..GNHAVRQSKSAQGLN...SRYAGVRKASAAQRSPNLSTFAEAGAL
Aspergillus_flavus       PTTRL..PSY..GSAPGVRQSKSAQCLN...SRYAGVRKASAAQRSPNLSTFAEAGAL
Aspergillus_oryzae       PTTRL..PSY..GSAPGVRQSKSAQCLN...SRYAGVRKASAAQRSPNLSTFAEAGAL
Penicillium_zonata        PTTRL..PSSLGATGHSVRQSKSAQSLN...SRYAGVRKVSVAQRSPNLFSTFAEAGAL
Penicillium_rubens       PTARM..PSS..GAGNSMRQSKSTQSLN...SRYAGVRKASAAQRSPNLFSTFAESGI
Talaromyces_marneffei    PNARM..ATF..GAPHTTRHAKSSHTLG...SRYAGVRKLSATQRSPNLFSTFAEATA
Blastomyces_dermatitidis PTTTRI..HGM..GAGHVLRHAKSTONLSPSHTSRYPGIRKASAPQRSPLGITSTFAESLF
Histoplasma_capsulatum  PTTTRI..QGM..GAGHVLRHAKSTONLSP...NRYPGIRKASVAQRSPLGITSTFAESRF
Microsporium_canis       PTRRV..SSA..AWGGVRKSSQLAELS...PRYASVRKLISGSPRSPFPYSLEGRHAL
Trichophyton_tonsurans  PTRRV..SSA..AWSGVRKSSQLAELS...PRFGMRKISGSPRSPFPYSLEGRHAL
Ascosphaera_apis        PTLGVTRPGYGPGHCHTLRQTKSTQSLGHSARSLSGIRKTSYNSRSPNLSTFTENSL
Arthrotrrys_flagrans    HS...VHGTPLETPSSDADF...NATVQHKLKRKPSDLKSEHS...
Alternaria_alternata    PG...GNGDKVLRIRISSGIPNA..GGRV.QKSQPGSAQRSPMVSTFSDAAS
```

```
Aspergillus_nidulans      S.SAKT...ELSTMLOPV.TTNSLAPPTPLTPEDLHHLLPTTPSTDG.YCLS.AQPTAHLF
Aspergillus_niger        G.S.KA...DMSSMLQPAVTTGG.LAPPTPLTPEDLHHLLPTTPSDGG.YCLS.AQPTSQLF
Aspergillus_flavus       G.TSKP...EMSSMLSPAVTTGG.LAPPTPLTPDDLHHFIPNTPSDGG.YCLS.AQPTSQLF
Aspergillus_oryzae       G.TSKP...EMSSMLSPAVTTGG.LAPPTPLTPDDLHHFIPNTPSDGG.YCLS.AQPTSQLF
Penicillium_zonata        S.AAKA...EMLOP.SVSANTLAPPTPLTPEDFQHLLEPASPFGG.YCLSAHPASQQL
Penicillium_rubens       K.....KAEKMLRPSISTTSLAPPTPLTPQDLQHFMPASPTD.SN.YCLS.AHSTAHFF
Talaromyces_marneffei    A.AANASSESRQKHLRHTSASVGNLAPPTPLTPEDFQHMLLPTTSTDMQN..FSTPHLT
Blastomyces_dermatitidis N.CANAT...DMMSVPLVLTITLAPPTPLTPEDLRTLPLTPND.SQ.YCVSPTDDM...
Histoplasma_capsulatum  N.CANTA...DLMSLPLGLVTTSLAPPTPLTPEDLQTLPLTPND.SQ.YCVSPTDDM...
Microsporium_canis       S.NT...DLAVPSSSTSSLPATPMTPEMQYLLEP.TPIDNQ.YCLSPQEM...
Trichophyton_tonsurans  S.NA...DLAVPSSSTSSLPATPMTPEMQYLLEP.TPIDNQ.YCLSPQEM...
Ascosphaera_apis        G.SALP...TISPLATPMTPDGARSLMP.TPNDAYA.YLSPQEC...
Arthrotrrys_flagrans    PKFART...FSTSSATTIHGGS.LAPPTPLTPQDFGNYWGGAA...L.VIRPHSAMPDH
Alternaria_alternata
```

```
Aspergillus_nidulans      . . . . . PTTOPMQINIAS. . . . . PPAATPLGMDTMS...SYPVHSVAP.PMSAPA
Aspergillus_niger        . . . . . PTTOPMQINIAS. . . . . PPAATPLAVDVL...SYPYQGVAP.PMSAPA
Aspergillus_flavus       . . . . . PTTOPMQINIAS. . . . . PPAATPMAMDMLS...TYQVHSVAP.PMSAPA
Aspergillus_oryzae       . . . . . PTTOPMQINIAS. . . . . PPAATPMAMDMLS...TYQVHSVAP.PMSAPA
Penicillium_zonata        FQSTTATTOPMQIHIAS. . . . . PPSPTLTMVLS...PFAYTTLAP.PMSAPA
Penicillium_rubens       . . . . . PTTOPMQVNMAS. . . . . PPAATPL..DIYS...PFVQNVAP.PMSAPA
Talaromyces_marneffei    DTQGNFVPTQSMQINVAS. . . . . PPEATPLTLDFVS...AMQYQNMVPLSATP
Blastomyces_dermatitidis GCARFFPTISOPMQVHIES. . . . . PPTPLHLGLVPS...HLOQYQSMGVP.PMSASP
Histoplasma_capsulatum  GCARLFPMSSOPVQVHIES. . . . . PPTPLHLGLVQS...HLOQYQSMGVP.PMSASP
Microsporium_canis       GYAHSPPTSQSMNFDENQ. . . . . ESKRQPPFVMVGMPH...AQSYQSFTE.PMSAPP
Trichophyton_tonsurans  GYSHSPPTSQSMTFDECO. . . . . ESKRQPPFVMVGMPH...PQSYQFTE.PMSAPP
Ascosphaera_apis        SYA..APPPTGMSLEPES. . . . . PPTGTFNFVGYGMPRPSLHHTSFVP.PRSAPP
Arthrotrrys_flagrans    . . . . . TNEADAPIKQNLAS. . . . . PPTPETAGLAAVNGSFKEL...E.SSDQ
Alternaria_alternata    . . . . . NSESMHTNWSDDQAGNVIAKTTSPPSSSLDLQSRFVND..ALYRDT.PMSAPA
```



```
Aspergillus_nidulans      NFTSFP.DYS...CD.GSFQGRNWE.ATSMPSPEVVPFQSQCH...Q..M
Aspergillus_niger        HYSFPA.DYAS...CE.APLTGRSWTDATSMPSPEASFQSPCQ...LPQA
Aspergillus_flavus       HYSFPA.DYVT...CEGAPLGRSWTGANSMPSPPEAFAQNRVP...ITQA
Aspergillus_oryzae       HYSFPA.DYVT...CEGAPLGRSWTGANSMPSPPEAFAQNRVP...ITQA
Penicillium_zonata        QYTFPPEYAG...SCEVPLTARSWAEAVPMPSPPEMNFVAGMP...PSIA
Penicillium_rubens       QVSSFP.EYLT...CDSVPIPARSWADTGSISSPPEYPAGLQVP...HS.S
Talaromyces_marneffei    QYASFT.DYSP...ITSEPLTGVSWAVSTPDSL...FP.SLQ...SQR
Blastomyces_dermatitidis QRTTFQEYALS...VPTNPMNGGLWSDVSSMPSPETSHLQOPT..IHMPPQ
Histoplasma_capsulatum  QHTTFQEYQLS...IPTNPMSSGGHWDPASSMSPPETSHLQOPT..IHMPPQ
Microsporium_canis      NFTTFNELIPD...CGQQQQOQQALTSSEADPS...LNV..IHMPPR
Trichophyton_tonsurans  NFTTFNELIPG...CGQQQQOQQALTSSEAEPS...LNV..IHMPPR
Ascosphaera_apis        NQHGF.PDSNNF.MPIPGSAPGTAAGPHAMS.DYAWSTGSS.LSPDTPHLQNSH..FSPHPT
Arthrotrrys_flagrans    QYTNFTKMLEQSSPIVSTAAFDGHSDFQPNVEHWISGHQPYPYWTPTHQSDNHVNDISQQ
Alternaria_alternata    TQQHF.PRTSYMQ...QPQMRAA...FHS.TTDLTIQPKP
```

		540		550		
<i>Aspergillus nidulans</i>	N	F	STPYDHALDQ.....	SOSENG.....P	
<i>Aspergillus niger</i>	D	L	TTMSYEQAMEQ.....	GA...DHVPVIGS	
<i>Aspergillus flavus</i>	D	V	SSLSYGQALEQ.....	GRQPADSLSAAGS	
<i>Aspergillus oryzae</i>	D	V	SSLSYGQALEQ.....	GRQPADSLSAAGS	
<i>Penicillium zonata</i>	Q	HS	I...SPIP.....	DQ.....GL...SMD.....	
<i>Penicillium rubens</i>	T	V	SPMGYNATTDH.....	TGQAFGMESVSGS	
<i>Talaromyces marneffei</i>	Q	QP	PIIYIEQDDDEHQ.....	DPKWTLSGDDGSSLYGSTKASAT	
<i>Blastomyces dermatitidis</i>	TH	I	SPITYEESFD	SGN.....PA.....LVEDTVTSQSES	
<i>Histoplasma capsulatum</i>	TH	I	SPITYGESL	SGN.....PA.....LVEDMMTGQRES	
<i>Microsporium canis</i>	TH	I	SPVAYDDQM	QGEQVEDSAATEDWQGGQQS.....T...QSTS	
<i>Trichophyton tonsurans</i>	TH	I	SPIAYDDVQ	GEH.VEESAPAEWQGGQPS.....STHSTTG	
<i>Ascospaera apis</i>	H	QMLPQQHNGTAQH	I	SPSMTYESPF	E	FENNHP.....MMEGMNPLSPQGTIRAASP
<i>Arthrobotrys flagrans</i>	S	QQAYGIHDGRQRH	
<i>Alternaria alternata</i>	EH	F	RRRSLP	DTAAQAG.N.....DSSSQYMQAGNMHYDEFKDVLSLGI	



	560	570	580	590
<i>Aspergillus nidulans</i>	SQSPFGD...ADI...QAPGDASKA	TEFHLYEFPDQE	EAHRFVA
<i>Aspergillus niger</i>	PSLVYHT...SDVDMP TSAAFCGDSKQ	TEFHIYEFPEQQ	EAHRFVA
<i>Aspergillus flavus</i>	PPLMYTT...DADMHTSSGSFHGDAKP	TEFYIHEFPEQQ	EAHRFVA
<i>Aspergillus oryzae</i>	PPLMYTT...DADMHTSSGSFHGDAKP	TEFYIREFPEQQ	EAHRFVA
<i>Penicillium zonata</i>HETGSYSGSPPGSHAKE	TEFRIOEFPEQQ	EAHRFVA
<i>Penicillium rubens</i>	PSLIYSI...EDTDIPGSAELA.ERKR	PEFMMHEFP	EHDT	HFG.....GH
<i>Talaromyces marneffei</i>	PPA.....NMMTVSEEHDPNGMT	TFHIHEFPKQ	EAHRNVA
<i>Blastomyces dermatitidis</i>	PQCTVKSCGTP TTSSP.QGSTPGSKV	TEFLIOEFPEQQ	EAHRRAA
<i>Histoplasma capsulatum</i>	PHCAVKNCNTP TASSP.QGSSPGSRRV	TEFLIOEFPEQQ	EAHRRAA
<i>Microsporium canis</i>	PNSPISE.....SGQYNSTGKGAS	TEFYIOEFPPQD	EAL	K.....MAA
<i>Trichophyton tonsurans</i>	PNSPVSE.....SSQAYNSAGKS.N	TEFYIOEFPPQD	EAM	K.....VAA
<i>Ascospaera apis</i>	PQMD.....QSGYPVQG...QNF	AQQSMKAQT	PEFS	FMQQLTGADERDDGTS
<i>Arthrobotrys flagrans</i>
<i>Alternaria alternata</i>	H.....HN.VPFAPQVSAMP	DFLVH	QYTP	POGTD.....HGNLL

	600	610
<i>Aspergillus nidulans</i>	QQLPNQKPKAYTFADNR	TPTNFG.....
<i>Aspergillus niger</i>	QQLPSQKPKAYTFNNQ	TPNDF.....
<i>Aspergillus flavus</i>	QQLP.QKPKAYTFNNQ	TPSDWRGN.....
<i>Aspergillus oryzae</i>	QQLP.QKPKAYTFNNQ	TPSDWRGN.....
<i>Penicillium zonata</i>	AQLSVQPKPKAYTFINNA	TGHVEA.....
<i>Penicillium rubens</i>	HLSMSQKPKAYTFANN	TPSNYPS.....
<i>Talaromyces marneffei</i>	QQLAPQIPKNYTFNSNO	TPSDF.....
<i>Blastomyces dermatitidis</i>	EQLPPQKPMNYTFNSNH	TPNDF.....
<i>Histoplasma capsulatum</i>	EQLPPQKPMNYTFNSNH	TPNDF.....
<i>Microsporium canis</i>	QQLPPQRARTYTFNTNO	TPNDFYRTAIFPPI
<i>Trichophyton tonsurans</i>	QQLPPQRARTYTFNTNO	TPNDFYRTAIFPPI
<i>Ascospaera apis</i>	AMLTPYDQGTVMFSTP	TTHLLA.....
<i>Arthrobotrys flagrans</i>
<i>Alternaria alternata</i>	RRITTEPQPKSYIFANQ	GFGDFERGQ.....



A



B




C




WT




$\Delta hbxA$




hbxA-com




OEhbxA








A




B




A**B****C****D****E**

A. flavus




A. nidulans


A



B




C




A

Fun Cat




B

Fun Cat




A

Nidulanin A




B

Nidulanin B




C

Nidulanin D




D




A

Sterigmatocystin




B




A

Austinol




B


Dehydroaustinol





C



D



A**B****C**