



**HAL**  
open science

## Evapotranspiration mapping from remote sensing data: uncertainties and ensemble estimates based on multimodel – multidata simulations

Albert Oliosio, Aubin Allies, Hugo Desrutins, Simon Carrière, Nesrine Farhani,  
José Sobrino, Drazen Skoković, Jérôme Demarty, Jordi Etchanchu, Gilles  
Boulet, et al.

### ► To cite this version:

Albert Oliosio, Aubin Allies, Hugo Desrutins, Simon Carrière, Nesrine Farhani, et al.. Evapotranspiration mapping from remote sensing data: uncertainties and ensemble estimates based on multimodel – multidata simulations. International Workshop on High-Resolution Thermal EO, ESA (European Space Agency), May 2023, ESRIN, Frascati,, Italy. hal-04153021

**HAL Id: hal-04153021**

**<https://hal.inrae.fr/hal-04153021>**

Submitted on 6 Jul 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Evapotranspiration mapping from remote sensing data: uncertainties and ensemble estimates based on multimodel-multidata simulations

Albert Olioso<sup>1 (2\*)</sup>, Aubin Allies<sup>3 (4\*)</sup>, Hugo Desrutin<sup>2</sup>, Simon Carrière<sup>5(2\*)</sup>,  
Nesrine Farhani<sup>4 (7\*)</sup>, José Sobrino<sup>6</sup>, Dražen Skoković<sup>6</sup>, Jérôme Demarty<sup>4</sup>, Jordi Etchanchu<sup>4</sup>,  
Gilles Boulet<sup>7</sup>, Samuel Buis<sup>2</sup>, Marie Weiss<sup>2</sup>

1 INRAE (Mediterranean Forest Ecology Lab, URFM), Avignon, France

2 INRAE – Avignon Université (Mediterranean Agroecosystem Modeling Lab, EMMAH), Avignon, France

3 EarthDaily Agro, Balma, France

4 IRD (HydroScience Lab, HSM), Montpellier, France

5 Sorbone University (METIS), Paris, France

6 University of Valencia (Global Change Unit), Burjassot, Spain

7 IRD (CESBIO), Toulouse, France

(\* Previous affectation)

Preparation of the future missions



LSTM @ esa



## Context

- ❑ ET is a major component of the hydrological cycle, **but**

{  
→ in many situations, ET and its evolution are not well known  
→ there is a lot of uncertainties in ET monitoring


- ❑ Many models exist, but none looks satisfactory in every situations (season, type of climate, type of surfaces...

→ **ensemble modelling**

Ensemble modelling approaches were developed in various field of research (ex. Climate, hydrology, agronomy...) assuming they are providing an optimal or suboptimal solution

- ❑ Work done in the frame of TRISHNA and LSTM mission preparations

**Evapotranspiration (ET)** can be derived using various models based on **thermal infrared data**



No consensus on a best model => ensemble averaging considering both models and data sources

Many unknowns remain concerning the uncertainties in the derivation of ET  
in particular for discriminating uncertainties from input data and models

ET estimate = average of ensemble members


Uncertainty =  
standard deviation of ensemble members

## Models of Latent Heat Flux (LE)

[see Lagouarde and Boulet 2016]

-> **Evaporative fraction (EF) model:**  $LE \sim EF \times (Rn - G)$   
with  $EF = BC / BA$

$EF$  = evaporative fraction <-  $T_s$  vs. *albedo or NDVI or fraction cover*  
 $Rn$  = net radiation <- { *albedo, emissivity,  $T_s$ , solar irradiance*  
*atmospheric irradiance*  
 $G$  = ground heat flux <-  *$Rn$ , NDVI, fCOVER*




ex: S-SEBI (Roerink et al. 2000)

Upscaling to daily level based on

$$ET \sim LE \times Rgd / Rgi$$

-> **Residual aerodynamic equation:**  $LE = Rn - H - G$

$H$  = sensible heat flux <- {  $T_s, T_a$  (air temperature),  
 $u_a$  (wind speed)  
 $zom, zoh$  (roughness)



ex: SSEBE (Oliosio et al. 2006)

## Application:

ESA experiment in Grosseto (Italy) in support of the LSTM program : July 2018


Airborne images in the solar and the thermal domains on two different days

Various sources for input data :



- incident radiations,
- LAI,
- fCOVER,
- surface temperature...

Various « models » for

- Ground heat flux
- Evaporative fraction EF
- albedo



## Example of variations (=> uncertainties) in data sources and models




### Uncertainty

**Novice case** -> all available data or models are used

**Expert case** -> previous knowledge => { some of the data sources and algorithms are dropped

**Global uncertainty** -> standard deviation of ET (pixel basis) all together, ~400 000 cases (*ET*)


**Factor uncertainty** -> standard deviation of ET for variations of one factor only






## Uncertainty in Ts:

several processings were performed mainly differing in the atmospheric profiles used for the atmospheric corrections



Ts standard deviation map Day 2 (K)







## Evapotranspiration map

Average of all the calculations for Day 2

Novice case – average ET = 3.4 mm d<sup>-1</sup>




Expert case - average ET = 2.8 mm d<sup>-1</sup>





**Uncertainty in ET**  
(standard deviation  $\text{mm d}^{-1}$ )

Day 2 case




**Expert case**

**Novice case**




**Uncertainty in ET (mm d<sup>-1</sup>) Day 2 case**


Expert case




Novice case



**Uncertainty related to Ts:**




### ET uncertainty break-down




Ts

### ET uncertainty break-down




## Applications of EVASPA to the Crau area

EVASPA is an implementation of the method that will be used for the level 2 ET product for TRISHNA (together with the STIC model (Mallick et al.))



Monthly ET obtained over the Crau aquifer (600 km<sup>2</sup>) with the evaporative fraction models for MODIS TERRA and AQUA data :





## Summary

Ensemble modelling applied to multi-data source – multi-model (or algorithm) may be used for:

- monitoring ET
  - providing uncertainty in the estimates
- (however this uncertainty is only epistemic and does not include estimation errors)
- providing information on the main uncertainty factors:
    - in all analysed cases, surface temperature was not the main limitation in ET estimations
    - for contextual models, the main sources of uncertainty concern algorithm (EF and ground heat flux)
    - for other models, including aerodynamic equations, meteorological forcing of wind speed and air temperature have also a strong impact

The EVASPA algorithm is a simple algorithm that will be the basis for implementing ET products in the frame of the TRISHNA program