



**HAL**  
open science

## Metabolization of flavan-3-ol oligomers by human gut bacteria

Halifa Ruben, Agnes Cornu, Pierre Peyret, Claire Dufour, Carine Le Bourvellec, Pascale Mosoni

► **To cite this version:**

Halifa Ruben, Agnes Cornu, Pierre Peyret, Claire Dufour, Carine Le Bourvellec, et al.. Metabolization of flavan-3-ol oligomers by human gut bacteria. 13th International Gut Microbiology Symposium, Jun 2023, Aberdeen, United Kingdom. hal-04158076

**HAL Id: hal-04158076**

**<https://hal.inrae.fr/hal-04158076v1>**

Submitted on 10 Jul 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Metabolization of flavan-3-ol oligomers by human gut bacteria

HALIFA Ruben, CORNU Agnes, PEYRET Pierre, DUFOUR Claire, LE BOURVELLEC Carine, MOSONI Pascale

UMR454 MEDIS, Université Clermont Auvergne, INRAE, Clermont-Ferrand, France

UMR1213 Herbivores, VetAgro Sup-Université Clermont Auvergne, INRAE, Clermont-Ferrand, France

UMR408 SQPOV, INRAE, Avignon Université, Avignon, France

Flavan-3-ols are a largely consumed subclass of flavonoids and are involved in the prevention of cardiovascular diseases [1]. The contribution of phenolic metabolites produced by the gut microbiota in the health effects of polyphenols (including flavan-3-ols) is currently an open field of investigation. Although a few bacterial species metabolize flavan-3-ol monomers [2] no bacterial isolates active on oligomers (called procyanidins) have been described. Knowing that the gut microbiota hydrolyzes procyanidins [3], our aim was to identify bacteria degrading flavan-3-ol oligomers and the degradation products. From human stools of three healthy individuals, culturomic approaches combined with screening for the metabolic activity of bacterial isolates by HPLC-DAD allowed us to obtain four strains of *Eggerthella lenta* and one strain of *Flavonifractor plautii* degrading (+)-catechin and (-)-epicatechin. The activity of these strains was then tested on B-type (DP2 to 4) and A-type (DP2) procyanidins and the metabolites generated were characterized by LC-ESI-MS / MS. These two species co-metabolized (+)-catechin and (-)-epicatechin into hydroxyphenylvaleric acid derivatives. Only *E. lenta* converted procyanidins while *F. plautii* alone or in co-culture with *E. lenta* did not show any activity towards procyanidins. The reaction catalyzed by *E. lenta* on dimers (B-type and A-type) corresponded to the opening of the C-ring of the terminal unit. This work is the first report of flavan-3-ol oligomers metabolization by the human gut bacterium *E. lenta*.

1. Fraga et al (2019). Food Funct. 10, 514-528.
2. Braune & Blaut (2016), Gut Microbes, 7, 216-234
3. Le Bourvellec et al (2019) Nutrients, 11, 664.