



HAL
open science

Tanned but not burned: A negative feedback loop controls Citrus fruit coloration

Louis-Valentin Méteignier

► **To cite this version:**

Louis-Valentin Méteignier. Tanned but not burned: A negative feedback loop controls Citrus fruit coloration. *The Plant cell*, 2023, 35 (4), pp.1163-1164. 10.1093/plcell/koad017. hal-04160154


HAL Id: hal-04160154

<https://hal.inrae.fr/hal-04160154v1>

Submitted on 12 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Tanned but not burned: A negative feedback loop controls *Citrus* fruit coloration

Louis-Valentin Méteignier ^{1,2,*}

¹ Assistant Features Editor, The Plant Cell, American Society of Plant Biologists, USA

² PHIM Plant Health Institute, Univ Montpellier, INRAE, CIRAD, Institut Agro, IRD, Montpellier, France

*Author for correspondence: louis-valentin.meteignier@inrae.fr

Fruits are essential for the reproductive success of angiosperms by promoting seed maturation and dispersal. The attraction of seed dispersers is influenced by fruit scent and coloration, and the biosynthesis of carotenoids is central for the production of volatile precursors and pigments (Rodríguez et al., 2013). In specific orange varieties such as *Citrus sinensis*, the accumulation of the apocarotenoid β -citraurin (β -C) in the fruit peel confers a deep orange color. These specific varieties possess a distinct miniature inverted-repeat transposable element allele in the promoter of the gene encoding the last step of β -C biosynthesis, creating a basic helix-loop-helix protein-binding element (E-box) associated with increased gene expression in orange peels (Zheng et al., 2019).

The phytohormone jasmonic acid (JA) promotes carotenoid accumulation in tomato (Liu et al., 2012), and the basic helix-loop-helix transcription factor (TF) MYC2 is a master regulator of the JA response (Lorenzo et al., 2004). However, how JA influences β -C accumulation in *Citrus* remained unknown. In this issue, Pengtao Yue and colleagues (Yue et al., 2022) provide evidence that β -C biosynthesis is regulated by a JA-coordinated negative feedback regulatory module involving CsMYC2 (see Figure) in *Citrus sinensis*.

The authors first confirmed that endogenous levels of β -C, JA, and the transcripts encoding carotenoid biosynthetic enzymes showed some degree of correlation in orange peels, as originally shown by Liu et al. (2012). In addition, pretreatment of detached fruits with ethylene or abscisic acid inhibitors did not prevent the promotion of fruit coloration by methyl-jasmonate (MeJA) treatment, suggesting that MeJA-induced fruit coloration was independent of ethylene or ABA. In their quest for the identification of a TF required for JA-induced fruit coloration, the authors observed that

among the three detected MYC-encoding genes in *Citrus*, only CsMYC2 transcripts accumulated more in fruit peels after MeJA treatment and during natural fruit coloration. Through a compelling series of in vitro and in vivo experiments, the authors show that CsMYC2 interacted with E-box-containing promoters of carotenoid biosynthesis genes and stimulated their transcription in transactivation assays (see Figure). Importantly, MeJA treatment reinforced the association of CsMYC2 with E-boxes as observed by chromatin immunoprecipitation, followed by quantitative PCR (ChIP-qPCR) experiments. Interestingly, the authors also observed a rapid and transient increase in the accumulation of carotenoid gene transcripts after MeJA treatment, with the peak of expression preceding the peak of JA and β -C accumulation. The authors therefore posited that a negative feedback loop might occur to dampen an excessive JA response.

Among the 37 detected orthologs of JA repressors in Arabidopsis, transcript levels of *Mitogen-Activated Protein Kinase 6* (CsMAPK6) and two other putative repressors were increased by MeJA treatment in *C. sinensis*. The authors performed ChIP-qPCR and transactivation experiments suggesting that CsMYC2 interacted with the E-box-containing CsMAPK6 promoter to activate transcription. They also confirmed that CsMYC2 and CsMAPK6 interacted as observed in Arabidopsis (Sethi et al., 2014). To understand the molecular consequences of the CsMYC2-CsMAPK6 interaction, the authors hypothesized that CsMAPK6 phosphorylates CsMYC2. Using a phos-tag biotin probe, the researchers observed that phosphorylation of CsMYC2 occurred only when CsMAPK6 was co-expressed. Mass spectrometry analysis of CsMYC2 identified several phosphorylated amino acids. Site-directed mutagenesis of these amino acids to alanine prevented the detection of CsMYC2 by phos-tag biotin assay




Figure The JA-orchestrated MYC2-MAPK6 negative feedback regulatory module controls orange coloration. During maturation, JA levels increase and activate MYC2 expression. Then, MYC2 binds the E-boxes in carotenoid biosynthetic and *MAPK6* gene promoters to stimulate transcription. At this stage, fruit coloration occurs. Upon accumulation of excessive amounts of JA, MAPK6 binds and phosphorylates MYC2 to induce its degradation, causing an arrest in fruit coloration. Adapted from Yue et al. (2022) with images taken from bioicons.com and commons.wikimedia.org.

when co-expressed with *CsMAPK6*, thus revealing that they are *CsMAPK6*-dependent phosphorylation sites. The authors then confirmed the effect of phosphorylations on *CsMYC2* protein stability by transient expression of phospho-variants in *Citrus calli*. They further showed that co-expression of *CsMYC2* and *CsMAPK6* decreased the transactivation potential of *CsMYC2*, and co-incubation of *CsMYC2* with *CsMAPK6* reduced the binding of *CsMYC2* with the E-box, suggesting that the interaction alone could reduce *CsMYC2* TF activity by steric hindrance. Finally, the authors validated the role of *CsMYC2* and *CsMAPK6* in fruit coloration by transient overexpression or silencing experiments in orange peels. Transient overexpression of *CsMAPK6* or of an antisense *CsMYC2* transcript decreased β-C accumulation and orange coloration.

Collectively, these results led the authors to propose that in parallel to its role in activating carotenoid genes transcription upon increased JA levels, MYC2 also stimulates the transcription of *MAPK6*, which in turn binds and displaces MYC2 from biosynthesis genes. After protein interaction, MAPK6 phosphorylates MYC2 to induce its degradation and avoid an excessive JA response (see Figure).

References

- Liu L, Wei J, Zhang M, Zhang L, Li C, Wang Q (2012) Ethylene independent induction of lycopene biosynthesis in tomato fruits by jasmonates. *J Exp Bot* **63**(16): 5751–5761
- Lorenzo O, Chico JM, Saénchez-Serrano JJ, Solano R (2004) JASMONATE-INSENSITIVE1 encodes a MYC transcription factor essential to discriminate between different jasmonate-regulated defense responses in Arabidopsis. *Plant Cell* **16**(7): 1938–1950
- Rodríguez A, Alquézar B, Peña L (2013) Fruit aromas in mature fleshy fruits as signals of readiness for predation and seed dispersal. *New Phytol* **197**(1): 36–48
- Sethi V, Raghuram B, Sinha AK, Chattopadhyay S (2014) A mitogen-activated protein kinase cascade module, MKK3-MPK6 and MYC2, is involved in blue light-mediated seedling development in Arabidopsis. *Plant Cell* **26**(8): 3343–3357
- Yue P, Jiang Z, Sun Q, Wei R, Yin Y, Xie Z, Larkin RM, Ye J, Chai L, Deng X (2023) Jasmonate activates a *CsMPK6*-*CsMYC2* module that regulates the expression of β-citraurin biosynthetic genes and fruit coloration in orange (*Citrus sinensis*). *Plant Cell* **35**(4): 1167–1185
- Zheng X, Zhu K, Sun Q, Zhang W, Wang X, Cao H, Tan M, Xie Z, Zeng Y, Ye J, et al. (2019) Natural variation in *CCD4* promoter underpins species-specific evolution of red coloration in citrus peel. *Mol Plant* **12**(9): 1294–1307