



HAL
open science

A chromatin modification at the crossroads of cell fates in pollen

Louis-Valentin Méteignier

► **To cite this version:**

Louis-Valentin Méteignier. A chromatin modification at the crossroads of cell fates in pollen. The Plant cell, 2022, 34 (8), pp.2823-2824. 10.1093/plcell/koac145 . hal-04160165

HAL Id: hal-04160165


<https://hal.inrae.fr/hal-04160165v1>

Submitted on 23 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright



A chromatin modification at the crossroads of cell fates in pollen

Louis-Valentin Méteignier  ^{1,2,*}

¹ Assistant Features Editor, *The Plant Cell*, American Society of Plant Biologists, USA

² Université de Tours, EA2106 “Biomolécules et Biotechnologies Végétales,” Tours 37000, France

*Author for correspondence: meteignier@univ-tours.fr

In Brief

During gametogenesis in angiosperms, each microspore (i.e. the product of meiosis) asymmetrically divides to form a bicellular pollen grain composed of the vegetative cell (VC) and the generative cell. The generative cell then divides symmetrically to generate two sperm cells (i.e. the male germline [MG] cells), thus giving rise to the mature tricellular pollen grain. The VC is responsible for pollen tube formation, which is required for the transport of the two sperm cells to the embryo sac. During fertilization, one sperm cell fertilizes the egg cell to generate the zygote, while the other fuses with the central cell to initiate endosperm development. Although both the VC and MG cells play essential but different roles during sexual reproduction in flowering plants, the mechanisms controlling differentiation of the microspore toward each distinct cell type are not well understood.

Previous work has shown that extensive chromatin remodeling occurs during MG differentiation. Notably, MG-specific elimination of H3K27me₃, a repressive histone modification associated with facultative heterochromatin, mediates the transcriptional reprogramming that is required for sperm cell specification. By contrast, H3K27me₃ is retained in the VC, albeit the importance of this modification in that cell remained unclear (Borg et al., 2020). To understand the role of H3K27me₃ in VC differentiation, Xiaorong Huang and Meng-Xiang Sun (Huang and Sun, 2022) applied molecular cytology and genome-wide sequencing approaches in the context of VC-specific H3K27me₃ erasure.

The authors used the *LAT52* promoter to drive expression of the gene encoding the H3K27me₃ demethylase REF6 specifically in the VC. Using immunofluorescence microscopy, the authors observed the loss of H3K27me₃ in REF6-

expressing VCs relative to control VCs. The selective erasure of H3K27me₃ in VCs led to pollen germination defects (see Figure). Interestingly, MG-specific chromatin markers appeared in REF6-targeted VC nuclei, including the repressive histone modification H3K9me₂, which is characteristic of constitutive heterochromatin. These results suggested that removal of H3K27me₃ can induce the transition from a VC state to an MG state in pollen.

To understand the effects of H3K27me₃ erasure from VCs on gene expression, Huang and Sun then mapped H3K27me₃ by CUT&Tag, assayed chromatin accessibility by ATAC-seq, and probed gene expression by RNA-seq in sorted VC and MG cells of untransformed plants and plants expressing REF6 in VCs. First, the authors confirmed that H3K27me₃ depletion in REF6-expressing VCs was comparable to H3K27me₃ levels in wild-type MG cells, in contrast to untransformed VCs, in which H3K27me₃ is present at higher levels. Then, the authors observed that chromatin accessibility in transgenic REF6-expressing VCs was more affected in genomic regions corresponding to developmental and/or signaling genes in pollen. Indeed, H3K27me₃ removal decreased chromatin accessibility at genes involved in VC differentiation, while it increased chromatin accessibility at genes required for establishing the MG cells. Finally, expression of genes involved in VC and MG cell identity was decreased or increased, respectively, in the REF6-expressing VCs, a result consistent with the observed changes in chromatin accessibility (see Figure). Together, these results support a model in which VC-targeted H3K27me₃ demethylation induces a transition of VC toward MG cell identity. To confirm these findings, the authors analyzed untransformed and *ProLAT52:REF6-GFP* pollen grains by transmission electron microscopy. Strikingly, the chromatin of




Figure H3K27me3 contributes to pollen germination by enabling chromatin accessibility and expression of VC genes. Upper panel, Photographs of *in vitro* pollen germination assays. Sperm cells are marked in red and REF6-GFP delineate VCs (adapted from Huang and Sun [2022], Figure 2). Lower panel, Model for H3K27me3-mediated VC differentiation (adapted from Huang and Sun [2022], Figure 7; nucleosome images are licensed CC-BY 4.0 and freely available at bioicons.com). Pollen: grey, VC: blue, and SC: dark yellow.

REF6-expressing VCs, but not control VCs, formed abnormally condensed nuclear foci reminiscent of the densely packed, H3K9me2-enriched chromatin of wild-type MG cells (Pinon et al., 2017).

In conclusion, this study expands on the well-established role of H3K27me3 during development to include the specification of all the different types of cells in pollen grains, while raising other exciting questions. H3K27me3 has been shown to work in a combinatorial manner with other histone modifications such as H3K9me2, which suggests a possible interplay between H3K27me3 and H3K9me2 in cell fate specification in pollen (Pinon et al., 2017). For instance, are the mechanisms involved in chromatin condensation and repression of genes required for VC differentiation (after H3K27me3 erasure) related? Does H3K27me3 prevent the spread of H3K9me2 in genes required for VC differentiation?

Future studies should enable a comprehensive understanding of the contribution of epigenetic mechanisms in specifying cell fate in the germline of plants.

References

- Borg M, Jacob Y, Susaki D, LeBlanc C, Buendía D, Axelsson E, Kawashima T, Voigt P, Boavida L, Becker J, et al. (2020) Targeted reprogramming of H3K27me3 resets epigenetic memory in plant paternal chromatin. *Nat Cell Biol* **22**: 621–629
- Huang X, Sun M-X (2022) H3K27 methylation regulates the fate of two cell lineages in male gametophytes. *Plant Cell* **34**: 2989–3005
- Pinon V, Yao X, Dong A, Shen WH (2017) SDG2-mediated H3K4me3 is crucial for chromatin condensation and mitotic division during male gametogenesis in Arabidopsis. *Plant Physiol* **174**: 1205–1215