



HAL
open science

Mapping black Locust stands with Sentinel-2 and Ven μ s time series images

Zoé Ginter, Baptiste Hautdidier

► **To cite this version:**

Zoé Ginter, Baptiste Hautdidier. Mapping black Locust stands with Sentinel-2 and Ven μ s time series images: A test in Médoc, France. Non-native Trees for European Forests, Sep 2018, Vienne, Austria. hal-04160611

HAL Id: hal-04160611

<https://hal.inrae.fr/hal-04160611>

Submitted on 12 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MAPPING BLACK LOCUST STANDS WITH SENTINEL-2 AND VENUS TIME SERIES IMAGES

A test in Médoc, France

Zoé Ginter ^{1,2}, Baptiste Hautdidier ¹
¹: UR ETBX, Irstea ²: UMR Passages, Bordeaux-Montaigne University

COST Action FP1403 NNEXT
 International Conference
 NON-NATIVE TREE SPECIES
 for EUROPEAN FORESTS
 VIENNA, AUSTRIA
 12-14 September 2018





WHY SATELLITE DATA?

In France, *Robinia* sp. has been included in the latest version of the NFI large-scale forest mapping (*BD Forêt V2*). Considering the shortness of the list of broadleaved species, this addition provides an invaluable asset for management and geographical analysis. Due to a production process based on the interpretation of aerial photos, the **space-time coverage is yet uneven**. With a minimum mapping unit of 0,5 ha (and a width of 20 m), it may underestimate the **diffuse spatial dynamics** of pioneer tree species.

WHY NOW?


The recent earth observation satellites Sentinel-2 (launched in jun. 2015 & mar. 2017) and Venus (aug. 2017) have been hailed as game changers for the monitoring of vegetation: a high revisit rate (5 days) and a multispectral high resolution camera allows discriminating forest stands by the analysis of the **spatial time-series (SITS)** of their surface reflectance ^[1].



IS ROBINIA SP. A GOOD CANDIDATE FOR THIS?


Black Locust is able to expand in urban fringes and peripheries, cropland-vineyards interfaces, along linear infrastructure... It has also a **specific phenological pattern** (peak in chlorophyllous activity in summer, flowering in May ^[2]).

LOCATION




TENTATIVE RESULTS

As proposed by Karasiak & al. ^[1], a machine-learning method delivers a relevant supervised classification: the **Support Vector Machine** algorithm of the Orfeo ToolBox (with a Radial Basis Function kernel). It discerns most *Robinia* forests from their neighbouring Pine & Oak-dominated stands. It does also a good job on other broadleaved (e.g. Poplar Vs. Alder/Ash).



METHODS



MAY THE NFI MAPS OVER-ESTIMATE ROBINIA STANDS?

Part of the discrepancies between the classification and the NFI map can be attributed to time (a 4 year gap) and scale (with a marked improvement on edges and small woodlands). But Robinia forests are here actually **far from monocultures**: along with young coppices, it is also very present in mixed stands under *Q. robur* & *P. pinaster*. More originally, it was also found under high concurrence by Aspen (*Populus tremula*) and Alder Burckthorn (*Frangula alnus*).

THE NEXT STEPS

Densifying the set of ground truth points for proper cross-validation & area comparison (NFI). Testing Sentinel-2 3A & Venus 2A over a season once images are available. Replicating the model on neighbouring regions.

MASKING THE CLOUDS OUT, AN ESSENTIAL TASK WITH 'SITS' DATA


Clouds are the main concern of remote sensors, but the need is even more pressing with time-series. The use of mask clouds (with level 2A data), followed by a linear interpolation was here instrumental. But frontier effects (& differences between ESA & Muscate algorithms) leave out **spatial artefacts**, to which some algorithms are highly sensitive (e.g. K-means unsupervised classifications). Seamless monthly composites provided by Theia (Level 3A) might be a trade-off to consider.

REFERENCES

- [1] Karasiak, Sheeren, Dejoux, Willm, & Monteil (2018) Monitoring and understanding the green-leaf phenology of tree species with Sentinel-2, *Sentinel-2 Theia Workshop*, Toulouse.
- [2] Somodi, Čarni, Ribeiro & Podobnikar T (2012) Recognition of the invasive species *Robinia pseudacacia* from combined remote sensing and GIS sources. *Biological Conservation* 150(1):59-67.

IT'S ALL WELL AND GOOD, BUT WHERE IS THE NICE FLOWERING!?

Robinia's flowering event is **short**: typically 10 days around the 15/05 in SW France. In case of a **cloudy spring**, whatever the merits of the observation platform, assuming a favourable window is a risky gamble: no image over Médoc in 2017 and 2018! The biological trait of black Locust that is the most evident for the neighbouring dwellers is thus the one that would remain elusive from the above.



CONTACT

Baptiste Hautdidier | baptiste.hautdidier@irstea.fr
 +331 5 57 89 01 90
 Institut national de recherche en sciences et technologies pour l'environnement et l'agriculture
 Centre de Bordeaux - UR ETBX
 www.irstea.fr/en/institute/centers/bordeaux

FUNDING BY

