



HAL
open science

Improving Weed Management Based on the Timing of Emergence Peaks: A Case Study of Problematic Weeds in Northeast USA

Bryan Brown, Eric R Gallandt, Antonio Ditommaso, Paul Salon, Richard G Smith, Matthew R Ryan, Stéphane Cordeau

► **To cite this version:**

Bryan Brown, Eric R Gallandt, Antonio Ditommaso, Paul Salon, Richard G Smith, et al.. Improving Weed Management Based on the Timing of Emergence Peaks: A Case Study of Problematic Weeds in Northeast USA. *Frontiers in Agronomy*, 2022, 4, pp.art. 888664. 10.3389/fagro.2022.888664 . hal-04163681

HAL Id: hal-04163681

<https://hal.inrae.fr/hal-04163681>

Submitted on 17 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Improving Weed Management Based on the Timing of Emergence Peaks: A Case Study of Problematic Weeds in Northeast USA

OPEN ACCESS

Edited by:

Karla Leigh Gage,
Southern Illinois University
Carbondale, United States

Reviewed by:

Vipan Kumar,
Kansas State University, United States
Panagiotis Kanatas,
University of Patras, Greece
Virginia Nichols,
Arizona State University, United States

*Correspondence:

Stéphane Cordeau
stephane.cordeau@inrae.fr

†ORCID:

Bryan Brown
orcid.org/0000-0002-7666-2209
Eric R. Gallandt
orcid.org/0000-0001-6684-6250
Antonio DiTommaso
orcid.org/0000-0001-8215-2777
Richard G. Smith
orcid.org/0000-0002-3571-5888
Matthew R. Ryan
orcid.org/0000-0001-5667-9452
Stéphane Cordeau
orcid.org/0000-0003-1069-8388

Specialty section:

This article was submitted to
Weed Management,
a section of the journal
Frontiers in Agronomy

Received: 03 March 2022

Accepted: 13 May 2022

Published: 21 June 2022

Citation:

Brown B, Gallandt ER, DiTommaso A,
Salon P, Smith RG, Ryan MR and
Cordeau S (2022) Improving
Weed Management Based on
the Timing of Emergence Peaks:
A Case Study of Problematic
Weeds in Northeast United States.
Front. Agron. 4:888664.
doi: 10.3389/fagro.2022.888664

Bryan Brown^{1†}, Eric R. Gallandt^{2†}, Antonio DiTommaso^{3†}, Paul Salon⁴,
Richard G. Smith^{5†}, Matthew R. Ryan^{3†} and Stéphane Cordeau^{3,6*†}

¹ New York State Integrated Pest Management Program, Cornell University, Cornell AgriTech, Geneva, NY, United States, ² School of Food and Agriculture, University of Maine, Orono, ME, United States, ³ Soil and Crop Sciences, School of Integrative Plant Science, Cornell University, Ithaca, NY, United States, ⁴ United States Department of Agriculture – Natural Resources Conservation Service (USDA-NRCS) Soil Health Division, Big Flats Plant Materials Center, Corning, NY, United States, ⁵ Department of Natural Resources and the Environment, University of New Hampshire, Durham, NH, United States, ⁶ Agroécologie, INRAE, Institut Agro, Univ. Bourgogne, Univ. Bourgogne Franche-Comté, Dijon, France

We reviewed the timing of the peak rate of emergence for 15 problematic weed species as well as ways to use this knowledge to improve control. Much of the previous literature modeled emergence based on growing-degree-days. For these models, we input average temperature data from several zones of Northeast USA. Within species, model-predicted peak emergence in the warmest and coolest zones differed by an average of 39 days. Also within species, there was some variation between models, likely reflecting different conditions in study locations and population-level differences that will need to be addressed in future modelling efforts. Summarizing both observed and modelled results, emergence typically peaked early-season for barnyardgrass, Canada thistle, common lambsquarters, common ragweed, giant foxtail, large crabgrass, perennial sowthistle, and smooth crabgrass. Emergence typically peaked mid-season for hairy galinsoga, mouseear chickweed, and red sorrel. Emergence typically peaked late-season for annual bluegrass. Several species emerged in a protracted manner, including common chickweed, quackgrass, and redroot pigweed. With this improved knowledge, farmers may target key problematic species of a particular field in several ways. Weed seedling control efforts can be timed at the highest densities or most vulnerable phenological stage. Residual herbicides and suppressive mulches can be timed to maximize effectiveness prior to their breakdown. And if management flexibility allows, crop selection and associated planting dates may be adjusted to improve crop competition or facilitate seedbank depletion through timely bare fallow periods. Such improvements to weed management based on timing of emergence will likely become even more impactful as predictive model reliability continues to improve.

Keywords: emergence periodicity, seedbank, tillage, annual weed, ecologically-based management, stale seedbed

INTRODUCTION

The timing of weed emergence is among the most important variables determining how species respond to management (Ryan et al., 2010; Cordeau et al., 2017c). This is especially pertinent for weeds germinating from seed, which are most vulnerable to management at the time of emergence (Mohler, 2001). The timing of emergence can also correlate with other biological traits of weeds that impact management success, such as seed weight, cotyledon type, and photosynthetic pathway (Cordeau et al., 2017b). Therefore, improved knowledge of weed emergence periodicity may be used to enhance management tactics (Bastiaans et al., 2008; Norsworthy et al., 2012; Reinhardt Piskackova et al., 2021). For example, the timing of weed control efforts (Forcella, 1999; Batlla et al., 2020), crop planting dates (Royo-Esnal et al., 2018; Sousa-Ortega et al., 2020), and seedbank depletion tactics (Nordell and Nordell, 2009) may all be adjusted to disadvantage problematic weed species.

But emergence periodicity varies greatly between species, reflecting contrasting patterns of dormancy, germination requirements, and pre-emergence growth rates, which are mediated by many abiotic factors (Batlla et al., 2020) so that even within species, empirically observed patterns of emergence can range widely (Cordeau et al., 2017b; Cordeau et al., 2017a). To improve management, modeling efforts have attempted to predict cumulative emergence for individual species, typically using Weibull or Gompertz functions (Royo-Esnal et al., 2020) with growing-degree-days (Forcella et al., 1997; Renner et al., 1999; Sousa-Ortega et al., 2020) or cooling-degree-days (Taylor et al., 2021) calculated using base temperatures required for germination of each modeled species (Masin et al., 2012). Addition of base water potential (Cao et al., 2011; Šoštarčić et al., 2021) or other edaphic parameters (Archer et al., 2006; Davis et al., 2013) may be used to further improve model accuracy.

In previous work investigating emergence periodicity at several sites in Northeast USA, Cordeau et al. (2017b) tilled new plots every two weeks and recorded weed densities by species. It was observed that some weed species exhibited prominent emergence peaks (**Figure 1**), which may be targeted for improved management. Emergence peaked early for most summer annual species, including common lambsquarters (*Chenopodium album* L.), common ragweed (*Ambrosia artemisiifolia* L.), giant foxtail (*Setaria faberi* Herrm.), large crabgrass [*Digitaria sanguinalis* (L.) Scop.], and smooth crabgrass (*Digitaria ischaemum* Schreb. ex Muhl.). But redroot pigweed (*Amaranthus retroflexus* L.), barnyardgrass [*Echinochloa crus-galli* (L.) Beauv.], and hairy galinsoga (*Galinsoga quadriradiata* Cav.) demonstrated a

delayed emergence peak compared to the other summer annuals. Winter annuals, including common chickweed [*Stellaria media* (L.) Vill.] and annual bluegrass (*Poa annua* L.), as well as perennials, including mouseear chickweed (*Cerastium fontanum* Baumg.), perennial sowthistle (*Sonchus arvensis* L.), red sorrel (*Rumex acetosella* L.), and quackgrass [*Elymus repens* (L.) Gould] generally exhibited later or bimodal emergence peaks. The exception was the perennial, Canada thistle [*Cirsium arvense* (L.) Scop.], which emerged early, and for an extended period.

This review further examines the emergence periodicity of these 15 species – many of which have been cited by farmers as the most problematic weeds of not only Northeast USA (Jabbour et al., 2014), but also Midwest USA (Gibson et al., 2006), and Europe (Bourgeois et al., 2019). For each species, we review the empirically-observed and model-derived emergence peaks and provide an updated synthesis of emergence-based management recommendations applicable to Northeast USA and similarly temperate regions.

REVIEW METHODS

We conducted a literature search using ProQuest and Google Scholar with search terms “weed,” AND “emergence,” AND “timing,” “periodicity,” OR “phenology.” Additional searches replaced “weed” with the scientific name of each of our 15 species. Species entries in the series, *The Biology of Canadian Weeds*, were also reviewed. Pertinent sources cited in the resulting articles were also reviewed. No geographic limitations were imposed on our search so we could contextualize the emergence of each species over a broad range of locations. Since many of the empirically-based emergence periodicity results were qualitative, we presented them in table form (**Table 1**). This literature search also provided the basis for our discussion of management tactics that can be improved using knowledge of emergence periodicity.

Some results of our literature search predicted 50% cumulative emergence based on growing-degree-days. This timing typically coincides with the peak, or greatest rate of emergence (Royo-Esnal et al., 2020). To determine the timings associated with the growing-degree-days required for 50% emergence, we used weather data representing four prominent USDA Plant Hardiness Zones found in Northeast USA (USDA, 2012). Specifically, from coldest to warmest, we used weather data representing Zones 4a, 5a, 6a, and 7a, with average annual minimum temperatures of -34.4 to -31.7°C, -28.9 to -26.1°C, -23.3 to -20.6°C, and -17.8 to 15.0°C, respectively. Within each zone, we selected three representative weather station sites from the Network for Environment and Weather Applications (<https://newa.cornell.edu>) to encompass a wide geographic range. Zone 4a included Milan, NH, Morrisville, VT, and Saranac Lake, NY. Zone 5a included Bangor, ME, Ceres, NY, and Saratoga Springs, NY. Zone 6a included Cabot, PA, Geneva, NY, and Northborough, MA. Zone 7a included Atlantic City, NJ, Philadelphia, PA, and Riverhead, NY. Growing-degree-days were calculated in accordance with each prior study, but using mean daily maximum and minimum temperature data from 2018 to

Abbreviations: annual bluegrass, *Poa annua* L.; barnyardgrass, *Echinochloa crus-galli* (L.) Beauv.; Canada thistle, *Cirsium arvense* (L.) Scop.; common chickweed, *Stellaria media* (L.) Vill.; common lambsquarters, *Chenopodium album* L.; common ragweed, *Ambrosia artemisiifolia* L.; giant foxtail, *Setaria faberi* Herrm.; hairy galinsoga, *Galinsoga quadriradiata* Cav.; large crabgrass, *Digitaria sanguinalis* (L.) Scop.; mouseear chickweed, *Cerastium fontanum* Baumg.; perennial sowthistle, *Sonchus arvensis* L.; quackgrass, *Elymus repens* (L.) Gould; redroot pigweed, *Amaranthus retroflexus* L.; red sorrel, *Rumex acetosella* L.; smooth crabgrass, *Digitaria ischaemum* Schreb. ex Muhl.




FIGURE 1 | Violin plots of emergence of the five most abundant weeds following tillage on different dates and locations; Old Town, ME (**A**); Durham, NH (**B**), Big Flats, NY (**C**), and Aurora, NY (**D**) (adapted from Cordeau et al., 2017b). Box plots (white) are presented within violin plots (black) and means represented by red dots.

2020 over the three locations representing each zone. If no start date was provided, we began the accumulation on January 1. One of the prediction models required additional information on the soil conditions (Archer et al., 2006). For this model we input

typical conditions for Northeast USA – chisel plowed loam following corn, with wet initial conditions, and adequate rainfall. Predicted dates for 50% (peak) emergence were graphed by species and model (**Figure 2**).

TABLE 1 | Comparison of peak emergence of target weeds from previous studies.

Species (Latin name)	Location	USDA plant hardiness zone*	Observed peak emergence	Reference
Annual bluegrass (<i>Poa annua</i> L.)	California, USA	9	November 5	Shem-Tov and Fennimore, 2003
	Illinois, USA	6	Spring and autumn	Branham, 1991
	Legnaro, Italy	8	Late February	Masin and Macolino, 2016
	New York, USA	6	June 23 and Aug. 4	Cordeau et al., 2017b
	Maryland, USA	7	Late September to mid October	Kaminski and Demooeden, 2007
	Scotland, UK	7	September	Lawson et al., 1974, as reported by Mohler, 2001
Barnyardgrass (<i>Echinochloa crus-galli</i> (L.) Beauv.)	Tennessee, USA	7	Mid October	Taylor et al., 2021
	Arkansas, USA	7	Late May	Bagavathiannan et al., 2011
	Central Czech Republic	6	May and June	Jursik et al., 2014
	Massachusetts, USA	6	June, but through September	Vengris, 1965
	New York, USA	6	June 23	Cordeau et al., 2017b
	Northern Greece	7	April 20	Vasileiadis et al., 2016
Canada thistle (<i>Cirsium arvense</i> (L.) Scop.)	Ontario, Canada	5	June	Maun and Barrett, 1986
	Idaho, USA	6	March and April	Hodgson, 1955
	Montana, USA	4	Early May	Hodgson, 1964
	New York, USA	6	April 28	Cordeau et al., 2017b
Common chickweed (<i>Stellaria media</i> (L.) Vill)	North Dakota, USA	4	Early May	Donald, 2000
	England, UK	8	Early spring or late fall	Roberts and Dawkins, 1967
	Maine, USA	5	July 7	Cordeau et al., 2017b
	New Hampshire, USA	6	July 21	Cordeau et al., 2017b
	New York, USA	6	Mid September	DiTommaso, 2016
Common lambsquarters (<i>Chenopodium album</i> L.)	Scotland, UK	7	August to October	Lawson et al., 1974, as reported by Mohler, 2001
	Central Czech Republic	6	March and April	Jursik et al., 2014
	Maine, USA	5	April 28	Cordeau et al., 2017b
	Mid-Atlantic USA	8	May 11 to June 1	Myers et al., 2004
	Minnesota, USA	4	Late April	Harvey and Forcella, 1993
	New York, USA	6	Mid May	DiTommaso, 2016
	Northern/central Italy	8	Late April	Masin et al., 2012
	Ontario, Canada	5	Early June	Roman et al., 2000
Common ragweed (<i>Ambrosia artemisiifolia</i> L.)	Quebec, Canada	4	May 30	Leblanc et al., 2004
	Scotland, UK	7	May	Lawson et al., 1974, as reported by Mohler, 2001
	Wisconsin, USA	4	June 11	Buhler et al., 1996
	Illinois, USA	6	April and May, none after June 1	Stoller and Wax, 1973
	Mid-Atlantic, USA	8	April 7 to May 1	Myers et al., 2004
	Nebraska, USA	5	April 28	Barnes et al., 2017
	New York, USA	6	Early May	DiTommaso, 2016
	New York, USA	6	May 12	Cordeau et al., 2017b
Giant foxtail (<i>Setaria faberi</i> Herm.)	New York, USA	6	Prior to June 9	Dickerson, 1968
	Ontario, Canada	5	90% emergence prior to June 15	Bassett and Crompton, 1975
	Mid-Atlantic, USA	8	Sites ranged May 2 to May 24	Myers et al., 2004
	New York, USA	6	April 28	Cordeau et al., 2017b
Hairy galinsoga (<i>Galinsoga quadriradiata</i> Cav.)	Ohio, USA	6	May 17	Cardina et al., 2007
	Wisconsin, USA	4	June 8	Buhler et al., 1996
	Central Czech Republic	6	June and July	Jursik et al., 2014
Large crabgrass (<i>Digitaria sanguinalis</i> (L.) Scop.)	Maine, USA	5	May 26	Cordeau et al., 2017b
	Multiple locations	-	May and June, but continues throughout growing season	Warwick and Sweet, 1983
	New York, USA	6	Early October (delayed due to drought)	DiTommaso, 2016
	Arkansas, USA	8	Spring	King and Oliver, 1994
	Legnaro, Italy	8	Mid to late May	Masin et al., 2005
	Mid-Atlantic, USA	8	Sites ranged May 18 to June 8	Myers et al., 2004
	New Hampshire, USA	6	April 28	Cordeau et al., 2017b
	New York, USA	6	April 28	Cordeau et al., 2017b
Ohio, USA	6	June 1	Cardina et al., 2011	

(Continued)

TABLE 1 | Continued

Species (Latin name)	Location	USDA plant hardiness zone*	Observed peak emergence	Reference
Mouseear chickweed (<i>Cerastium fontanum</i> Baumg.)	New Hampshire, USA	6	August 4	Cordeau et al., 2017b
Perennial sowthistle (<i>Sonchus arvensis</i> L.)	New York, USA	6	July 21 and August 18	Cordeau et al., 2017b
	Uppsala, Sweden	5	Late April (from rhizomes)	Håkansson, 1969
	USA and Canada	5	Late May (from seed)	Lemna and Messersmith, 1990
Quackgrass (<i>Elymus repens</i> (L.) Gould)	Canada and USA	5	Early spring (from seed)	Werner and Rioux, 1977
	England, UK	8	Spring and autumn	Williams, 1971
	Maine, USA	5	August 4 and September 1	Cordeau et al., 2017b
	New York, USA	6	June 9 and September 1	Cordeau et al., 2017b
	New Hampshire, USA	6	September 1	Cordeau et al., 2017b
Red sorrel (<i>Rumex acetosella</i> L.)	Victoria, Australia	9	Autumn (from seed)	Amor, 1985
Redroot pigweed (<i>Amaranthus retroflexus</i> L.)	Central Czech Republic	6	Late April and May	Jursík et al., 2014
	Maine, USA	5	July 7 and August 18	Cordeau et al., 2017b
	New York, USA	6	April 28	Cordeau et al., 2017b
	Wisconsin, USA	4	June 8	Buhler et al., 1996
Smooth crabgrass (<i>Digitaria ischaemum</i> Schreb. ex Muhl.)	Maryland, USA	8	Early June	Fidanza et al., 1996
	New Hampshire, USA	6	June 9	Cordeau et al., 2017b
	New York, USA	6	May 12	Cordeau et al., 2017b
	Ohio, USA	6	May 10	Cardina et al., 2011

*Approximated based on USDA (2012) for sites in USA and Magarey et al. (2008) for all other sites.

Many of these studies reported 50% cumulative emergence, which is not necessarily equivalent to peak emergence, but often represents the most rapid rate of emergence.

COMPARISON OF EMERGENCE PEAKS FROM PREVIOUS RESEARCH

The previous empirically-based emergence results (Table 1) were generally in agreement as to the season of peak emergence for a given species. But within such general trends, peak emergence varied by study, likely reflecting differences in growing-degree-day accumulation between sites.

Model-based predictions of peak emergence using weather data from Northeast USA (Figure 2) were in general agreement with empirically observed results. Spring-emerging species were predicted to emerge earlier in the warmer modelled zones. Within each species, the mean difference in predicted peak emergence between warmest and coolest modeled USDA Plant Hardiness Zones of Northeast USA was 39 days.

Results by Species

In this section, we will draw from empirical observations (Table 1) and model-based predictions (Figure 2) to summarize the typical emergence patterns of our 15 reviewed species. Although the climate of Northeast USA is primarily composed of USDA plant hardiness zones 4 through 7, our review found several previous results from zones 8 and 9, which we include for further context (Table 1).

Annual bluegrass generally exhibited peak emergence in autumn, but spring, summer, and even winter emergence peaks were also observed (Table 1, Figure 2). Emergence models predicted that warmer climates would experience earlier spring and later autumn emergence of annual bluegrass compared with colder climates (Figure 2). These results reflect the winter annual life cycle of

annual bluegrass. Indeed, Håkansson (2003) reports that it germinates whenever dormancy has been broken and environmental conditions are adequate.


Barnyardgrass demonstrated emergence peaks in April or May from studies conducted in warm climates (Table 1). But in cooler climates, a June emergence peak was more common. For example, modelling efforts based in Minnesota and Iowa, USA showed a mean peak emergence of June 1 (Figure 2). Though emergence may occur for an extended period (Vengris, 1965; Gołębiowska and Kieloch, 2016), barnyardgrass phenological traits differ by location, and cold winter conditions likely select against late and untimely emergence (Martinková et al., 2021).

Canada thistle typically exhibited emergence peaks in early to mid-spring in studies from northern USA (Table 1, Figure 2). Emergence is mostly from overwintering root stock rather than seed (Hodgson, 1964), which likely contributes to vigorous early growth and establishment success.

Common chickweed has shown emergence peaks in spring, summer, and autumn (Table 1, Figure 2), reflecting its relatively weak dormancy (Grundy et al., 2003). Likewise, its emergence can be protracted over most of the growing season (Figure 1).

Common lambsquarters demonstrated emergence peaks mostly in mid-spring (Table 1, Figure 2). Earlier or later emergence was generally observed in warmer or colder climates, respectively. Common lambsquarters has distinct dormancy-breaking requirements (Grundy et al., 2003), but emergence can be somewhat protracted (Figure 1, Gołębiowska and Kieloch, 2016).

Common ragweed was generally the earliest emerging of the annual weeds we reviewed, with peaks mostly in early to mid-spring (Table 1, Figure 2). In several studies emergence was observed to curtail sharply in June (Table 1, Figure 1).



Giant foxtail emergence peaks were mostly in mid-spring (**Table 1, Figure 2**). Cordeau et al. (2017b) observed that emergence occurred over a short period, resulting in a prominent emergence peak (**Figure 1**).

Hairy galinsoga demonstrated emergence peaks mostly in late spring and summer (**Table 1**), congruent with anecdotal results from Jernigan et al. (2017). Though hairy galinsoga has no seed dormancy (Warwick and Sweet, 1983), it does require adequate soil conditions for germination, as evident in the emergence peak observed in autumn following a summer drought (**Table 1**).

Large crabgrass emerged in mid to late spring (**Table 1, Figure 2**). Cordeau et al. (2017b) found that emergence peaked sharply at one site, but was slightly protracted at another (**Figure 1**). In both cases, emergence was earlier than expected based on other findings from similar latitudes. This could perhaps be explained by the six-week window between tillage and sampling, which may have skewed results.

Mouseear chickweed demonstrated a mid-summer emergence peak (**Table 1**), but only one study was found reporting emergence periodicity.

Perennial sowthistle emerged primarily in mid spring in two of three studies (**Table 1**), reflecting its initiation of rhizome shoot elongation at that time (Torssell et al., 2015). But Cordeau et al. (2017b) found an emergence peak in summer, which may possibly be explained by spring field preparation with a glyphosate application that may have set back early-emerging perennial sowthistle.

Quackgrass exhibited emergence peaks in spring, spring and autumn, or summer and autumn (**Table 1**). But based on the

experimental design of these studies, some of the summer and autumn emergence likely represents regrowth from earlier-emerged plants.


Red sorrel emergence, which is primarily from rhizomes, peaked in summer, whereas emergence from seed peaked in autumn (**Table 1, Figure 2**).

Redroot pigweed demonstrated a range of emergence peaks from spring to summer (**Table 1, Figure 2**). This protracted emergence pattern of *Amaranthus* species has been well-described in the literature (Jha and Norsworthy, 2009; Teasdale and Mirsky, 2015; Chahal et al., 2021; Reinhardt Piskackova et al., 2021).

Smooth crabgrass emergence peaks were primarily in May and June, similar to large crabgrass (**Table 1, Figure 2**).

MANAGEMENT RECOMMENDATIONS

In this section, we will provide a synthesis of management tactics that may be improved with knowledge of weed emergence. The suite of applicable management tactics depends on the season and pattern of emergence of the target weed species. For example, direct control of weed seedlings (**Figure 3A**), overlapping residual herbicides (**Figure 3B**), residue mulch (**Figure 3C**), and delayed planting (**Figure 3D**) are most applicable to early- and mid-season weeds. Early planting to improve crop competition (**Figure 3E**) is most applicable to mid- and late-season weeds. And feasibility of fallow or cover cropped periods (**Figures 3F, G**) may vary by season and



cropping system (Figure 4). Therefore, we present the following management recommendations by season of weed emergence. Within each season, management tactics are organized in ascending order of management flexibility required, in accordance with Figure 3. Although many of the studies of emergence-based

management tactics were conducted outside of Northeast USA, most were conducted in similarly temperate climates, and their conclusions are applicable to this review.

Weeds With Early-Season Peak Emergence

Many of the species we reviewed exhibit emergence peaks in the early part of the growing season – typically April and May (Table 1) – including barnyardgrass, Canada thistle, common lambsquarters, common ragweed, giant foxtail, large crabgrass, perennial sowthistle, and smooth crabgrass. These weeds are especially problematic since they have the potential to compete with spring-planted crops for the entire growing season.

Direct control of early-season weed seedlings with herbicides or cultivation should be guided by weed emergence to maximize effectiveness (Forcella, 1999; Zimdahl, 2013). Farmers relying on a single post-emergence herbicide application should time it well after peak weed emergence so that the greatest number of weeds are sprayed (Fidanza et al., 1996; Masin et al., 2005; Archer et al., 2006). But for early emerging summer annual weeds, like common lambsquarters, larger weeds can survive herbicide applications, so earlier timings may be more effective in such cases (DeGreeff et al., 2018). Likewise, weed seedling control using cultivation is most effective in the early “white thread” stage, especially for intra-row control (Gallandt et al., 2017), so efficiency of cultivation could be maximized if conducted around peak emergence periods of problem weeds.

Timing of early-season residual herbicide applications is also of great importance. A premature application may allow the chemical to break down prior to peak emergence, whereas a late application may not affect emerged weeds. Application and planting dates can be adjusted to optimize residual herbicide effectiveness (Culpepper et al., 2004; Webster et al., 2009; Rosario-Lebron et al., 2019). Though preplant applications of residual herbicides may not be needed if most emergence occurs prior to seedbed preparation, they may be used prior to stale or false seedbed periods to bolster seedbank depletion (Coleman et al., 2016).

Cover crop residue, or residue applied in the form of hay or straw mulch, can effectively reduce weed emergence *via* physical (Carrera et al., 2004) or allelopathic mechanisms (Weston and Duke, 2003). Timing of cover crop termination for creation of grown-in-place mulch may interact with many other factors to affect weed biomass following termination (Wayman et al., 2015). Teasdale and Mirsky (2015) noted that weeds emerging prior to termination have an advantage in this system, with giant foxtail becoming the dominant weed when a hairy vetch (*Vicia villosa* Roth.) cover crop was terminated with rolling, but in disk-killed hairy vetch, smooth pigweed (*Amaranthus hybridus* L.) eventually dominated, likely due to warmer soil temperatures caused by tillage, which may have stimulated germination of this later-emerging species. Roller-crimped cereal rye (*Secale cereale* L.) can provide effective weed control in organic soybean [*Glycine max* (L.) Merr.], but common ragweed often escapes suppression since it emerges before the rye is rolled into a




FIGURE 4 | Examples of crops that may be shifted to avoid of peak weed emergence periods occurring in the early (A, April-May), middle (B, June-July), or late (C, August-September) growing season of the cool, temperate climate of Northeast USA. Height of vertical bars represents the relative stage of crop growth.

suppressive mulch (Wallace et al., 2018). Thus, fields dominated by the earliest-emerging weeds should not rely on roller-crimped cover crops for weed control since they need to be terminated at anthesis, which is typically mid to late spring in Northeast USA. On small-scale farms, hay or straw mulch is sometimes used, and may be applied before planting robust crops or well after planting to ensure crops have achieved a sufficient height to withstand the mulching (Brown and Gallandt, 2018). When considering the best timing of mulch application relative to the peak emergence of problem weeds, farm managers should consider that it is most effective soon after application, before it decomposes (Law et al., 2006).

Delayed planting may be used to avoid peak emergence of problematic early-season weeds (Gill and Holmes, 1997). This would allow delayed usage of tillage or burndown herbicides to control most of the targeted species before the crop is planted (Chahal et al., 2021). Royo-Esnal et al. (2018) found that by delaying planting dates, they were able to avoid 50-100% of the emergence of their target weed. Common ragweed was less abundant in corn (*Zea mays* L.) and soybean crops when planting dates were delayed (Wallace et al., 2018). Delaying soybean planting date to June can decrease weed density at the time of post-emergence herbicide application (DeWerff et al., 2015). Delayed planting can also extend the activity and improve

the control provided by pre-emergence herbicides (Rosario-Lebron et al., 2019). The timing of crop planting can also be adjusted to avoid crop losses due to the timing of weed competition (Mohler, 2001; Knezevic et al., 2002). Delaying planting may reduce the yield potential in some crops, which must be weighed against weed management benefits. Since crops differ in planting dates and associated management (Liebman and Gallandt, 1997), planting date may factor into crop or variety choice. Crops that can be planted in late-spring to avoid early-season weeds include corn, cucurbits, dry beans (*Phaseolus* spp.), solanaceous crops, soybeans, and sunflowers (*Helianthus* spp.) (Figure 4). Alternatively, alfalfa (*Medicago sativa* L.) and winter cereals may be planted the previous year to establish a competitive advantage over weeds with early-season peak emergence.

Bare fallow periods, including false or stale seedbeds, are some of the most powerful methods to reduce the weed seedbank (Gallandt, 2006), especially if the timing corresponds with the emergence peak of targeted weed species. Fallow periods are perhaps most feasible during the spring since more than one tillage pass or burndown herbicide application is sometimes necessary to prepare for planting, and spacing out these events allows emergence flushes between control timings. Although it is best to finish the sequence with minimal disturbance, such as with herbicides or flaming (Caldwell and Mohler, 2001), initiating with shallow tillage set to the germination depth of the target weed species can enhance the depletion of the seedbank (De Cauwer et al., 2019). But while tillage may affect the emergence of some species, the timing and magnitude of common ragweed emergence has been shown to be unaffected (Barnes et al., 2017). In such cases, it would be best to allow sufficient time for emergence rather than expecting an early peak due to tillage.

Weeds With Mid-Season Peak Emergence

Weeds with mid-season emergence peaks in June or July, included hairy galinsoga, mouseear chickweed, and red sorrel. These species may not be exposed to early season weed control efforts. Furthermore, opportunities to control these mid-season weeds may be curtailed if canopy closure or crop height restrict tractor entry into fields. Likely for these reasons, mid-season weeds can be the most abundant weeds in corn crops (Fried et al., 2020).

Direct control of mid-season weed seedlings through post-emergence herbicides or cultivation may be possible until the “layby” stage of the crop, immediately prior to canopy closure. Use of overlapping residual herbicides at this stage would further reduce emergence until full crop canopy closure (Culpepper et al., 2004).

Early planting of spring-planted crops may be used to establish a size advantage over mid-season weeds that can be used to improve control (Mohler, 2001). This size advantage is especially important for keeping weeds as small as possible for late post-emergence herbicide applications (Reinhardt Piskackova et al., 2021) and improving the selectivity of intra-row cultivation (Gallandt et al., 2017). The effect of early planting

can perhaps be strengthened by additional weed suppressive tactics. Increasing crop planting density can be used to hasten canopy closure and facilitate mid-season shading (Mohler, 1996). Residue mulches (Brown and Gallandt, 2018; Wallace et al., 2018) or living mulches (Hartwig and Ammon, 2002; Westbrook et al., 2021) can provide further shading. Due to increased crop competition, late-emerging barnyardgrass and common lambsquarters had no effect on corn yield (Gołębiewska and Kieloch, 2016). Thus, planting earlier can perhaps be used to ensure mid-season weeds emerge after the critical period for weed control, thereby nullifying their threat – but only if seed production is prevented (Reinhardt Piskackova et al., 2020). For example, to reduce barnyardgrass seed production by 99%, rice crops required a 23-day advantage (Singh et al., 2017).

When planting dates were delayed, Wallace et al. (2018) found that composition of weeds with mid-season emergence periodicity increased in organic no-till corn and soybeans. They suggest integration of winter grains or perennial forages to reduce seedbanks of these species. Alternatively, in specialty crop systems, cool season crops that are established before or planted after peak emergence may avoid mid-season weeds entirely (DeVore et al., 2011). Such crops include beets, carrots, green peas, leafy greens, or lettuce (Figure 4).

Bare fallow periods targeting the seedbank of mid-season weeds may be difficult for farmers to implement in the middle of the growing season but could be integrated with cover cropping to simultaneously satisfy other pest management or soil health goals (Nordell and Nordell, 2009). Aligning the timing of cover crop seedbed preparation with the peak emergence of problem weeds would encourage weed seed germination, while termination of the cover crop prior to weed seed production would ensure a seedbank reduction (Figure 3G). Bare fallow periods can also be used in conjunction with cool season crops (Figure 4). Bare fallow periods in the mid to late season are especially important when targeting mid-season perennial weeds, such as quackgrass (Table 1), to exhaust the carbohydrate reserves in their rhizomes (Ringselle et al., 2020).

Weeds With Late-Season Peak Emergence

Weeds with peak emergence in late summer and autumn included the winter annual species, annual bluegrass. Weeds emerging at this timing provide minimal competition with established spring-planted crops (Knezevic et al., 2002; Cordeau et al., 2015; Gołębiewska and Kieloch, 2016). In interviews about problematic weeds, organic vegetable farmers made almost no mentions of winter annuals (Jabbour et al., 2014), reflecting the control of winter annuals by spring tillage. But late-season weeds can be problematic in winter cereals (Royo-Esnal et al., 2018). And in no-till systems, autumn-established winter annuals can grow too large the following spring to control with post-emergence herbicides (Wallace et al., 2019).

Autumn-applied burndown herbicides can reduce overwintering weed densities the following spring (Hasty et al., 2004). And autumn-applied residual herbicides in conjunction

with winter cover crops can reduce the emergence of winter annual weeds (Walters et al., 2007) and keep them at a manageable height the following spring (Wallace et al., 2019). Interseeding may be used to expedite cover crop establishment and improve late season weed suppression, but compatibility with other post-harvest weed control tactics may be limited.

As grain corn or soybean crops mature and leaves wither, the open canopy can allow successful establishment of weeds with late-season emergence peaks. Harvest of these crops often occurs in late autumn, forcing farm managers to decide whether to terminate the weeds in autumn, or use the weeds as winter soil cover (Jabbour et al., 2014) but risk their escape in spring (Wallace et al., 2018).

Shifting spring-planted crops earlier (Figure 4) can allow for a late-season bare fallow period, which would target the seedbank of winter annuals, and help exhaust carbohydrate reserves of perennial weeds to minimize their survival over winter (Andersson et al., 2013). Late fallow periods may also necessitate a delay in autumn-planted crops. This could perhaps be offset with denser seeding rates. Royo-Esnal et al. (2018) found that by delaying winter cereal planting dates, they were able to avoid most emergence of their target weed. But in a dry year, weed emergence was delayed, and an earlier planting date would have been preferable.

Weeds With Protracted Emergence

Rather than emerging in a short, easily manageable cohort, several species exhibited a protracted emergence pattern. These included common chickweed, quackgrass, and redroot pigweed. The protracted emergence of *Amaranthus* species is well-known (Jha and Norsworthy, 2009; Chahal et al., 2021). Common lambsquarters (Gołębiewska and Kieloch, 2016) and hairy galinsoga (Jernigan et al., 2017) may also sometimes display protracted emergence. This emergence pattern allows these species to avoid early-season control efforts. Although late emerging weeds may not affect crop yield (Knezevic et al., 2002), they should still be controlled from a seedbank management standpoint (Norris, 1999; Brown and Gallandt, 2018) and because they are subject to sublethal herbicide doses, which can select for resistance (Norsworthy et al., 2012). Perennial weeds may also appear to have a protracted emergence due to their survival and resprouting after successive control efforts (Ringselle et al., 2020).

Control of weeds with protracted emergence requires an extended effort. We suggest a multifaceted strategy that includes as many of the following tactics as is feasible. In chronological order: densely planting crops that quickly close canopy (Mohler, 1996); overlapping residual herbicides to extend control through canopy closure (Chahal et al., 2018); ensuring seedling control is as effective as possible by timing control efforts while weeds are less than 10 cm tall (Reinhardt Piskackova et al., 2020) and using a final pass as late as possible (Crow et al., 2015); minimizing seed production of escapes using hand removal, mowing, rescue herbicide applications (Hill et al., 2016), electrical discharge systems, or artificial pollination (Lidor-Nili and Noivirt-Brik, 2017); using harvest weed seed control if the escapes have already produced seed (Walsh et al., 2012); avoiding

post-harvest tillage to allow for seed predation (Birthisel et al., 2015); and moldboard plowing to bury remaining seeds below germinable depth (Mohler, 1996).

Management of Multiple Problem Weeds

As seen in Figure 1, farms often contain species with different weed emergence patterns. Therefore, farmers may need to prioritize targeting the emergence peaks of the most problematic species. Ideally, the most competitive or difficult-to-control weeds would be avoided entirely by adjusting planting dates or crop selection to allow for a bare fallow (Figure 3F) or short-term cover crop (Figure 3G). Lower priority weeds could be addressed with adjusted in-crop weed control (Figures 3A–C) combined with tactics that improve the competitive advantage of the crop (Figures 3D, E). Fields with profuse weed emergence over the entire growing season may need to be taken out of production for a full year to allow for extended bare fallow periods (Nordell and Nordell, 2009) or rapid succession cover crops that are incorporated before weeds have time to set seed (Sarrantonio and Gallandt, 2003; Gallandt, 2006; Mirsky et al., 2010).

Conversely, fields may have weed communities dominated by species with similar phenology, reflecting the history of management (Ryan et al., 2010). For example, an organic vegetable rotation and a conventional corn and alfalfa rotation resulted in two different weed communities with very different phenology (Mohler et al., 2018). Likewise, corn and soybean are more conducive to summer annual broadleaf weed emergence than wheat and alfalfa (Goplen et al., 2017). Therefore, in some cases it may be possible to target all of the problematic weeds by temporarily switching to a cropping system that allows for new management timings – such as the occasional rotation of spring versus autumn-planted crops.

OUTLOOK

Variation in Peak Emergence Within Each Species

From our review of previous research, the peak emergence of each species varied by study (Figure 1, Table 1). Indeed, for a given weed species, emergence timing is expected to differ by location due to different weather conditions (Royo-Esnal et al., 2020). But there was also variation between modelling results of a given species when we imposed identical weather conditions (Figure 2). This may be due to other differences between study sites where models were validated, such as soil texture (Leblanc et al., 2004), surface residue (Oreja et al., 2020), crop canopy (Cardina et al., 2011), or type of tillage employed (Mohler, 2001). Additionally, depth of burial can influence not only the time required by a seedling to reach the soil surface (Gonzalez-Andujar et al., 2016), but also the accumulation of conditions required to break seed or bud dormancy (Cao et al., 2011). Furthermore, within a species there may be genetic-based differences in emergence between populations resulting from varying selection pressure in different microclimates (Dorado

et al., 2009; Papiernik et al., 2020; Liu et al., 2021). For example, barnyardgrass emergence differed based on the elevation of the seed collection site (Martinková et al., 2021) and common lambsquarters emergence varied by the slope of the landscape where seeds developed (Papiernik et al., 2020).

While Grundy et al. (2003) used a single thermal time model to sufficiently predict cumulative emergence of common chickweed from different sites brought to a single location, similar common garden experiments have found differences in emergence periodicity between populations (Papiernik et al., 2020; Martinková et al., 2021). Likewise, for large crabgrass (Dorado et al., 2009) and annual ryegrass (*Lolium rigidum* Gaud., Sousa-Ortega et al., 2019), single models could accurately predict emergence at single but not multiple locations, suggesting that different models should be developed for different regions.

But models may only require minor adjustments to accurately predict emergence in new regions. For example, the Italian emergence model, AlertInf (Masin et al., 2012), was modified with the base temperature and water potential required for germination of barnyardgrass in Croatia, to effectively predict emergence in Croatian corn (Šoštarčić et al., 2021). To further improve emergence predictions, other site-specific factors may need to be incorporated into models to maximize their accuracy and utility to farmers (Royo-Esnal et al., 2020). Inclusion of real-time *in-situ* information may be useful as well, such as phenological indicators (Cardina et al., 2007; Cardina et al., 2011), degree of crust formation, shading provided by the crop, and weed emergence scouting data. Models may also need to incorporate an adjustment factor to account for genetic tendencies of some populations within a species to emerge earlier or later than predicted. Farmers may achieve a simpler version of such a process by comparing their observed emergence to model predicted emergence and adjusting the prediction accordingly in subsequent years.

Considerations for Future Modelling Efforts

Other improvements to emergence models may be gained by improving the parameter estimates through statistically independent sampling and reducing the time between sampling events (Gonzalez-Andujar et al., 2016). For example, work of Cordeau et al. (2017b) was based on statistically independent samples as each sample through the season was from a new plot. But sampling occurred six weeks after each plot was tilled, which was ideal for investigating cohort effects resulting from different tillage timings, but was too imprecise for emergence modeling.

For this review, we focused on peak emergence or 50% cumulative emergence. The two terms are similar since the rate of emergence is theoretically highest at 50% cumulative emergence (Royo-Esnal et al., 2020). But the terms are not interchangeable since the greatest empirically observed emergence peak may occur before or after 50% cumulative emergence if the distribution of emergence is skewed by drought or other random variables. In cumulative emergence

models, emergence peaks would be represented by large, predicted flushes of emergence resulting from favorable weather, which may or may not bring the cumulative total above 50%. Such predictions of large flushes may be just as useful for planning purposes as cumulative emergence because weed seedling control will need to be prioritized while the cohort remains small. Thus, farmer-oriented emergence models should allow for incorporation of weather forecasts to provide corresponding weed emergence forecasts.

We have made the case for managing weeds based on their emergence peaks, but accurate predictions of the start, end, or any other point on the emergence continuum can also be useful for management purposes. For example, to maximize efficiency, weed seedling controls with nearly 100% efficacy should not be conducted until most of the weeds have emerged (Fidanza et al., 1996; Masin et al., 2005; Archer et al., 2006), whereas controls that are only effective on small weeds should be initiated when relatively few have emerged (Oriade and Forcella, 1999).

Feasibility of Adjusting Management Based on Weed Emergence

We have presented a range of techniques to improve weed management based on knowledge of peak emergence (Figure 3). Certainly, slight adjustments to the timing of seedling control may be more feasible for many farmers than making drastic changes to the cropping system. But farmers are very responsive to adjusting their management based on the biology and timing of their most problematic weeds (Jabbour et al., 2014) and will sometimes go to extreme efforts to regain control. Since most of the tactics we present embody a “weed smarter, not harder” approach, they may be easily adopted by farmers.

For farmers unable to shift the timing of their planting, it may still be possible to alter the edaphic environment to shift weed emergence earlier or later. Prior to planting, weed emergence may be expedited through practices that increase soil warming, such as tillage (Travlos et al., 2020) or solarization. Timing of irrigation may also be used to alter weed emergence (Kanas et al., 2021). Conversely, delaying the emergence of problematic weeds may be achieved through limiting absorption of sunlight *via* surface residue management (Oreja et al., 2020) or crop canopy establishment.

The feasibility of emergence-based management may be impacted by other factors warranting future research. Such as the weed community shifts that may occur after long-term emergence-based management. Or changing weather patterns that will likely lengthen the growing season to the benefit of many weed species (Peters and Gerowitt, 2015). The increased rainfall expected in Northeast USA may decrease the number of days when soils are dry enough to support tractor-drawn equipment, thereby increasing the importance of access to ample equipment and labor to conduct timely weed control efforts (Birthisel et al., 2021). Likewise, timely control may be challenging on large-scale farms since it may require several days to cover the entire acreage. But perhaps fields or

microclimatic areas with consistently early weed emergence could be managed first, and so forth, to allow control timing to mirror emergence trends.

Overall, previous studies were generally in agreement as to the season of peak emergence for each species. Variation in observed results likely reflected the climatic conditions of each study site. But our use of identical weather data in previous models also showed variation in emergence trends within species, highlighting the importance of local edaphic conditions and population-level differences. Nonetheless, prediction accuracy will improve as models become more advanced (Royo-Esnal et al., 2020) and farmers can further improve prediction reliability based on their own ground truthing and adjustments to model predictions in subsequent years. Furthermore, even our broad characterizations of peak emergence for each species (i.e., “late spring”) may be used to improve management. Depending on management flexibility, knowledge of the season and pattern of weed emergence be used to improve the timing of control efforts, enhance crop competition, or expedite weed seedbank depletion.

AUTHOR CONTRIBUTIONS

BB conducted the literature search. All authors were involved in the interpretation of the results. BB, EG, and SC wrote the initial draft. All authors were involved in improving the subsequent

drafts. All authors contributed to the article and approved the submitted version.

FUNDING

This work was partially supported by a joint research and extension project funded by the Cornell University Agricultural Experiment Station (Hatch funds) and Cornell Cooperative Extension (Smith Lever funds) received from the National Institutes for Food and Agriculture (NIFA) U.S. Department of Agriculture (Project: 2013-14-425). This work was financially supported by the Maine Agriculture and Forestry Experiment Station, French ANR COSAC project (ANR-14-CE18-0007), and the Burgundy Region in France (FABER project Couv’Herbi). Additional funding was provided by the French program Investissements d’Avenir ANR PPR SPECIFICS project (ANR-20-PCPA-0008) and the Ecophyto Dephy EXPE ABC (Agroécologie en Bourgogne et Région centre). Additional funding was provided by the New York State Department of Agriculture and Markets.

ACKNOWLEDGMENTS

The authors acknowledge field assistance from Joseph Cannon, Paul Stachowski, Scott Morris, Meredith Ward, Christopher Pelzer, Jeff Liebert, Kristine Averill, and Nicholas Warren.

REFERENCES

- Amor, R. (1985). Seasonal Emergence of Weeds Typically Occurring in the Victorian Cereal Belt. *Plant Prot. Q.* 1, 18–20.
- Andersson, L., Boström, U., Forkman, J., Hakman, I., Liew, J., and Magnuski, E. (2013). Sprouting Capacity From Intact Root Systems of *Cirsium arvense* and *Sonchus arvensis* Decrease in Autumn. *Weed Res.* 53, 183–191. doi: 10.1111/wre.12013
- Archer, D., Forcella, F., Korth, A., Kuhn, A., Eklund, J., and Spokas, K. (2006). *WeedCast Version 4.0* (Morris, MN: USDA ARS).
- Bagavathiannan, M. V., Norsworthy, J. K., Smith, K. L., and Burgos, N. (2011). Seedbank Size and Emergence Pattern of Barnyardgrass (*Echinochloa crus-galli*) in Arkansas. *Weed Sci.* 59, 359–365. doi: 10.1614/WS-D-10-00149.1
- Barnes, E., Werle, R., Sandell, L., Lindquist, J., Knezevic, S., Sikkema, P., et al. (2017). Influence of Tillage on Common Ragweed (*Ambrosia artemisiifolia*) Emergence Pattern in Nebraska. *Weed Technol.* 31, 623–631. doi: 10.1017/wet.2017.38
- Bassett, I. J., and Crompton, C. W. (1975). The Biology of Canadian Weeds: 11. *Ambrosia artemisiifolia* L. And *A. psilostachya* DC. *Can. J. of Plant Sci.* 55, 463–476. doi: 10.4141/cjps75-072
- Bastiaans, L., Paolini, R., and Baumann, D. T. (2008). Focus on Ecological Weed Management: What is Hindering Adoption? *Weed Res.* 48, 481–491. doi: 10.1111/j.1365-3180.2008.00662.x
- Batlla, D., Malavert, C., Farnocchia, R. B. F., and Benech-Arnold, R. (2020). “Modelling Weed Seedbank Dormancy and Germination,” in *Decision Support Systems for Weed Management*. Eds. G. R. Chantre and J. L. González-Andújar New York: Springer. doi: 10.1007/978-3-030-44402-0_4
- Birthisel, S. K., Clements, R. S., and Gallandt, E. R. (2021). Review: How Will Climate Change Impact the ‘Many Little Hammers’ of Ecological Weed Management? *Weed Res.* 61, 327–341. doi: 10.1111/wre.12497
- Birthisel, S. K., Gallandt, E. R., Jabbour, R., and Drummond, F. A. (2015). Habitat and Time are More Important Predictors of Weed Seed Predation Than Space on a Diversified Vegetable Farm in Maine, USA. *Weed Sci.* 63, 916–927. doi: 10.1614/WS-D-15-00057.1
- Bourgeois, B., Munoz, F., Fried, G., Mahaut, L., Armengot, L., Denelle, P., et al. (2019). What Makes a Weed a Weed? A Large-Scale Evaluation of Arable Weeds Through a Functional Lens. *Am. J. Bot.* 106, 90–100. doi: 10.1002/ajb2.1213
- Branham, B. (1991). Dealing With *Poa annua*. *Golf Course Management.* 59, 46–60.
- Brown, B., and Gallandt, E. (2018). A Systems Comparison of Contrasting Organic Weed Management Strategies. *Weed Sci.* 66, 109–120. doi: 10.1017/wsc.2017.34
- Buhler, D., Mester, T., and Kohler, K. (1996). The Effect of Maize Residues and Tillage on Emergence of *Setaria faberi*, *Abutilon theophrasti*, *Amaranthus retroflexus* and *Chenopodium album*. *Weed Res.* 36, 153–165. doi: 10.1111/j.1365-3180.1996.tb01811.x
- Caldwell, B., and Mohler, C. L. (2001). Stale Seedbed Practices for Vegetable Production. *HortScience* 36, 703–705. doi: 10.21273/HORTSCI.36.4.703
- Cao, R., Francisco-Fernandez, M., Anand, A., Bastida, F., and Gonzalez-Andujar, J. L. (2011). Computing Statistical Indices for Hydrothermal Times Using Weed Emergence Data. *J. Agric. Sci.* 149, 701–712. doi: 10.1017/S002185961100030X
- Cardina, J., Herms, C. P., and Herms, D. A. (2011). Phenological Indicators for Emergence of Large and Smooth Crabgrass (*Digitaria sanguinalis* and *D. ischaemum*). *Weed Technol.* 25, 141–150. doi: 10.1614/WT-D-10-00034.1
- Cardina, J., Herms, C. P., Herms, D. A., and Forcella, F. (2007). Evaluating Phenological Indicators for Predicting Giant Foxtail (*Setaria faberi*) Emergence. *Weed Sci.* 55, 455–464. doi: 10.1614/WS-07-005.1
- Carrera, L. M., Abdul-Baki, A. A., and Teasdale, J. R. (2004). Cover Crop Management and Weed Suppression in No-Tillage Sweet Corn Production. *HortScience* 39, 1262–1266. doi: 10.21273/HORTSCI.39.6.1262

- Chahal, P., Barnes, E., and Jhala, A. (2021). Emergence Pattern of Palmer Amaranth (*Amaranthus palmeri*) Influenced by Tillage Timings and Residual Herbicides. *Weed Tech.* 35, 433–439. doi: 10.1017/wet.2020.136
- Chahal, P. S., Ganie, Z. A., and Jhala, A. J. (2018). Overlapping Residual Herbicides for Control of Photosystem II and 4-Hydroxyphenylpyruvate Dioxygenase (HPPD) Inhibitor-Resistant Palmer Amaranth (*Amaranthus palmeri* S. Watson) in Glyphosate-Resistant Maize. *Front. Plant Sci.* 8. doi: 10.3389/fpls.2017.02231
- Coleman, L. B., Chaudhari, S., Jennings, K. M., Schultheis, J. R., Meyers, S. L., and Monks, D. W. (2016). Evaluation of Herbicide Timings for Palmer Amaranth Control in a Stale Seedbed Sweetpotato Production System. *Weed Technol.* 30, 725–732. doi: 10.1614/WT-D-15-00133.1
- Cordeau, S., Guillemin, J.-P., Reibel, C., and Chauvel, B. (2015). Weed Species Differ in Their Ability to Emerge in No-Till Systems That Include Cover Crops. *Ann. Appl. Biol.* 166, 444–455. doi: 10.1111/aab.12195
- Cordeau, S., Smith, R. G., Gallandt, E. R., Brown, B., Salon, P., Ditommaso, A., et al. (2017a). Disentangling the Effects of Tillage Timing and Weather on Weed Community Assembly. *Agriculture* 7, 66. doi: 10.3390/agriculture7080066
- Cordeau, S., Smith, R. G., Gallandt, E. R., Brown, B., Salon, P., Ditommaso, A., et al. (2017b). How do Weeds Differ in Their Response to the Timing of Tillage? A Study of 61 Species Across the Northeastern United States. *Ann. Appl. Biol.* 171, 340–352. doi: 10.1111/aab.12377
- Cordeau, S., Smith, R. G., Gallandt, E. R., Brown, B., Salon, P., Ditommaso, A., et al. (2017c). Timing of Tillage as a Driver of Weed Communities. *Weed Sci.* 65, 504–514. doi: 10.1017/wsc.2017.26
- Crow, W. D., Steckel, L. E., Hayes, R. M., and Mueller, T. C. (2015). Evaluation of POST-Harvest Herbicide Applications for Seed Prevention of Glyphosate-Resistant Palmer Amaranth (*Amaranthus palmeri*). *Weed Technol.* 29, 405–411. doi: 10.1614/WT-D-14-00146.1
- Culpepper, A. S., Flanders, J., York, A. C., and Webster, T. M. (2004). Tropical Spiderwort (*Commelina benghalensis*) Control in Glyphosate-Resistant Cotton. *Weed Technol.* 18, 432–436. doi: 10.1614/WT-03-175R
- Davis, A. S., Clay, S., Cardina, J., Dille, A., Forcella, F., Lindquist, J., et al. (2013). Seed Burial Physical Environment Explains Departures From Regional Hydrothermal Model of Giant Ragweed (*Ambrosia trifida*) Seedling Emergence in US Midwest. *Weed Sci.* 61, 415–421. doi: 10.1614/WS-D-12-00139.1
- De Cauwer, B., De Cuyper, T., De Ryck, S., Delanote, L., Dewaele, K., Willekens, K., et al. (2019). Reduction in Field Emergence and Seedbank Density of *Galinsoga quadriradiata* and Other Weeds After Contrasting False Seedbed Strategies in Organic Vegetable Fields. *Weed Res.* 59, 265–278. doi: 10.1111/wre.12363
- DeGreeff, R. D., Varanasi, A. V., Dille, J. A., Peterson, D. E., and Jugulam, M. (2018). Influence of Plant Growth Stage and Temperature on Glyphosate Efficacy in Common Lambsquarters (*Chenopodium album*). *Weed Technol.* 32, 448–453. doi: 10.1017/wet.2018.38
- DeVore, J., Norsworthy, J., Johnson, D., Starkey, C., Wilson, M., and Griffith, G. (2011). “Palmer Amaranth Emergence as Influenced by Soybean Production System and Deep Tillage,” in *2011 Proceedings, Southern Weed Science Society*, San Juan, PR. 239.
- DeWerff, R. P., Conley, S. P., Colquhoun, J. B., and Davis, V. M. (2015). Weed Control in Soybean as Influenced by Residual Herbicide Use and Glyphosate-Application Timing Following Different Planting Dates. *Weed Technol.* 29, 71–81. doi: 10.1614/WT-D-14-00040.1
- Dickerson, C. T. J. (1968). *Studies on the Germination, Growth, Development, and Control of Common Ragweed (Ambrosia artemisiifolia L.)* (Ithaca (NY: Cornell University).
- DiTommaso, A. (2016). *Weed Emergence Patterns in Response to Disturbance on Two Soil Types* (New York State Integrated Pest Management). Available at: <https://hdl.handle.net/1813/52046> (Accessed 2 November 2021).
- Donald, W. W. (2000). A Degree-Day Model of *Cirsium arvense* Shoot Emergence From Adventitious Root Buds in Spring. *Weed Sci.* 48, 333–341. doi: 10.1614/0043-1745(2000)048[0333:ADDMOC]2.0.CO;2
- Dorado, J., Sousa, E., Calha, I., Gonzalez-Andujar, J. L., and Fernandez-Quintanilla, C. (2009). Predicting Weed Emergence in Maize Crops Under Two Contrasting Climatic Conditions. *Weed Res.* 49, 251–260. doi: 10.1111/j.1365-3180.2008.00690.x
- Fidanza, M., Dernoeden, P., and Zhang, M. (1996). Degree-Days for Predicting Smooth Crabgrass Emergence in Cool-Season Turfgrasses. *Crop Sci.* 36, 990–996. doi: 10.2135/cropsci1996.0011183X0036000400029x
- Forcella, F. (1999). *Estimating the Timing of Weed Emergence. Site Specific Management Guidelines* (Atlanta: Potash & Phosphate Institute).
- Forcella, F., Wilson, R. G., Dekker, J., Kremer, R. J., John, C., Randy, L. A., et al. (1997). Weed Seed Bank Emergence Across the Corn Belt. *Weed Sci.* 45, 67–76. doi: 10.1017/S0043174500092493
- Fried, G., Chauvel, B., Munoz, F., and Reboud, X. (2020). Which Traits Make Weeds More Successful in Maize Crops? Insights From a Three-Decade Monitoring in France. *Plants* 9, 40. doi: 10.3390/plants9010040
- Gallandt, E. R. (2006). How Can We Target the Weed Seedbank? *Weed Sci.* 54, 588–596. doi: 10.1614/WS-05-063R.1
- Gallandt, E. R., Brainard, D. C., and Brown, B. (2017). “Developments in Physical Weed Control,” in *Integrated Weed Management for Sustainable Agriculture*. Ed. R. L. Zimdahl (Sawston, UK: Burleigh Dodds Science Publishing), 261–283.
- Gibson, K., Johnson, W., and Hillger, D. (2006). Farmer Perceptions of Weed Problems in Corn and Soybean Rotation Systems. *Weed Technol.* 20, 751–755. doi: 10.1614/WT-05-089R.1
- Gill, G. S., and Holmes, J. E. (1997). Efficacy of Cultural Control Methods for Combating Herbicide-Resistant *Lolium rigidum*. *Pest Manage. Sci.* 51, 352–358. doi: 10.1002/(SICI)1096-9063(199711)51:3<352::AID-PS648>3.0.CO;2-M
- Golebiowska, H., and Kieloch, R. (2016). The Competitive Ability of *Chenopodium album* and *Echinochloa crus-galli* in Maize Crops Depending on the Time of Their Occurrence or Removal. *Acta Agrobot.* 69, 1688. doi: 10.5586/aa.1688
- Gonzalez-Andujar, J. L., Chantre, G. R., Morvillo, C., Blanco, A. M., and Forcella, F. (2016). Predicting Field Weed Emergence With Empirical Models and Soft Computing Techniques. *Weed Res.* 56, 415–423. doi: 10.1111/wre.12223
- Goplen, J. J., Sheaffer, C. C., Becker, R. L., Coulter, J. A., Breitenbach, F. R., Behnken, L. M., et al. (2017). Seedbank Depletion and Emergence Patterns of Giant Ragweed (*Ambrosia trifida*) in Minnesota Cropping Systems. *Weed Sci.* 65, 52–60. doi: 10.1614/WS-D-16-00084.1
- Grundy, A., Peters, N., Rasmussen, I. A., Hartmann, K., Sattin, M., Andersson, L., et al. (2003). Emergence of *Chenopodium album* and *Stellaria media* of Different Origins Under Different Climatic Conditions. *Weed Res.* 43, 163–176. doi: 10.1046/j.1365-3180.2003.00330.x
- Haj Seyed Hadi, M. R., and Petersonzalez-Andujar, J. L. (2009). Comparison of Fitting Weed Seedling Emergence Models With Nonlinear Regression and Genetic Algorithm. *Comput. Electr. Agric.* 65, 19–25. doi: 10.1016/j.compag.2008.07.005
- Håkansson, S. (1969). Experiments With *Onchus Arvensis* L.: I. Development and Growth, and the Response to Burial and Defoliation in Different Developmental Stages. *Uppsala Lantbrukshogsk Ann.* 35, 989–1030.
- Håkansson, S. (2003). *Weeds and Weed Management on Arable Land: An Ecological Approach* (Wallingford, UK: CABI).
- Hartwig, N. L., and Ammon, H. U. (2002). Cover Crops and Living Mulches. *Weed Sci.* 50, 688–699. doi: 10.1614/0043-1745(2002)050[0688:AIACCA]2.0.CO;2
- Harvey, S., and Forcella, F. (1993). Vernal Seedling Emergence Model for Common Lambsquarters (*Chenopodium album*). *Weed Sci.* 41, 309–316. doi: 10.1017/S0043174500076220
- Hasty, R. F., Sprague, C. L., and Hager, A. G. (2004). Weed Control With Fall and Early-Preplant Herbicide Applications in No-Till Soybean. *Weed Technol.* 18, 887–892. doi: 10.1614/WT-03-041R3
- Hill, E. C., Renner, K. A., and Sprague, C. L. (2014). Henbit (*Lamium amplexicaule*), Common Chickweed (*Stellaria media*), Shepherd’s-Purse (*Capsella bursa-pastoris*), and Field Pennycress (*Thlaspi arvense*): Fecundity, Seed Dispersal, Dormancy, and Emergence. *Weed Sci.* 62, 97–106. doi: 10.1614/WS-D-13-00074.1
- Hill, E., Renner, K., VanGessel, M., Bellinder, R., and Scott, B. (2016). Late-Season Weed Management to Stop Viable Weed Seed Production. *Weed Sci.* 64, 112–118. doi: 10.1614/WS-D-15-00096.1
- Hodgson, J. (1955). *Whiteweed and its Control* (Moscow, ID: Idaho Agricultural Extension Service Experiment), p.1–p.19.
- Hodgson, J. M. (1964). Variations in Ecotypes of Canada Thistle. *Weeds* 13, 167–171. doi: 10.2307/4040720
- Jabbour, R., Zwickle, S., Gallandt, E. R., McPhee, K. E., Wilson, R. S., and Doohan, D. (2014). Mental Models of Organic Weed Management: Comparison of New

- England US Farmer and Expert Models. *Renew. Agric. Food Syst.* 29, 319–333. doi: 10.1017/S1742170513000185
- Jernigan, A., Caldwell, B., Cordeau, S., DiTommaso, A., Drinkwater, L., Mohler, C., et al. (2017). Weed Abundance and Community Composition Following a Long-Term Organic Vegetable Cropping Systems Experiment. *Weed Sci.* 65, 639–649. doi: 10.1017/wsc.2017.33
- Jha, P., and Norsworthy, J. K. (2009). Soybean Canopy and Tillage Effects on Emergence of Palmer Amaranth (*Amaranthus palmeri*) From a Natural Seed Bank. *Weed Sci.* 57, 644–651. doi: 10.1614/WS-09-074.1
- Jursik, M., Holec, J., and Soukup, J. (2014). Field Emergence of Weeds During the Growing Season of Sugar Beet. *Listy Cukrovarnické a Reparské* 130, 166–172.
- Kaminski, J. E., and Dernoeden, P. H. (2007). Seasonal *Poa annua* L. Seedling Emergence Patterns in Maryland. *Crop Sci.* 47, 775–779. doi: 10.2135/cropsci2006.03.0191
- Kanatas, P., Gazoulis, I., and Travlos, I. (2021). Irrigation Timing as a Practice of Effective Weed Management in Established Alfalfa (*Medicago sativa* L.) Crop. *Agronomy* 11, 550. doi: 10.3390/agronomy11030550
- King, C. A., and Oliver, L. R. (1994). A Model for Predicting Large Crabgrass (*Digitaria sanguinalis*) Emergence as Influenced by Temperature and Water Potential. *Weed Sci.* 42, 561–567. doi: 10.1017/S0043174500076955
- Knezevic, S. Z., Evans, S. P., Blankenship, E. E., Van Acker, R. C., and Lindquist, J. L. (2002). Critical Period for Weed Control: The Concept and Data Analysis. *Weed Sci.* 50, 773–786. doi: 10.1614/0043-1745(2002)050[0773:CPFWCT]2.0.CO;2
- Law, D. M., Rowell, A. B., Snyder, J. C., and Williams, M. A. (2006). Weed Control Efficacy of Organic Mulches in Two Organically Managed Bell Pepper Production Systems. *Horttechnology* 16, 225–232. doi: 10.21273/HORTTECH.16.2.0225
- Lawson, H. M., Waister, P. D., and Stephens, R. J. (1974). Patterns of Emergence of Several Important Arable Weed Species. *Br. Crop Pr.* 9, 121–135.
- Leblanc, M. L., Cloutier, D. C., Stewart, K. A., and Hamel, C. (2004). Calibration and Validation of a Common Lambsquarters (*Chenopodium album*) Seedling Emergence Model. *Weed Sci.* 52, 61–66. doi: 10.1614/P2002-109
- Lemna, W. K., and Messersmith, C. G. (1990). The Biology of Canadian Weeds. 94. *Sonchus arvensis* L. *Can. J. Plant Sci.* 70, 509–532. doi: 10.4141/cjps90-060
- Lidor-Nili, E., and Noivirt-Brik, O. (2017). *Composition, Kits and Methods for Weed Control. US Patent 5,925,808* (Washington, DC: U.S. Patent and Trademark Office).
- Liebman, M., and Gallandt, E. (1997). “Many Little Hammers: Ecological Management of Crop-Weed Interactions,” in *Ecology in Agriculture*. Ed. L. Jackson (New York: Academic Press), 291–343.
- Liu, R., Kumar, V., Jha, P., and Stahlman, P. W. (2021). Emergence Pattern and Periodicity of Palmer Amaranth (*Amaranthus palmeri*) Populations From Southcentral Great Plains. *Weed Technol* 36:110–117. doi: 10.1017/wet.2021.81
- Magarey, R. D., Borcherdt, D. M., and Schlegel, J. W. (2008). Global Plant Hardiness Zones for Phytosanitary Risk Analysis. *Sci. Agric.* 65, 54–59. doi: 10.1590/S0103-90162008000700009
- Martinková, Z., Honěk, A., Pekár, S., and Leišova-Svobodová, L. (2021). Geographic Differentiation of Adaptive Phenological Traits of Barnyardgrass (*Echinochloa crus-galli*) Populations. *Weed Sci.* 69, 353–361. doi: 10.1017/wsc.2021.11
- Masin, R., Loddio, D., Benvenuti, S., Otto, S., and Zanin, G. (2012). Modeling Weed Emergence in Italian Maize Fields. *Weed Sci.* 60, 254–259. doi: 10.1614/WS-D-11-00124.1
- Masin, R., and Macolino, S. (2016). Seedling Emergence and Establishment of Annual Ryegrass (*Poa annua*) in Turfgrasses of Traditional and Creeping Perennial Ryegrass Cultivars. *Weed Technol.* 30, 238–245. doi: 10.1614/WT-D-15-00070.1
- Masin, R., Zuin, M. C., Archer, D. W., Forcella, F., and Zanin, G. (2005). WeedTurf: A Predictive Model to Aid Control of Annual Summer Weeds in Turf. *Weed Sci.* 53, 193–201. doi: 10.1614/WS-04-066R1
- Maun, M. A., and Barrett, S. C. H. (1986). The Biology of Canadian Weeds. 77. *Echinochloa crus-galli* (L.) Beauv. *Can. J. Plant Sci.* 66, 739–759. doi: 10.4141/cjps86-093
- Mirsky, S. B., Gallandt, E. R., Mortensen, D. A., Curran, W. S., and Shumway, D. L. (2010). Reducing the Germinable Weed Seedbank With Soil Disturbance and Cover Crops. *Weed Res.* 50, 341–352. doi: 10.1111/j.1365-3180.2010.00792.x
- Mohler, C. L. “Weed Life History: Identifying Vulnerabilities.” in *Ecological Management of Agricultural Weeds*. Eds. M. Liebman, C. L. Mohler and C. P. Staver (2001) (Cambridge, UK: Cambridge University Press), 40–98.
- Mohler, C. L. (1996). Ecological Bases for the Cultural Control of Annual Weeds. *J. Prod. Agric.* 9, 468–474. doi: 10.2134/jpa1996.0468
- Mohler, C. L., Caldwell, B. A., Marschner, C. A., Cordeau, S., Maqsood, Q., Ryan, M. R., et al. (2018). Weed Seedbank and Weed Biomass Dynamics in a Long-Term Organic Vegetable Cropping Systems Experiment. *Weed Sci.* 66, 611–626. doi: 10.1017/wsc.2018.52
- Myers, M. W., Curran, W. S., Vangessel, M. J., Calvin, D. D., Mortensen, D. A., Majek, B. A., et al. (2004). Predicting Weed Emergence for Eight Annual Species in the Northeastern United States. *Weed Sci.* 52, 913–919. doi: 10.1614/WS-04-025R
- Nordell, A., and Nordell, E. (2009). Weed the Soil, Not the Crop. *Acres U.S.A.* 40, 21–28.
- Norris, R. F. (1999). Ecological Implications of Using Thresholds for Weed Management. *J. Crop Prod.* 2, 31–58. doi: 10.1300/J144v02n01_03
- Norsworthy, J. K., Ward, S. M., Shaw, D. R., Llewellyn, R. S., Nichols, R. L., Webster, T. M., et al. (2012). Reducing the Risks of Herbicide Resistance: Best Management Practices and Recommendations. *Weed Sci.* 60, 31–62. doi: 10.1614/WS-D-11-00155.1
- Oreja, F. H., Batlla, D., and de la Fuente, E. B. (2020). *Digitaria sanguinalis* Seed Dormancy Release and Seedling Emergence are Affected by Crop Canopy and Stubble. *Weed Res.* 60, 111–120. doi: 10.1111/wre.12392
- Oriade, C., and Forcella, F. (1999). Maximizing Efficacy and Economics of Mechanical Weed Control in Row Crops Through Forecasts of Weed Emergence. *J. Crop Prod.* 2, 189–205. doi: 10.1300/9785537
- Papiernik, S. K., Forcella, F., and Amundson, G. B. (2020). Emergence of Common Lambsquarters (*Chenopodium album* L.) is Influenced by the Landscape Position in Which Seeds Developed. *Agric. Environ. Lett.* 5, e20000. doi: 10.1002/ael2.20000
- Peters, K., and Gerowitz, B. (2015). Weed Growth Properties of *Amaranthus retroflexus*, *Echinochloa crus-galli* and *Setaria viridis* as Influenced by Shifts in the Maize Cropping Season. *J. Plant Dis. Prot.* 122, 49–55. doi: 10.1007/BF03356530
- Reinhardt Piskackova, T. A., Reberg-Horton, S. C., Richardson, R. J., Jennings, K. M., Franca, L., Young, B. G., et al. (2021). Windows of Action for Controlling Palmer Amaranth (*Amaranthus palmeri*) Using Emergence and Phenology Models. *Weed Res.* 61, 188–198. doi: 10.1111/wre.12470
- Reinhardt Piskackova, T. A., Reberg-Horton, C., Richardson, R. J., Jennings, K. M., and Leon, R. G. (2020). Integrating Emergence and Phenology Models to Determine Windows of Action for Weed Control: A Case Study Using *Senna obtusifolia*. *Field Crops Res* 258, e107959. doi: 10.1016/j.fcr.2020.107959
- Renner, K. A., Swinton, S. M., and Kells, J. J. (1999). Adaptation and Evaluation of the WEEDSIM Weed Management Model for Michigan. *Weed Sci.* 47, 338–348. doi: 10.1017/S0043174500091876
- Ringselle, B., De Cauwer, B., Salonen, J., and Soukup, J. (2020). A Review of non-Chemical Management of Couch Grass (*Elymus repens*). *Agronomy* 10, e11178. doi: 10.3390/agronomy10081178
- Roberts, H. A., and Dawkins, P. A. (1967). Effect of Cultivation on the Numbers of Viable Weed Seeds in Soil. *Weed Res.* 7, 290–301. doi: 10.1111/j.1365-3180.1967.tb01384.x
- Roman, E. S., Murphy, S. D., and Swanton, C. J. (2000). Simulation of *Chenopodium album* Seedling Emergence. *Weed Sci.* 48, 217–224. doi: 10.1614/0043-1745(2000)048[0217:SOCASE]2.0.CO;2
- Rosario-Lebron, A., Leslie, A. W., Yurchak, V. L., Chen, G., and Hooks, C. R. R. (2019). Can Winter Cover Crop Termination Practices Impact Weed Suppression, Soil Moisture, and Yield in No-Till Soybean [*Glycine max* (L.) Merr.]? *Crop Prot.* 116, 132–141. doi: 10.1016/j.cropro.2018.10.020
- Royo-Esnal, A., Recasens, J., Garrido, J., and Torra, J. (2018). Rigput Brome (*Bromus diandrus* Roth.) Management in a No-Till Field in Spain. *Agronomy* 8, e251. doi: 10.3390/agronomy8110251
- Royo-Esnal, A., Torra, J., and Chantre, G. R. (2020). “Weed Emergence Models,” in *Decision Support Systems for Weed Management*. Eds. G. R. Chantre and J. L. González-Andújar (Cham, Switzerland: Springer), 85–116.
- Ryan, M. R., Smith, R. G., Mirsky, S. B., Mortensen, D. A., and Seidel, R. (2010). Management Filters and Species Traits: Weed Community Assembly in Long-

- Term Organic and Conventional Systems. *Weed Sci.* 58, 265–277. doi: 10.1614/WS-D-09-00054.1
- Sarrantonio, M., and Gallandt, E. (2003). The Role of Cover Crops in North American Cropping Systems. *J. Crop Prod.* 8, 53–74. doi: 10.1300/J144v08n01_04
- Scherer, A., Melander, B., Jensen, P. K., Kudsk, P., and Avila, L. A. (2017). Germination of Winter Annual Grass Weeds Under a Range of Temperatures and Water Potentials. *Weed Sci.* 65, 468–478. doi: 10.1017/wsc.2017.7
- Shem-Tov, S., and Fennimore, S. A. (2003). Seasonal Changes in Annual Bluegrass (*Poa annua*) Germinability and Emergence. *Weed Sci.* 51, 690–695. doi: 10.1614/0043-1745(2003)051[0690:SCIABP]2.0.CO;2
- Singh, M., Bhullar, M. S., and Chauhan, B. S. (2017). Relative Time of Weed and Crop Emergence is Crucial for Managing Weed Seed Production: A Study Under an Aerobic Rice System. *Crop Prot.* 99, 33–38. doi: 10.1016/j.cropro.2017.05.013
- Šoštarić, V., Masin, R., Loddo, D., Svečnjak, Z., Rubinić, V., and Ščepanović, M. (2021). Predicting the Emergence of *Echinochloa crus-galli* (L.) P. Beauv. In Maize Crop in Croatia With Hydrothermal Model. *Agronomy* 11, e2072. doi: 10.3390/agronomy11102072
- Sousa-Ortega, C., Chamber, E., Urbano, J. M., Izquierdo, J., Loureiro, I., Mari, A. I., et al. (2019). Should Emergence Models for *Lolium rigidum* be Changed Throughout Climatic Conditions? The Case of Spain. *Crop Prot* 68, 268–277. doi: 10.1016/j.cropro.2019.105012
- Sousa-Ortega, C., Royo-Esnal, A., DiTommaso, A., Izquierdo, J., Loureiro, I., Mari, A. I., et al. (2020). Modeling the Emergence of North African Knapweed (*Centaurea diluta*), an Increasingly Troublesome Weed in Spain. *Weed Sci.* 68, 268–277. doi: 10.1017/wsc.2020.22
- Stoller, E. W., and Wax, L. M. (1973). Periodicity of Germination and Emergence of Some Annual Weeds. *Weed Sci.* 21, 574–580. doi: 10.1017/S0043174500032513
- Taylor, D. R., Prorock, M., Horvath, B. J., and Brosnan, J. T. (2021). Modeling Seasonal Emergence of *Poa annua* in Urban Greenspace. *Sci. Rep.* 11, e18960. doi: 10.1038/s41598-021-98525-4
- Teasdale, J., and Mirsky, S. (2015). Tillage and Planting Date Effects on Weed Dormancy, Emergence, and Early Growth in Organic Corn. *Weed Sci.* 63, 477–490. doi: 10.1614/WS-D-14-00112.1
- Tiwari, R., Reinhardt Piskackova, T. A., Devkota, P., Mulvaney, M. J., Ferrell, J. A., and Leon, R. G. (2021). Emergence Patterns of Winter and Summer Annual Weeds in Ethiopian Mustard (*Brassica carinata*) Cropping System. *Weed Sci.* 69, 446–453. doi: 10.1017/wsc.2021.20
- Torrsell, B., Eckersten, H., Anbari, S., Lundkvist, A., and Verwijst, T. (2015). Modelling Below-Ground Shoot Elongation and Emergence Time of *Sonchus Arvensis* Shoots. *Acta Agriculturae Scandinavica.* 65, 582–588. doi: 10.1080/09064710.2015.1044463
- Travlos, I., Gazoulis, I., Kanatas, P., Tsekoura, A., Zannopoulos, S., and Papastilianou, P. (2020). Key Factors Affecting Weed Seeds' Germination, Weed Emergence, and Their Possible Role for the Efficacy of False Seedbed Technique as Weed Management Practice. *Front. Agron.* 2. doi: 10.3389/fagro.2020.00001
- USDA (2012) *USDA Plant Hardiness Zone Map* (Agricultural Research Service, U.S. Department of Agriculture). Available at: <http://planthardiness.ars.usda.gov> (Accessed 10 November 2021).
- Vasileiadis, V., Froud-Williams, R., Loddo, D., and Eleftherohorinos, I. (2016). Emergence Dynamics of Barnyardgrass and Jimsonweed From Two Depths When Switching From Conventional to Reduced and No-Till Conditions. *Span. J. Agric. Res.* 14, e1002. doi: 10.5424/sjar/2016141-8752
- Vengris, J. (1965). Seasonal Occurrence of Barnyardgrass in Potato Fields in Massachusetts. *Weeds* 13, 374–375. doi: 10.2307/4040902
- Wallace, J., Curran, W., and Mortensen, D. (2019). Cover Crop Effects on Horseweed (*Erigeron canadensis*) Density and Size Inequality at the Time of Herbicide Exposure. *Weed Sci.* 67, 327–338. doi: 10.1017/wsc.2019.3
- Wallace, J. M., Keene, C. L., Curran, W., Mirsky, S., Ryan, M. R., and VanGessel, M. J. (2018). Integrated Weed Management Strategies in Cover Crop-Based, Organic Rotational No-Till Corn and Soybean in the Mid-Atlantic Region. *Weed Sci.* 66, 94–108. doi: 10.1017/wsc.2017.53
- Walsh, M. J., Harrington, R. B., and Powles, S. B. (2012). Harrington Seed Destructor: A New Nonchemical Weed Control Tool for Global Grain Crops. *Crop Sci.* 52, 1343–1347. doi: 10.2135/cropsci2011.11.0608
- Walters, S. A., Young, B. G., and Nolte, S. A. (2007). Cover Crop and Pre-Emergence Herbicide Combinations in No-Tillage Fresh Market Cucumber Production. *J. Sustain. Agric.* 30, 5–19. doi: 10.1300/J064v30n03_03
- Warwick, S. I., and Sweet, R. D. (1983). The Biology of Canadian Weeds. 58. *Galinsoga parviflora* and *Galinsoga quadriradiata* (= *G. Ciliata*). *Can. J. Plant Sci.* 63, 695–709. doi: 10.4141/cjps83-087
- Wayman, S., Cogger, C., Benedict, C., Burke, I., Collins, D., and Bary, A. (2015). The Influence of Cover Crop Variety, Termination Timing and Termination Method on Mulch, Weed Cover and Soil Nitrate in Reduced-Tillage Organic Systems. *Renew. Agric. Food Syst.* 30, 450–460. doi: 10.1017/S1742170514000246
- Webster, T. M., Grey, T. L., Flanders, J. T., and Culpepper, A. S. (2009). Cotton Planting Date Affects the Critical Period of Benghal Dayflower (*Commelina benghalensis*) Control. *Weed Sci.* 57, 81–86. doi: 10.1614/WS-08-118.1
- Werle, R., Sandell, L. D., Buhler, D. D., Hartzler, R. G., and Lindquist, J. L. (2014). Predicting Emergence of 23 Summer Annual Weed Species. *Weed Sci.* 62, 267–279. doi: 10.1614/WS-D-13-00116.1
- Werner, P. A., and Rioux, R. (1977). The Biology of Canadian Weeds. 24. *Agropyron repens* (L.) Beauv. *Can. J. Plant Sci.* 57, 905–919. doi: 10.4141/cjps77-130
- Westbrook, A. S., Bhaskar, V., and DiTommaso, A. (2021). Weed Control and Community Composition in Living Mulch Systems. *Weed Res* 62, 12–23. doi: 10.1111/wre.12511
- Weston, L. A., and Duke, S. O. (2003). Weed and Crop Allelopathy. *Crit. Rev. Plant Sci.* 22, 367–389. doi: 10.1080/713610861
- White, S. N., Boyd, N. S., and Van Acker, R. C. (2015). Temperature Thresholds and Growing-Degree-Day Models for Red Sorrel (*Rumex acetosella*) Ramet Sprouting, Emergence, and Flowering in Wild Blueberry. *Weed Sci.* 63, 254–263. doi: 10.1614/WS-D-14-00048.1
- Williams, E. (1971). Germination of Seeds and Emergence of Seedlings of *Agropyron repens* (L.) Beauv. *Weed Res.* 11, 171–181. doi: 10.1111/j.1365-3180.1971.tb00993.x
- Zimdahl, R. L. (2013). *Fundamentals of Weed Science* (Boston: Elsevier/Academic Press).

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Brown, Gallandt, DiTommaso, Salon, Smith, Ryan and Cordeau. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.