



HAL
open science

Phenotyping of soybean phenology to temperature and photoperiod

Pierre Maury, Béatrice Quinquiry, Dorian Bourgeois, Neila Ait-Kaci, Lin Kang, Céline Schoving, Philippe Debaeke

► **To cite this version:**

Pierre Maury, Béatrice Quinquiry, Dorian Bourgeois, Neila Ait-Kaci, Lin Kang, et al.. Phenotyping of soybean phenology to temperature and photoperiod. Johann Vollmann, Marjana Vasiljević, Leopold Rittler, Jegor Miladinović, Donal Murphy-Bokern. 11th World Soybean Research Conference (WSRC11), Jun 2023, Vienna, Austria. , pp.416, 2023, Soybean Research for Sustainable Development. Abstracts of the World Soybean Research Conference 11 (WSRC 11), 18- 23 June 2023, Vienna, Austria. 10.5281/zenodo.7974681 . hal-04165324

HAL Id: hal-04165324

<https://hal.inrae.fr/hal-04165324>

Submitted on 18 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

P. Maury¹, B. Quinquiry¹, D. Bourgeois², N. Ait-Kaci¹,
L. Kang², C. Schoving^{2,3}, P. Debaeke²

Background & Aims



Improving soybean production in Europe under climate change needs a good prediction of cultivar phenology under different temperature and photoperiod conditions. For that purpose, a **Simple Phenology Algorithm (SPA)** was developed (Schoving et al., 2020). Before being applied, SPA requires the calibration of 7 genotypic parameters. Therefore, a **simple phenotyping method** was designed and set up under natural and controlled conditions to determine these parameters.

SPA algorithm

- Formalism of development rate (Rdev)
- 7 genotypic parameters

$$R_{dev} = R_{dev,max} \cdot f(T) \cdot f(P)$$

- 3 parameters for the **thermal function F(T)**: minimum T0, optimum Topt, and maximum Tmax
- 3 parameters for the **photoperiodic function F(P)**: optimal and critical daylength for development, and S sensitivity coefficient
- 1 parameter for the **maximal development rate**: optimal Physiological Development Days to complete a given phenological phase (PDD opt)


Phenotyping of F(T) parameters

- Experiment in controlled conditions
- Determination of the rate and speed of seed germination

Climatic chambers



4 x 25 seeds ; 11 temperatures



Fitting according to Yin et al (1995)

Mean cardinal temperatures (T0, Topt, and Tmax) for germination were ca. 2.6, 28.8, and 38.7°C, respectively with significant differences among cultivars (GM 000 to II)

Phenotyping /optimization of F(P) parameters

- Experiment in natural climatic conditions
- Recording of main phenological stages (VE, R1, R5, R7, R8)


Heliaphen platform (10 and 6 cultivars in 2017 and 2022)



4 to 6 planting dates (from March to September)

Observed day of appearance of a main stage
Daily or hourly air temperatures were tested as input data for parameters optimisation

SPA Calibration (e.g. 6 cultivars - 2022)




S and PDD opt were set by optimisation, i.e. minimize the RMSE between predicted and observed day of R8 stage for all planting dates of a given cultivar


SPA Evaluation

- Multi-environment trials: SW France and France levels
- hourly or daily temperatures for PDD optimisation

2013-2018 (4 years) – SW France



2010-2021 (12 years) – France




Observed (OBS) and predicted (SIM) days after planting for maturity (R7) in SW France trials (A) days of year for maturity R8 of Post-registration trials at France level (B).

Indicators of performance of SPA during its field evaluation

Multi-environment trials/temperatures input for optimisation	MBE (days)	RMSE (days)	RRMSE (%)
SW France (R7 prediction)	-0,70	5,61	5
France level (R8 prediction)			
Daily temperature	12,50	23,50	16
Hourly temperature	-2,40	19,70	14

SPA Application

e.g. Areas suitable for a soybean cultivar x sowing date x climate



Conclusion & perspectives

A simple cultivar phenotyping method was designed and applied on a first set of 15 cultivars commonly grown in France. The optimization of the PDD plant parameter from hourly temperature values instead of daily values improved the prediction of phenology by SPA. This approach - simple cultivar phenotyping and photothermal algorithm - is currently evaluated on several grain legumes to design more diversified and agroecological cropping systems.

UMR AGIR – INRAE Toulouse,
24 chemin de Borde-Rouge
31320 Castanet-Tolosan
<https://www6.toulouse.inrae.fr/agir>

References

Schoving C et al., 2020. *Frontiers in Plant Science* 10: 1755

¹ INP-ENSAT, Université de Toulouse, UMR AGIR, France

² INRAE, Université de Toulouse, UMR AGIR, Castanet-Tolosan, France

³ Terres Inovia, Baziège, France



Contacts :
pierre.maury@ensat.fr
philippe.debaeke@inrae.fr



ECODIV Project

