



HAL
open science

Long-term regulation of fat sensing in rainbow trout (*Oncorhynchus mykiss*) fed a vegetable diet from the first feeding: focus on free fatty acid receptors and their signaling

Elodie Baranek, Cécile Heraud, Laurence Larroquet, Anne Surget, Anthony Lanuque, Frédéric Terrier, Sandrine Skiba-Cassy, Jérôme Roy

► To cite this version:

Elodie Baranek, Cécile Heraud, Laurence Larroquet, Anne Surget, Anthony Lanuque, et al.. Long-term regulation of fat sensing in rainbow trout (*Oncorhynchus mykiss*) fed a vegetable diet from the first feeding: focus on free fatty acid receptors and their signaling. *British Journal of Nutrition*, 2024, 131 (1), pp.1-16. <10.1017/S0007114523001599>. <hal-04166623>

HAL Id: hal-04166623

<https://hal.inrae.fr/hal-04166623v1>

Submitted on 20 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Long-term regulation of fat sensing in rainbow trout (*Oncorhynchus mykiss*) fed a vegetable diet from the first feeding: focus on free fatty acid receptors and their signaling

Elodie Baranek, Cécile Heraud, Laurence Larroquet, Anne Surget, Anthony Lanuque, Frederic Terrier, Sandrine Skiba-Cassy and Jérôme Roy

INRAE, Université de Pau et des Pays de l'Adour, E2S UPPA, UMR1419 Nutrition Metabolism and Aquaculture, Aquapôle, F-64310 Saint-Pée-sur-Nivelle, France.

Corresponding author: Jérôme Roy, Phone: +33 5 59 51 59 65, E-mail: jerome.roy@inrae.fr

ACKNOWLEDGMENTS: Authors are very grateful to F. Sandres, who took care of and fed the fish during the trial. The authors also thank K. Dias and V. Véron for their help in training in western blotting and enzyme activity analysis, respectively.

FUNDING: This research was funded by the French National Research Agency (grant no. ANR-21-CE20-0009-01 “FEEDOMEGA”).

DATA AVAILABILITY STATEMENT: The data are available on request from the corresponding author.

COMPETING INTERESTS: The authors declare no competing or financial interests.



This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI

10.1017/S0007114523001599

The British Journal of Nutrition is published by Cambridge University Press on behalf of The Nutrition Society

ABSTRACT

Taste plays a fundamental role in an animal's ability to detect nutrients and transmits key dietary information to the brain, which is crucial for its growth and survival. Providing alternative terrestrial ingredients early in feeding influences the growth of rainbow trout (RT, *Oncorhynchus mykiss*). Thus, the present study aimed to assess the influence, *via* long-term feeding (for 8 months from the first feeding), of alternative plant ingredients (V diet for vegetable diet *vs.* C diet for a control diet) in RT on the mechanism of fat sensing at the gustatory level. After the feeding trial, we studied the pathways of the fat sensing mechanism in tongue tissue and the integrated response in the central nervous system. To this end, we analyzed the expression pattern of free fatty acid receptors (*ffar*) 1 and 2, markers of calcium-signaling pathways (phospholipase C β , Orai, Stim or Serca), the serotonin level (a key neurotransmitter in taste buds), and the expression pattern of appetite-regulating neuropeptides in the hypothalamus (central area of appetite regulation). The results revealed that the V diet modified the expression pattern of *ffar1* and paralogs of *ffar2* genes in tongue tissue, along with differential regulation of calcium-signaling pathways and a defect in serotonin level and brain turnover, without influencing neuropeptide expression. This study is the first to support that changes in growth performance and food intake of RT fed a V diet could be due to the influence of differences in nutrient sensing and a decrease in hedonic sensation. We revealed that RT have similar fat-detection mechanisms as mammals.

KEYWORDS: first feeding; food sensing; alternative aquafeed; taste receptor; farmed fish

INTRODUCTION

As production of fish for human consumption continues to increase due to higher demand, the expansion of aquaculture has increased the demand for aquafeed. To optimize fish growth, aquaculture relies on feeds that meet the nutritional requirements of the fish while also being as inexpensive as possible. The availability of traditional aquafeed ingredients, especially fishmeal (FM) and fish oil (FO), has not increased with demand ⁽¹⁾. Thus, to address this growth and limit environmental impacts of aquaculture, FM and FO must be replaced with renewable, eco-friendly, and less expensive ingredients ⁽²⁾. For production of rainbow trout (RT, *Oncorhynchus mykiss*), the total replacement of FM and FO by alternative plant-products negatively affect the feed utilization of fish, it reduces survival and growth rates compared to commercial diets, due to changes in feeding behavior among other factors ⁽³⁾. This kind of replacement changes the fatty acid (FA) profile. FM and FO are rich in ω -3 long-chain polyunsaturated FAs (ω -3 LC-PUFAs), such as eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), which are not found in plants. Due to their textural properties, dietary lipids such as ω -3 LC-PUFAs can regulate feeding behavior and influence hedonic sensation or pleasure in mammals, thus specifically influencing the regulation of feeding behavior after a meal ⁽⁴⁻⁶⁾. Thus, the absence of particular FAs in plant products could influence feed intake (FI) regulation in fish.

Nutrient sensing plays an essential role in the selection, evaluation, and intake of foods by vertebrates ⁽⁷⁾. This sensory system can detect nutrients, which influences the regulation of feeding behavior ⁽⁸⁾. The perception and discrimination of palatable nutrients by the central nervous system is initiated by the activation of specific taste receptors in taste bud cells (TBC). The mechanisms of taste sensation in mammals is well documented ⁽⁹⁾. The gustatory system distinguishes five fundamental taste qualities: sweet, salty, umami, sour, and bitter. However, when it comes to detecting fats, the reliance shifts towards factors like texture, olfaction, and oral irritation rather than only the sense of taste ⁽¹⁰⁾. Free FAs are detected after cleavage of triglycerides by lingual lipase ⁽⁹⁾ due to the activation of specific G protein-coupled receptors (GPCR) called “free FA receptors” (FFARs). Depending on the length of their carbon chain, free FAs activate different members of the FFAR family. In mammals, medium- and long-chain free FAs (6 – 12 and > 12 carbons, respectively) activate FFAR1 (GPR40) and/or FFAR4 (GPR120), while short-chain free FAs (< 6 carbons) activate FFAR2 (GPR43) and/or FFAR3 (GPR41) ⁽¹¹⁾. Activation of these receptors in TBC induces an intracellular signaling cascade (e.g. involving phospholipase C, calcium release-activated calcium modulator (Orai), stromal interaction molecule (Stim), sarco/endoplasmic reticulum Ca^{2+} (Serca)). This increases the amount of intracellular calcium, which depolarizes the membrane, facilitating the

release of neurotransmitters such as serotonin, which is considered the key neurotransmitter in TBC⁽¹²⁾. This signal transduction is then transmitted to the nucleus of the solitary tract in the myelencephalon (medulla oblongata) via gustatory nerves (VII and IX)⁽¹³⁾. This gustatory signal is propagated to the central level and processed, which regulates feeding behaviour⁽¹⁴⁾.

For fish, little is known about fat detection or its involvement in regulating FI especially concerning fat sensing⁽¹⁵⁾. Nevertheless, a distinct level of dietary ω -3 LC-PUFAs⁽¹⁶⁾ and alternative plant ingredients⁽¹⁷⁾ have been shown to influence RT feeding behavior, which suggests that palatability influences the perception of fat taste. Although oro-gustatory structures in RT are similar to those in mammals (i.e. the presence of TBC⁽¹⁸⁾ and some sensing receptors⁽¹⁷⁾), few studies have focused on the molecular mechanisms underlying lingual fat detection and integrated feeding responses in this species.

In farmed fish, understanding the mechanisms that regulate FI is essential to ensure the development of sustainable aquaculture in a context of changing availability of feed ingredients. This could help to develop specific diets that trigger a positive appetite response (via craving, hedonic sensation, or pleasure) and maximize growth performance. Here, we hypothesize that the alteration in growth performance observed in RT fed a V diet from their first meal may be linked to a disruption of their feeding behavior, which could be itself due to a differential regulation of the nutrient detection system, particularly with respect to ω -3 LC-PUFAs.

The present study evaluated the long-term influence (for eight months from the first feeding) of a vegetable diet devoid of EPA and DHA on the mechanisms that regulate fat detection in RT. To this end, we studied the expression of *ffars* and their signaling pathways in tongue tissue and assessed the integrated response in the central nervous system. We focused on the signaling pathways involving calcium and serotonin.

MATERIALS AND METHODS

Ethics statement

The trial was conducted in accordance with the Guidelines of the National Legislation on Animal Care of the French Ministry of Research (decree no. 2013-118, 1 February 2013) and following European Union legal frameworks related to the protection of animals used for scientific purposes (i.e. Directive 2010/63/EU). The scientists in charge of the trial received training and personal authorization. The trial was conducted at INRAE's NuMeA facilities (<https://doi.org/10.15454/GPYD-AM38>) and approved by the ethical committee (C2EA-73) of

INRAE “Comité d’éthique Aquitaine poissons oiseaux” (INRAE agreement no. 21699, 19 December 2019). All efforts were made to minimize the number of fish used and their suffering.

Fish rearing and experimental design

The fry used in this study came from diploid RT females of the same parental stock reared at INRAE’s experimental facilities at Lees-Athas (permit no. A64.104.1, Aspe valley, France). Swimming fry with an initial body weight of 104 ± 0.1 mg were transferred to INRAE’s experimental facilities at Donzacq (permit no. A40-228.1, Landes, France) for the feeding trial. Fingerlings were randomly allocated to 5 tanks per dietary treatment, with 150 fish per tank. Tanks ranging from 50-200 L were used, depending on fish size (maximum stocking density: 22 kg m^{-3} ; to avoid any selection, reduction in the number of fish was done randomly), to minimize the influence of tank size on feeding behavior⁽¹⁹⁾. The feeding trial was conducted for eight months (238 days) from the first meal at a constant water temperature ($17 \pm 1^\circ\text{C}$), with a natural photoperiod and a water flow rate adjusted to ensure a dissolved oxygen concentration greater than 90% of saturation. Fish were fed until apparent satiety. Fish were fed 7 times a day during the first days then the feeding frequency was gradually decreased to reach 2 meals per day from 10 grams until the end of the trial. Throughout the trial, dead fish were removed daily and weighed. Every three weeks (21 days, for a total of 11 periods), fish biomass and the quantity of feed distributed per tank were measured to calculate mean body weight, estimated daily FI (in units per day), and estimated feed efficiency (FE), as calculated below. For FI, due to the difference in body weight between the two diets throughout the trial (Figure 1), these values were expressed per kg metabolic body weight (MBW) per day. MBW was calculated as the geometric mean body weight, as previously described^(20–22).

$$\text{Growth} = \frac{Wf}{NBf}$$

$$\text{Feed intake} = \frac{\text{Feed distributed}}{\left(\frac{\text{Biomass} + Wi}{2}\right)^{0.75} / 1000 * \text{day}}$$

$$\text{Feed efficiency} = \frac{(\text{Biomass} + Wd) - Wi}{\text{Feed distributed}}$$

Wf: Final weight

NBf: Final number of fish

Wi: Initial weight

Wd: Death weight

For sampling, fish were first anaesthetized in a benzocaine bath at 30 mg L⁻¹ and then sacrificed in a benzocaine bath at 60 mg L⁻¹. After eight months of feeding, fish were sampled 20 min and 4 h after the last meal. To characterize the expression and regulation of the gustatory-sensing pathways, tissues were collected after frozen in liquid nitrogen and stored at -80°C. A total of 78 fish were sampled with seven or eight fish per tank and per diet (39 for C, and 39 for V). Twenty-eight fish were collected for qPCR analysis (tongue, hypothalamus and myelencephalon), with seven fish per diet (C or V) and per timing (20 min or 4 h after meal). Fourteen fish were collected 4 h after meal for WB analysis (tongue, and myelencephalon), with seven fish per diet (C or V). Thirty-six fish were collected for HPLC (tongue and myelencephalon) and MAO analysis (myelencephalon), either nine per diet (C or V), 4 h after meal and for each analysis (HPLC and MAO).

Experimental diets

Diets were manufactured at INRAE's experimental facilities at Donzacq using a twin-screw extruder (BC 45, Cleextral, Firminy, France). Pellet size was adapted to that of RT throughout the trial: 1 mm from 0–2 months, 2–3 mm from 2–4 months, and 4–5 mm from 4–8 months. Two control diets (C diet) and two vegetable diets (V diet) composed of 50% or 40% protein were formulated using Allix® software (A-Systems, Versailles, France). The diets (Table 1) were formulated to meet the nutritional requirements of RT as they grew (54.08% ± 0.45% protein in the two diet up to 20 g of BW, then 42.03% ± 0.83% until the end of the trial) ⁽²³⁾. To decrease the protein level, the proportion of whole wheat was increased, while the proportions of wheat and corn gluten were decreased.

The change in the diet profile occurred between period 4 and 5 (between day 84 and day 105) and between period 5 and 6 (between day 105 and day 126) for the C and V diets, respectively. This change in the proximate composition of the diet did not appear to have impacted the zootechnical parameters, as published elsewhere⁽³⁾. Furthermore, we accounted for differences in animal weight (geometric mean body weight) during measurements of the zootechnical parameters to avoid any potential biases in our analyses as previously published ^(20–22).

The C and V diets with identical protein content were also isoenergetic (24.79 kJ g⁻¹ ± 0.41% dry matter) and isolipidic (22.14% ± 0.38% dry matter) (Table 1). The ω-3 PUFAs were present in similar percentages in the C and V diets (ca. 30% FAs in the diet), but consisted mainly of DHA in the C diet and α-linolenic acid (ALA) in the V diet (Table 2). This percentage of ω-3 PUFAs was chosen to correspond to that found in the marine diet, as previously determined ⁽³⁾. The C diet contained FM (27%), FO (12.1%), and plant ingredients; the V diet was devoid of FM and FO, which were replaced with a blend of plant ingredients. The blend of vegetable oil in the V diet was chosen to

equal the percentages of the main FA classes in the C and V diets. To ensure the same amount of ω -3 PUFAs in the diets, DHA was replaced with ALA from linseed oil (10.8%) in the V diet. To obtain 26% DHA in the C diet, marine-based oil containing 70% DHA (OmegaVie® DHA 70 TG; Polaris, Quimper, France) was added (7.3%) in addition to fish oil (4.8%).

Synthetic L-lysine, L-methionine, dicalcium phosphate, and soy lecithin were added to all diets to correct deficiencies in essential amino acids, phosphorus, and phospholipids⁽²³⁾. A vitamin and mineral premix was also added to each diet. The profile of amino acid of the diet is already published⁽²⁴⁾.

Determination of proximate and FA composition

The nutrient composition of the diets was analyzed after 24 h of drying at 105°C. Ash content was determined by combustion in a muffle furnace for 16 h at 550°C. Gross energy was measured using an adiabatic bomb calorimeter (IKA, Heitersheim Gießheimer, Germany). Crude protein was determined using the Kjeldahl method⁽²⁵⁾. Crude lipids were determined using the Soxhlet method. Total lipid content was extracted, purified, and measured gravimetrically according to the Folch method⁽²⁶⁾ using dichloromethane instead of chloroform. All these analyses were performed following previously described protocols⁽²⁴⁾.

Total RNA isolation and real-time quantitative PCR

Total RNAs were isolated from the tongue and myelencephalon. Tissues were ground in TRIzol® reagent (Invitrogen, Carlsbad, CA, USA) using a tissue homogenizer (Precellys® 24, Bertin Technologies, Montigny-le-Bretonneux, France) and 1.4 mm ceramic beads for 2 × 20 s at 3,075× g, separated by a 15 s break, for the myelencephalon, or 2.8 mm ceramic beads for 2 × 30 s at 3,720× g, separated by a 30 s break, for the tongue. The quantities of extracted RNAs were measured using a spectrophotometer (ND-1000, Nanodrop, Thermo, Vantaa, Finland), and the quality of extracted RNAs was estimated using the 260 nm:280 nm OD ratio (purity of DNA and RNA) and 260 nm:230 nm OD ratio (purity of solution, to avoid phenol contamination). For gene-expression analysis, only total RNAs with a 260 nm:280 nm OD ratio exceeding 1.8 and a 260 nm:230 nm OD ratio exceeding 2.0 were utilized.

Two micrograms of total RNAs were reverse-transcribed into cDNA using the Super-Script III RNase H-Reverse transcriptase kit (Invitrogen) and random primers (Promega, Chartonnières-les-Bains, France) (Table 3) according to the manufacturer's recommendations.

The LightCycler® 480 system (Roche Diagnostics, Neuilly-sur-Seine, France) was used to measure mRNA levels via real-time quantitative PCR (RT-qPCR). The reaction mixture was 6 µL and consisted of 2 µL of diluted cDNA (1:10), 0.24 µL of each primer (10 µmol L⁻¹), 3 µL of LightCycler® 480 SYBR Green I Master mix, and 0.52 µL of DNase/RNase-free water (5 Prime GmbH, Hamburg, Germany). Thermal cycling was initiated with incubation at 95°C for 10 min for the initial denaturation of the cDNA and hot-start Taq-polymerase activation, followed by 45 cycles of a two-step amplification program (15 s at 95°C; 10 s at 60°C). Cycle thresholds greater than 35 cycles were not considered. Melting curves were controlled systematically (temperature gradient: 1.1°C/15 s from 65°C to 97°C) at the end of the last amplification cycle to ensure that only one fragment was amplified. During a run, each sample was deposited twice, and two negative controls were added: a pool of samples without reverse transcriptase and a pool of samples without cDNA. The efficiency of all qPCR reactions was 96-100%. The sequences of amplicons had been previously assessed via sequencing⁽²⁷⁾. Data were extrapolated from standard curves and normalized to the housekeeping elongation factor 1α gene (*eef1a*) in the myelencephalon and keratin 8b gene (*krt8b*) in the tongue. Relative expression of the target genes was determined using the $\Delta\Delta$ CT method⁽²⁸⁾.

Indolamine release measurement using ultra-high-pressure liquid chromatography

All reagents and standards were purchased from Merck KGaA (Darmstadt, Germany). All organic solvents used were gradient HPLC grade (ADL & Prochilab, Lormont, France). Ultrapure water was made daily using a water purification system (Simplicity®, Merck Millipore, Bedford, MA, USA).

The myelencephalons and tongues were homogenized using a tissue homogenizer (Precellys® 24) in 20 mM phosphate and 1 mM EDTA buffer (pH = 6.5 ± 0.05). An initial centrifugation (14,000×g, 20 min, 4°C) with a 10% volume-to-volume metaphosphoric acid solution deproteinized the supernatant. After a second centrifugation (14,000×g, 5 min, 4 °C), the supernatant was filtered with a 0.22 µm polyvinylidene fluoride (PVDF) unit. Metabolite mixtures were stored at -20°C (for no longer than one week) until analysis.

Indolamines from standard mixtures and polar metabolite extracts were separated and detected using ultra-high-pressure liquid chromatography (UHPLC) (ACQUITY H-Class PLUS System, Waters, Milford, MA, USA) equipped with a thermostatted autosampler supported by a fluorescence detector (ACQUITY Multi-λ, Waters). Data were obtained and quantified using Empower™ Pro software (Waters). Chromatographic separation was performed on a UHPLC column (150 × 4.6 mm, i.d. 3

µm) at 30°C (PFP(2), Phenomenex, Torrance, CA, USA). The injection volume was 10 µL, and the flow rate was set at 0.4 mL min⁻¹.

A quaternary solvent system was used, which consisted of (A) a 10 mM phosphate buffer (pH 4.3 ± 0.05), (B) methanol, (C) ultrapure water, and (D) acetonitrile. The mobile phase was filtered through in-line 0.2 µm membrane filters. The following linear gradient elution was used: 0–11 min: 80% A, 20% B; 16 min: 50% C, 50% D; 16–26 min: 50% C, 50% D; 27 min: 80% A, 20% B; and 27–35 min (column equilibration): 80% A, 20% B. The eluate was monitored using dual excitation/emission fluorescence detection: 285 nm and 355 nm for serotonin (5-hydroxytryptamine; 5-HT) and 5-hydroxyindolacetic acid (5-HIAA) respectively. Metabolites of interest were identified by comparing retention times to standards. Quantification was based on integrating peak areas and comparing them to standard calibration curves ($R^2 > 0.999$) of each metabolite of interest. Calibration curves were linear from 0.01–3 pmol per injection for serotonin and 5-HIAA.

Protein extraction and western blotting

Whole RT tongues were ground with a mortar and pestle under liquid nitrogen, and 200 mg of powder was collected. The tongue and myelencephalon were then homogenized in 1 ml T-Per extraction reagent (#78510, Thermo Fisher Scientific, Waltham, MA, USA) using a tissue homogenizer (Precellys® 24) at 4°C. The homogenates were subjected to two successive centrifugations (1,000× g, 15 min, 4°C and 20,000× g, 30 min, 4°C), and then supernatants containing cytoplasmic proteins were collected. Protein concentrations were determined using the bicinchoninic acid assay method (BCA, #BCA1-1KT, Sigma-Aldrich, Saint Louis, MO, USA). Proteins in the tongue and myelencephalon (20 µg of protein for each) were separated using SDS-PAGE gel electrophoresis with different percentages of acrylamide, depending on the size of the protein (6%, 7.5%, 12% and 15%) and transferred to a 0.45 µm PVDF membrane (#IPFL00010, Merk Millipore) via electrophoretic transfer. Membranes were blocked at room temperature for 1 h with an Intercept® phosphate buffered saline (PBS) blocking buffer and then incubated overnight at 4°C with the appropriate primary antibody: ORAI1 rabbit polyclonal IgG (PA5-26378, Invitrogen) (1:2000), STIM1 rabbit monoclonal IgG (D88E10#5668, Cell Signaling, Danvers, MA, USA) (1:1000), TPH2 rabbit polyclonal IgG (PA5-37803, Invitrogen) (1:1000), or VMAT2 mouse monoclonal IgG (Sc-374079, Santa Cruz Biotechnology, Dallas, TX, USA) (1:1000). The specificity of the mammalian antibodies used had been previously assessed via alignment of amino acid sequences of peptides using the SIGENAE database (<http://www.siginae.org>) to check for conservation of the antigen sequence with the corresponding sequence from mammals, which ensured good specificity (> 88%) of the mammalian antibodies used

to analyze the samples. The molecular weight of the band agreed with that of the desired protein. Membranes were washed three times with PBS containing 0.1% TWEEN® 20 (Sigma-Aldrich) and then incubated with the corresponding secondary antibodies for 1 h at room temperature. Membranes were washed three times with PBS containing 0.1% TWEEN® 20 and twice with PBS. Immunoreactivity was detected using enhanced chemiluminescence after incubating the membrane in Super Signal™ West Pico PLUS Chemiluminescent Substrate (#34580, Thermo Fisher Scientific) at room temperature. Chemiluminescence was acquired using an imager (iBright FL1500, Thermo Fisher Scientific). Two types of normalization were required due to the weight of the protein and the protein-loading control. ORAI1 and VMAT2 were normalized using tubulin, while STIM1 and TPH2 were normalized with total protein staining, using No-Stain™ Protein Labelling Reagent (#44717, Thermo Fisher Scientific).

Enzymatic activity

MAO-Glo™ Assay (#V141A, Promega) was used to measure enzymatic activity of monoamine oxidase (MAO) A and B, an enzyme responsible for degradation of serotonin to 5-HIAA. Whole RT tongues were ground with a mortar and pestle under liquid nitrogen, and 200 mg of powder was collected and placed in a centrifuge (Eppendorf, Hamburg, Germany) with an MAO reaction buffer. The whole myelencephalon was placed in the same centrifuge without prior grinding. Fractions were separated using centrifugation (5,000×g, 10 min, 4°C). Samples and analysis plates were prepared according to manufacturers' recommendations. Tongue tissue was diluted 2 times and myelencephalon 40 times. Mixtures of samples and the MAO solution without luciferin were used as negative controls. To calculate the MAO A signal, we subtracted the mean luminescence of the negative control and then subtracted that of the MAO B reaction.

Statistical analysis

All statistical analyses were performed using R software (version 3.6.1) (R Core Team (2019). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria) with the R Commander graphical user interface. The tank was the experimental unit for data on growth parameters and one-way ANOVA (repeated measures) was used for statistical analysis of growth parameters. In comparison, individual fish were the experimental unit for data on gene expression, indolamine release, protein abundance, and MAO activity. The results were calculated as means ± standard error of the mean (SEM). Gene expression data were statistically analyzed using two-way ANOVA considering both time and diet as factors.

When the interaction between time and diet was found significant, a one-way ANOVA was performed to assess the individual effects of diet or time. When appropriate, analysis was followed by Tukey's post-hoc test. When the normality and homogeneity of variances were confirmed (Shapiro-Wilk and Levene's test, respectively), *t*-test (student) were performed on the different variables obtained by WB, MAO, HPLC and qPCR (Myelencephalon ; Figure 4 and hypothalamus, Figure 5) analysis. Kruskal–Wallis non-parametric tests were performed when the conditions of application were not respected (for *npya*, *npyb*, *pomcb*, *cartpt*, MAO B). Differences were considered significant at $p < 0.05$.

RESULTS

Zootechnical parameters

Throughout the trial, fish fed the V diet grew statistically less than those fed the C diet for all periods measured (Figure 1A, $p < 0.05$). At the end of the trial, the difference in mean body weight between fish fed the C vs. V diet was nearly 34% ($p < 0.05$): 337.0 and 228.9 g, respectively. FI was significantly lower for fish fed the V diet for the first three periods (first feeding to 62 days) and from days 125-146 (Figure 1B). FI was significantly higher for fish fed the V diet from days 104-125 and 188-209. FE was significantly lower for fish fed the V diet for four 3-week periods (days 20-41, 62-104, and 188-209) (Figure 1C). The survival (data not shown) did not differ between RT fed the C vs. V diet ($87.0\% \pm 3.47\%$ vs. $87.1\% \pm 2.02\%$, respectively).

Long-term influence of the V diet on mRNA level of *ffar* in RT

Expression of *ffar1* and *ffar2* measured in the tongue of RT 20 min and 4 h after the last meal varied (Figure 2). The mRNA level of *ffar2b1.1*, *ffar2b1.2*, and *ffar2b2b2* had the same expression pattern. The highest expression was measured 20 min after feeding for fish fed the V diet. The expression decreased significantly from 20 min to 4 h, and reached that measured in fish fed the C diet, for which no significant difference was observed between 20 min and 4 h. The mRNA level also decreased from 20 min to 4 h for *ffar2a1a* in fish fed the V diet, but not for those fed the C diet. The mRNA level of *ffar1* was downregulated between 20 min and 4 h for both diets, but mRNA level at 20 min was significantly higher in fish fed the V diet than those fed the C diet. The mRNA level of the *ffar2a2*, *2b2a*, and *2b2b1* isoforms did not differ as a function of time or diet.

Calcium-signaling pathway gene expression and protein abundance in tongue tissue

The mRNA levels of calcium-signaling pathway markers in the proximal tongue of fish varied (Figure 3). Isoforms of phospholipase C showed different kinds of regulation. The mRNA

level of *plcb2* (phospholipase C β 2) showed a time effect, with higher expression at 4 h for both diets (Figure 3A). The mRNA level of *plcb3* increased from 20 min to 4 h after the last meal, but only in fish fed the C diet (Figure 3B). The mRNA level of *plcb4* showed higher expression in fish fed the C diet (Figure 3C). The expression of *stim1a* and *stim1b* was influenced by a postprandial effect, with increased expression from 20 min to 4 h, regardless of the diet (Figure 3D and E).

For fish fed the C diet, the mRNA level of *orai1a* and *orai1b* increased from 20 min to 4 h (Figure 3G and H). In contrast, *orai1a* expression decreased sharply from 20 min to 4 h after feeding, while *orai1b* expression increased the highest level of the four experimental conditions. The mRNA level of *serca3a* was upregulated at 4 h, but only for fish fed the C diet (Figure 3J). Protein abundance of STIM1 and ORAI1 was also investigated 4 h after the meal, but neither differed significantly (Figure 3F and I, respectively).

Serotonin-signaling pathway in tongue tissue and the myelencephalon

For indolamine turnover in the tongue 20 min after feeding, fish fed the V diet had less serotonin than fish fed the C diet, but more 5-HIAA ; thus, fish fed the V diet had a higher 5HIAA:serotonin ratio than fish fed the C diet (Figure 4A). The mRNA level of genes encoding components of the serotonin-signaling pathway in the myelencephalon varied little: of the genes analyzed, only the mRNA level of the serotonin receptor (*5ht1aa*) increased for fish fed the V diet (Figure 5B). The protein abundance of TPH2 (tryptophan hydroxylase, isozyme of serotonin synthesis from tryptophan) was similar for both diets (Figure 4C), while that for VMAT2 (vesicular monoamine transporter 2 of serotonin) was significantly higher in fish fed the C diet (Figure 4D). The enzymatic activity of MAO A in the myelencephalon was significantly higher in fish fed the V diet (Figure 4 E).

Hypothalamus neuropeptide gene expression profile

In the hypothalamus 6 h after the last meal, diets had no influence on the mRNA level of orexigen *npv* (neuropeptide y) (a and b paralogs) or *agrp1* and 2 (agouti-related neuropeptide) or on mRNA level of anorexigenic *pomc* (proopiomelanocortin) (a and b paralogs) or *cartpt* (cocaine and amphetamine regulated transcript protein) peptides (Figure 5). Likewise, diets had no influence on mRNA level in the hypothalamus of *mc4ra* (melanocortin 4 receptor), a gene-encoding receptor upregulated by anorexigenic POMC peptide and downregulated by AGRP orexigenic peptide.

DISCUSSION

In mammals, taste perception plays a key role in food acceptability and provides critical information about the quality and nature of nutrients. Nutrients interact with specific receptors and induce specific feeding behaviors (consumption or avoidance) that determine growth and survival⁽²⁹⁾. However, in fish, especially farmed fish, the mechanisms of feed detection that can ultimately influence the regulation of FI are largely unknown^(15,30). The present study is the first to identify the mechanism of feed sensing, specifically FAs sensing, in the tongue and brain of RT. Additionally, it highlights the regulation of this mechanism through a comparison between a V diet (with ALA as the primary source of ω -3 LC-PUFAs) and a diet containing FM and FO with a high level of DHA.

RT fed the V diet had significantly lower growth rates than those fed the C diet containing marine ingredients (i.e. mean body weight nearly 34% lower after 238 days). This decreased growth due to diets without FM and FO is well known in the literature, with several studies concluding that RT fed a plant-based diet grew less than those fed a diet containing FM and FO^(3,31,32). This decreased growth could be due to changes in feeding or reduced FE. We assumed that this decreased growth was related to changes in FI, which was lower for RT fed the V diet from first feeding to day 63. This decrease in growth performance due to the V diet could be related to the absence of marine products, such as FM and particularly FO^(33,34). Plant ingredients have other disadvantages, particularly the imbalance of certain nutrients such as ω -3 LC-PUFAs (DHA and EPA), which change feeding behavior (motivation to consume feed) and decrease growth performance greatly. In the present study, both diets were isolipidic but the C diet was rich in DHA (and to a lesser extent EPA), while the V diet had no EPA or DHA, but contained ALA to maintain the ω -3 LC-PUFAs content. Recently, RT were observed to prefer diets rich in EPA and DHA⁽³⁵⁾. Thus, we assume that the level of DHA and EPA in the present study's diet may have influenced the regulation of FI in trout. Nonetheless, although DHA and EPA were the only FAs in the C diet that were absent in the V diet, the formulation of the two diets differed; thus, the influence on FI may have been related to compounds other than ω -3.

Fish have been studied for their response to various taste modalities, including sourness, bitterness, saltiness, sweetness (sucrose), as well as substances such as acetic or hydrochloric acid, quinine, and sodium chloride. However, the compounds that have received the most attention are free amino acid (reviewed by Kasumyan and Døving, 2003)⁽³⁶⁾. Free amino acids elicit strong behavioral responses in all fish species, predominantly stimulatory, although some can act as deterrents and, very often, indifferent (no effect on fish behaviour). Although only a few amino acids

are known to stimulate the gustatory system in RT, a recent study demonstrated that a single oral administration of proline and glutamic acid decreased feed intake ⁽³⁷⁾. The effect of proline is consistent with classical electrophysiological studies in rainbow trout, where proline was found to be the most effective amino acid in stimulating both the extraoral surface, the rostral (palate) and posterior parts of the oral cavity ⁽³⁸⁾. Furthermore, existing literature indicates that carbohydrates are widely recognized as palatable to various herbivorous fish species such as grass carp (*Ctenopharyngodon idella*). Additionally, carbohydrates have been found to be palatable to omnivorous fish, including dace (*Leuciscus leuciscus*), roach (*Rutilus rutilus*), guppy (*Poecilia reticulata*), black molly (*Poecilia sphenops*), and platy (*Xiphophorus maculatus*), whose natural diet in the wild predominantly consists of algae. However, it is noteworthy that carnivorous fish tend to show indifference towards carbohydrates ⁽³⁶⁾. In RT, one study found that sugars elicited a positive behavioral response (cotton pellets flavoured or not ; swallowed vs rejected) ⁽³⁹⁾. In addition, fish are known to be attracted to various substances other than free amino acids and sugars. These include quaternary amines such as betaine (glycine betaine, tri-methylglycine), nucleotides, carboxylic acids (e.g., citric acid, lactic acid, butyric acid, etc.), bile salts, steroid hormones, and prostaglandins. For RT, Jones et al., have demonstrated a positive behavioral response to certain amides (acetamide, valeramide, and caproic amide), alcohols (hexanol and octanol), and four types of sugars ⁽³⁹⁾. These findings suggest that RT are sensitive to a wide range of compounds not only FAs, which may be important to study in the context of alternative plants products.

After 62 days, it is interesting to note that the decrease in FI was not observed. The initial changes observed in FI from day 1 to day 62 did not persist over the long term between 62 to 238 days. While 5 weeks periods were not different in FI between the two diets (including the last period of the trial corresponding to the analyses presented in this study), 2 weeks periods showed better FI on V diet and one for C diet. We postulate that the fish underwent adaptations to accommodate the provided diet, or alternatively, the juvenile fish were compelled to consume it in order to ensure their survival. In a recent study that included a similar formulation, we demonstrated that from the juvenile stage, RT had no preference for a commercial (with DHA and EPA) or plant-based (with ALA but without DHA and EPA) diet using a self-feeder method ⁽¹⁷⁾. Thus, we were interested in the postprandial regulation of FI in RT fed the V diet over the long term. FI are regulated in the central nervous system, where the neural circuitry integrates incoming sensory information via the lingual system to orchestrate an integrated feeding response. Thus, taste receptors in mammals have many functions ⁽⁷⁾. They guide appetite and trigger physiological processes for absorbing nutrients and facilitate the regulation of satiation ⁽⁴⁰⁾. In the present study, we observed differential expression of

ffar1 and paralogs of *ffar2* genes and their signaling pathways depending on the diet. First, *ffar1* and four out of seven paralogs of *ffar2* were upregulated 20 min after the meal in RT fed the V diet. This is in agreement with studies on rodents ⁽⁴¹⁾ that observed higher expression of the *ffar4* gene (GPR120) immediately after a meal, which could indicate the influence of postprandial mechanisms in fat-eating behavior. The literature indicates that activation of *ffar1* and *ffar4* could result in an anorectic effect that decreases FI. Indeed, Gorski et al. ⁽⁴²⁾ showed that early (20 min after meal) activation of FFAR1 (by an agonist) in mice decreases FI (by more than 38%) and body weight. FI is mainly centrally regulated, which raises questions about whether the influence of FFAR on body weight is due to direct or indirect effects on the central nervous system ⁽⁴²⁾. Many studies show that *ffars* are upregulated immediately after a meal and that FFAR1, 2, and 4, could play a critical role in several physiological homeostasis mechanisms, especially the regulation of appetite, eating disorders, or food preferences ^(11,43–45). Our results indicate this also occurs in RT. Ozdener et al. ⁽⁴⁶⁾ demonstrated that in humans and mice, incubation of TBC with ALA resulted in the upregulation of *ffar4*. These results agree with our results that indicate higher expression of the *ffars* gene 20 min after the meal for fish fed the V diet vs. the C diet. However, these studies are unable to relate the expression pattern of *ffars* to the signaling pathways (no difference in the expression pattern in response to calcium-signaling markers and a decrease in serotonin with the V diet) or the integrated response that we observed. This could be due to other taste receptors associated with the same signaling pathways that could mask the influence of *ffar* observed in the present study. However, our results suggest that FFAR1 and some paralogs of FFAR2 could be involved in the FA-sensing system. In mammals, this role is performed by FFAR1 ⁽⁴²⁾ and FFAR4 ⁽⁴⁷⁾. We speculate that FFAR1 and paralogs of FFAR2 in RT could have acquired the function of FFAR3 and FFAR4 reported in similar studies that used other models. However, this must be confirmed in future studies especially in short-term experiment from the first meal where FI appears to be particularly affected for fish fed a V diet.

The binding of free FAs on FFARs could activate the receptors, resulting in an increase in intracellular calcium concentrations. One of the first actors in this calcium-signaling pathway is *plcb*, which plays a role in GPCR-mediated signal transduction in RT ⁽⁴⁸⁾ and in the feeding response. The regulation of the three *plcb* isoforms followed the same trend: they were upregulated 4 h after the meal compared to 20 min in RT fed the C diet. Velasco et al. showed that *plcb1* expression was upregulated after 2 h of treatment with a FFAR4 agonist, but not the expression of *plcb 2, 3, or 4* in the hypothalamus of RT ⁽⁴⁹⁾. In contrast, our observation of the tongue suggested that *plcb* could be expressed in particular isoforms depending on the tissue considered. Store-operated calcium channels

(SOC), responsible for calcium entering the cell, are composed of Orai1 and Orai1/3 proteins⁽⁵⁰⁾. Two *orai1* isoforms were upregulated after 4 h with the C diet. The postprandial response of the *orai1a* and *orai1b* genes differed with the V diet, with *orai1a* expression downregulated 4 h after the meal, while *orai1b* expression was upregulated. Studies of rodents showed that *orai1* is expressed in TBC and functions in SOC channels to assure Ca²⁺ influx after depletion of the endoplasmic reticulum pool induced by LC-PUFAs⁽⁵¹⁾. Thus, our study confirms that the FA profile in the diet can also influence the regulation of *orai1* in RT. STIM is another actor involved in opening the SOC channels. In the present study, the two isoforms (*stim1a* and *stim1b*) had higher expression after 4 h for both diets. Shanmugamprema et al.⁽⁵¹⁾ suggest that *stim1* plays a key role in regulating fat taste in mice⁽⁵¹⁾. These actors of the calcium-signaling pathways are regulated by diet and the FA profile. In RT, they are involved in calcium-signaling pathways in intestinal tissues and are regulated by diet⁽⁵²⁾.

The processing and cerebral integration of gustatory sensory information is complex. The same cell-signaling cascades and molecular actors are involved in activating these receptors. However, the membrane depolarization that triggers the electrical signal, which is transmitted to the cerebral areas connected to different nerve bundles and cellular structures, differs. Thus, the brain areas involved must be able to process complex signals that are quantitative (i.e. signal intensity depending on the number of molecules bound and receptors activated), qualitative (i.e. discriminating receptors depending on the binder), and hedonic (e.g. good, bad, pleasant) in correlation with internal signals of the body (e.g. hunger, thirst). Thus, in animals (including fish), the large number of sensory receptors and paralogs for the same receptor (e.g. *ffar2*), combined with the strong discriminating ability of the animal (i.e. several thousand taste molecules), could reflect the presence of highly sophisticated and divergent mechanisms of amplification and coding of sensory information. After oral detection of a nutrient, the signals are transmitted to the brain and induce an integrated feeding response⁽⁷⁾. However, we observed no differences in the expression pattern of markers in FI in central areas of feeding behavior regulation. This is puzzling, because our results showed that reduced FI had a major influence on RT fed the V diet especially at early stage. It is possible that sampling was not performed during the most suitable postprandial period, even though 6 h postprandial is adequate to study the regulation of these genes by FFARs in the hypothalamus of RT⁽³⁰⁾. In addition, FI did not differ between the two dietary groups during the last period of the trial, which could also explain the lack of difference in FI markers in the brain.

The brain's reward circuitry involved in regulating FI is activated by taste and the consumption of palatable food⁽⁵³⁾, two essential criteria met by dietary lipids such as ω -3 LC-PUFAs. Studies have shown that sensory information is related to taste sensitivity and palatability⁽⁵⁴⁾. Dietary

lipids influence feeding behavior due to their textural properties in the mouth (e.g. gluiness, smooth, succulent, tender, and velvety). Thus, beyond a purely neuropeptide regulation, the influence of FI could be related more to hedonic sensations of the diet. A previous study based on the same trial⁽¹⁷⁾ indicated that changes in the FI of the V diet are related to a decrease in desire to eat rather than changes in the mechanisms of FI (i.e. regulation of appetite neuropeptide). RT can discriminate different levels of ω -3 LC-PUFAs in the diet and prefer and consume more of diets richer in ω -3 LC-PUFAs⁽³⁵⁾. Based on our results, ω -3 LC-PUFAs appear to provide textural properties to feed and render it more palatable, which ALA does not. In mammals, the sensation of fat plays an important role in the desire to eat due to sensory perception and/or pleasure, independent of regulation by the homeostatic regulatory circuit in the hypothalamus (i.e. regulation of appetite neuropeptide). This is called “hedonic eating behavior”⁽⁵⁵⁾. This hedonic regulation of FI is influenced by sensory and reward signals, which can dominate the homeostatic control and result in overconsuming foods with high palatability, even when energy requirements have been satisfied⁽⁵¹⁾. The reward circuit, such as the serotonin system, drives the hedonic motivation for FI. Serotonin signaling occurs in several regions of the brain involved in the homeostatic and hedonic circuits of FI regulation^(56,57). Serotonin also influences cognitive and behavioral functions, especially depressive disorders, motivation, and stress or anxiety-like behavior when low levels are detected⁽⁵⁸⁾. We observed a higher level of serotonin in its synthesis areas (i.e. raphe nucleus in the myelencephalon)⁽⁵⁶⁾ in RT fed the C diet. The higher level of serotonin associated with the C diet could be due to reuptake and storage in the vesicles⁽⁵⁸⁾, and not necessarily to an increase in synthesis. This mechanism and the brain turnover of serotonin regulation is the main prototypical neuromodulator involved in the adaptation (for the fine-tuning of most behaviors) of all animals⁽⁵⁸⁾. This is supported by the higher level of VMAT2, which controls entry of serotonin into the vesicles, and the lower degradation of serotonin related to lower MAO activity (an enzyme that degrades serotonin) in fish fed the C diet. However, the increase in serotonin in tongue tissue in RT fed the C diet cannot explain the increase in their hedonic eating behavior. In contrast, in previous study of the same RT, we observed a higher serotonin level in the telencephalon, which is the major center for decision-making that receives serotonin neurons inputs in fish, during a fasting period⁽²⁴⁾. The telencephalon can be responsible for this hedonic sensation in RT⁽⁵⁹⁾. In the same way, in this area, the precursor of dopamine (L-DOPA), a neurotransmitter that is a marker of the stress response, motivation, and ill-being when high in a chronic situation or due to long-term effects of feeding⁽²⁴⁾, was higher in RT fed the V diet. We also demonstrated that RT fed a diet enriched in ω -3 LC-PUFAs increased the markers of brain activity that may be associated with higher acceptance of these specific diets (e.g. decrease markers of inflammatory and oxidant status, and cortisol pathways; increase genes regulating normal animal behavior)⁽⁶⁰⁾. The increase in hedonic

sensation in RT fed the C diet in the long term may have increased the desire to eat (potentially induced by ω -3 LC-PUFAs in the diet). The response of fish fed a plant-based diet in the long term could be interpreted that they do not like the feed and that it does not provide pleasure or stimulate the appetite. Changes in FI would be visible in the initial stage after the first meal and would decrease over time due to the fish's need to eat to survive. The deregulation of serotonin turnover instead of regulation by the homeostatic regulatory circuitry in the hypothalamus (regulation of appetite neuropeptide) confirmed this long-term lack of desire to eat. Thus, a dynamic short-term experiment from the first meal could identify this defect in the brain and establishment of this defect in serotonin. It would also be very interesting to understand the relation between the decrease in the fat-detection sensing system in the tongue and the decrease in serotonin that causes a defect in hedonic sensation.

Overall, this study was the first to identify that changes in growth performance and FI of RT fed a vegetable diet could be related to an initial change in food detection and a decrease in long-term hedonic sensation (i.e. motivation to eat). We revealed that RT had the same fat-detection mechanisms as mammals (i.e. presence of nutrient receptors in the tongue, early activation of calcium-signaling pathways, and serotonin release followed by integrated feeding responses). The vegetable diet induces specific regulations at the lingual level, affecting fat receptors, calcium signaling, and early serotonin turnover, while neuropeptides remain unaffected. This modulation of fat sensing may play a role in detecting and integrating brain responses after meals over the long term. These findings have implications for the aquaculture sector, as they can assist in developing specialized diets that stimulate a positive appetite response (such as craving, hedonic sensation, or pleasure) and maximize the growth performance of farmed fish from their very first meal, using alternative plant-based ingredients.

ABBREVIATIONS

5-HIAA	5-hydroxyindolacetic acid
ω -3 LC-PUFA	ω -3 long chain polyunsaturated fatty acid
Agrp	agouti related neuropeptide
ALA	α -linolenic acid
BCA	bicinchoninic acid assay
C diet	control diet
DHA	docosahexaenoic acid
EPA	eicosapentaenoic acid

FAs	fatty acids
FFAR	free fatty acid receptor
FE	feed efficiency
FI	feed intake
FM	fish meal
FO	fish oil
GPCRs	G protein-coupled receptors
HVA	homovanillic acid
MAO	monoamine oxidase
MBW	metabolic body weight
NTS	nucleus of the solitary tract
Plcb	phospholipase C β
PVDF	polyvinylidene fluoride
RT-PCR	real-time quantitative PCR
Serca	sarco/endoplasmic reticulum Ca^{2+}
SOC	store-operated calcium
Stim	stromal interaction molecule
TBC	taste bud cells
Tph	tryptophan hydroxylase
Vmat	vesicular monoamine transporter
V diet	vegetable diet
RT	rainbow trout
UHPLC	ultra-high-pressure liquid chromatography

AUTHOR CONTRIBUTIONS

E.B.: methodology, software, data curation, writing-original draft preparation; **C.H.:** data curation, software, analysis of indolamine brain concentration; **L.L.:** data curation, analyzed lipid content; **A.S.:** data curation, analysis of diet composition; **A.L.** and **F.T.:** data curation, methodology; **S.SC.:** conceptualization, methodology, writing-review and editing. **J.R.:** conceptualization, methodology, software, data curation, writing-original draft preparation, funding acquisition. All authors have read and agreed to the published version of the manuscript.

BIBLIOGRAPHY

1. (2020) *The State of World Fisheries and Aquaculture 2020*. FAO.
2. Naylor RL, Hardy RW, Bureau DP, et al. (2009) Feeding aquaculture in an era of finite resources. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 15103–15110.
3. Lazzarotto V, Médale F, Larroquet L, et al. (2018) Long-term dietary replacement of fishmeal and fish oil in diets for rainbow trout (*Oncorhynchus mykiss*): Effects on growth, whole body fatty acids and intestinal and hepatic gene expression. *PLoS ONE* **13**, 25 [Martínez-Llorens S, editor].
4. Martínez-Ruiz NR, López-Díaz JA, Wall-Medrano A, et al. (2014) Oral fat perception is related with body mass index, preference and consumption of high-fat foods. *Physiol Behav* **129**, 36–42.
5. Cox DN, Hendrie GA & Carty D (2016) Sensitivity, hedonics and preferences for basic tastes and fat amongst adults and children of differing weight status: A comprehensive review. *Food Quality and Preference* **48**, 359–367.
6. Kindleysides S, Beck KL, Walsh DCI, et al. (2017) Fat Sensation: Fatty Acid Taste and Olfaction Sensitivity and the Link with Disinhibited Eating Behaviour. *Nutrients* **9**, 879.
7. Behrens M & Meyerhof W (2011) Gustatory and extragustatory functions of mammalian taste receptors. *Physiology & Behavior* **105**, 4–13.
8. Chaudhari N & Roper SD (2010) The cell biology of taste. *J Cell Biol* **190**, 285–296.
9. Jaime-Lara RB, Brooks BE, Vizioli C, et al. (2023) A systematic review of the biological mediators of fat taste and smell. *Physiological Reviews* **103**, 855–918.
10. Liu D, Archer N, Duesing K, et al. (2016) Mechanism of fat taste perception: Association with diet and obesity. *Progress in Lipid Research* **63**, 41–49.
11. Kimura I, Ichimura A, Ohue-Kitano R, et al. (2020) Free Fatty Acid Receptors in Health and Disease. *Physiological Reviews* **100**, 171–210.
12. Roper SD (2006) Signaling in the Chemosensory Systems: Cell communication in taste buds. *Cell. Mol. Life Sci.* **63**, 1494–1500.
13. Khan AS, Keast R & Khan NA (2020) Preference for dietary fat: From detection to disease. *Progress in Lipid Research* **78**.
14. Laugerette F, Passilly-Degrace P, Patris B, et al. (2005) CD36 involvement in orosensory detection of dietary lipids, spontaneous fat preference, and digestive secretions. *J Clin Invest* **115**, 3177–3184.

15. Morais S (2017) The Physiology of Taste in Fish: Potential Implications for Feeding Stimulation and Gut Chemical Sensing. *Reviews in Fisheries Science & Aquaculture* **25**, 133–149. Taylor & Francis.
16. Roy J, Mercier Y, Tonnet L, et al. (2020) Rainbow trout prefer diets rich in omega-3 long chain polyunsaturated fatty acids DHA and EPA. *Physiol. Behav.* **213**, 112692.
17. Roy J, Baranek E, Mercier Y, et al. (2022) Involvement of Taste Receptors in the Oro-Sensory Perception of Nutrients in Rainbow Trout (*Oncorhynchus Mykiss*) Fed Diets with Different Fatty Acid Profiles. *Aquaculture Nutrition* **2022**, 1–20 [Du Z-Y, editor].
18. Abbate F, Guerrera MC, Levanti M, et al. (2020) Anatomical, histological and immunohistochemical study of the tongue in the rainbow trout (*Oncorhynchus mykiss*). *Anat Histol Embryol* **49**, 848–858.
19. Roy J, Terrier F, Marchand M, et al. (2021) Effects of Low Stocking Densities on Zootechnical Parameters and Physiological Responses of Rainbow Trout (*Oncorhynchus mykiss*) Juveniles. *Biology* **10**, 1040.
20. Saravanan S, Schrama JW, Figueiredo-Silva AC, et al. (2012) Constraints on Energy Intake in Fish: The Link between Diet Composition, Energy Metabolism, and Energy Intake in Rainbow Trout. *PLoS ONE* **7**, e34743 [Blanc S, editor].
21. Saravanan S, Geurden I, Figueiredo-Silva AC, et al. (2012) Control of voluntary feed intake in fish: a role for dietary oxygen demand in Nile tilapia (*Oreochromis niloticus*) fed diets with different macronutrient profiles. *Br J Nutr* **108**, 1519–1529.
22. Figueiredo-Silva AC, Kaushik S, Terrier F, et al. (2012) Link between lipid metabolism and voluntary food intake in rainbow trout fed coconut oil rich in medium-chain TAG. *Br. J. Nutr.* **107**, 1714–1725.
23. National Research Council (2011) Nutrient requirements of fish and shrimp. *The National Academies Press, Washington, DC*.
24. Heraud C, Hirschinger T, Baranek E, et al. (2022) Detection and Modulation of Olfactory Sensing Receptors in Carnivorous Rainbow Trout (*Oncorhynchus mykiss*) Fed from First Feeding with Plant-Based Diet. *IJMS* **23**, 2123.
25. Kjeldahl J (1883) Neue Methode zur Bestimmung des Stickstoffs in organischen Körpern. *Fresenius, Zeitschrift f. anal. Chemie* **22**, 366–382.
26. Folch J, Lees M & Stanley GHS (1957) A SIMPLE METHOD FOR THE ISOLATION AND PURIFICATION OF TOTAL LIPIDES FROM ANIMAL TISSUES. *Journal of Biological Chemistry* **226**, 497–509.

27. Roy J, Baranek E & Marandel L (2023) Characterization of free fatty acid receptor family in rainbow trout (*Oncorhynchus mykiss*): towards a better understanding of their involvement in fatty acid signalisation. *BMC Genomics* **24**, 130.
28. Pfaffl MW (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Research* **29**, 2003–2007.
29. Besnard P, Passilly-Degrace P & Khan NA (2016) Taste of Fat: A Sixth Taste Modality? *Physiological Reviews* **96**, 151–176.
30. Velasco C, Otero-Rodiño C, Comesaña S, et al. (2017) Hypothalamic mechanisms linking fatty acid sensing and food intake regulation in rainbow trout. *Journal of Molecular Endocrinology* **59**, 377–390.
31. Panserat S, Kolditz C, Richard N, et al. (2008) Hepatic gene expression profiles in juvenile rainbow trout (*Oncorhynchus mykiss*) fed fishmeal or fish oil-free diets. *Br. J. Nutr.* **100**, 953–967.
32. Welker T, Barrows F, Overturf K, et al. (2016) Optimizing zinc supplementation levels of rainbow trout (*Oncorhynchus mykiss*) fed practical type fishmeal- and plant-based diets. *Aquacult Nutr* **22**, 91–108.
33. S. Kaushik, J. Cravedi, J. Lalles, J. Sumpter, B. Fauconneau, et al. Partial or total replacement of fish meal by soybean protein on growth, protein utilization, potential estrogenic or antigenic effects, cholesterolemia and flesh quality in rainbow trout, *Oncorhynchus mykiss*. *Aquaculture*. 133 (1995) 257–274. .
34. P. Gómez-Requeni, M. Mingarro, J. Calduch-Giner , F. Médale, S. Martin, et al. Protein growth performance, amino acid utilisation and somatotrophic axis responsiveness to fish meal replacement by plant protein sources in gilthead sea bream (*Sparus aurata*). *Aquaculture* 232 (2004) 493–510. .
35. Roy J, Mercier Y, Tonnet L, et al. (2020) Rainbow trout prefer diets rich in omega-3 long chain polyunsaturated fatty acids DHA and EPA. *Physiology & Behavior* **213**, 112692.
36. Kasumyan AO & Døving KB (2003) Taste preferences in fishes. *Fish and Fisheries* **4**, 289–347.
37. Comesaña S, Conde-Sieira M, Velasco C, et al. (2020) Oral and pre-absorptive sensing of amino acids relates to hypothalamic control of food intake in rainbow trout. *J Exp Biol* **223**, jeb221721.
38. Kohbara J & Caprio J (2001) Taste responses of the facial and glossopharyngeal nerves to amino acids in the rainbow trout. *Journal of Fish Biology* **58**, 1062–1072.

39. Jones KA (1990) Chemical requirements of feeding in rainbow trout, *Oncorhynchus mykiss* (Walbaum); palatability studies on amino acids, amides, amines, alcohols, aldehydes, saccharides, and other compounds. *Journal of Fish Biology* **37**, 413–423.
40. Roper SD & Chaudhari N (2017) Taste buds: cells, signals and synapses. *Nat Rev Neurosci* **18**, 485–497.
41. Hichami A, Khan AS & Khan NA (2021) Cellular and Molecular Mechanisms of Fat Taste Perception. In *The Pharmacology of Taste*, vol. 275, pp. 247–270 [Palmer RK, Servant G, editors]. Cham: Springer International Publishing.
42. Gorski JN, Pachanski MJ, Mane J, et al. (2017) GPR40 reduces food intake and body weight through GLP-1. *American Journal of Physiology-Endocrinology and Metabolism* **313**, E37–E47.
43. Sleeth ML, Thompson EL, Ford HE, et al. (2010) Free fatty acid receptor 2 and nutrient sensing: a proposed role for fibre, fermentable carbohydrates and short-chain fatty acids in appetite regulation. *Nutr. Res. Rev.* **23**, 135–145.
44. Sclafani A, Zukerman S & Ackroff K (2013) GPR40 and GPR120 fatty acid sensors are critical for postoral but not oral mediation of fat preferences in the mouse. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* **305**, R1490–R1497.
45. Ichimura A, Hirasawa A, Poulain-Godefroy O, et al. (2012) Dysfunction of lipid sensor GPR120 leads to obesity in both mouse and human. *Nature* **483**, 350–354.
46. Ozdener MH, Subramaniam S, Sundaresan S, et al. (2014) CD36- and GPR120-Mediated Ca²⁺ Signaling in Human Taste Bud Cells Mediates Differential Responses to Fatty Acids and Is Altered in Obese Mice. *Gastroenterology* **146**, 995–1005.
47. Auguste S, Fisette A, Fernandes MF, et al. (2016) Central Agonism of GPR120 Acutely Inhibits Food Intake and Food Reward and Chronically Suppresses Anxiety-Like Behavior in Mice. *IJNPPY* **19**.
48. Wang F, Liu F, Wang X, et al. (2020) Transcriptomic responses to yam (*Dioscorea oppositifolia* L.) extract dietary supplementation in rainbow trout (*Oncorhynchus mykiss*) liver. *Aquac Res* **51**, 932–945.
49. Velasco C, Conde-Sieira M, Comesaña S, et al. (2020) The long-chain fatty acid receptors FFA1 and FFA4 are involved in food intake regulation in fish brain. *Journal of Experimental Biology* **223**, jeb.227330.

50. Dramane G, Abdoul-Azize S, Hichami A, et al. (2012) STIM1 regulates calcium signaling in taste bud cells and preference for fat in mice. *J. Clin. Invest.* **122**, 2267–2282.
51. Shanmugamprema D, Muthuswamy K, Subramanian G, et al. (2020) Fat taste signal transduction and its possible negative modulator components. *Progress in Lipid Research* **79**.
52. Calo J, Blanco AM, Comesaña S, et al. (2021) First evidence for the presence of amino acid sensing mechanisms in the fish gastrointestinal tract. *Sci Rep* **11**, 4933.
53. Leigh S-J & Morris MJ (2018) The role of reward circuitry and food addiction in the obesity epidemic: An update. *Biol Psychol* **131**, 31–42.
54. Narukawa M1. Physiological responses to taste signals of functional food components. *Biosci Biotechnol Biochem.*
55. Cox DN, Hendrie GA & Carty D (2016) Sensitivity, hedonics and preferences for basic tastes and fat amongst adults and children of differing weight status: A comprehensive review. *Food Quality and Preference* **48**, 359–367.
56. Galen KA, Horst KW & Serlie MJ (2021) Serotonin, food intake, and obesity. *Obesity Reviews* **22**.
57. Tavares GA, do Amaral Almeida LC, de Souza JA, et al. (2020) Early weaning leads to disruption of homeostatic and hedonic eating behaviors and modulates serotonin (5HT) and dopamine (DA) systems in male adult rats. *Behavioural Brain Research* **383**, 112531.
58. Bacqué-Cazenave J, Bharatiya R, Barrière G, et al. (2020) Serotonin in Animal Cognition and Behavior. *IJMS* **21**, 1649.
59. Díaz-Rúa A, Chivite M, Comesaña S, et al. (2022) The Opioid System in Rainbow Trout Telencephalon Is Probably Involved in the Hedonic Regulation of Food Intake. *Front. Physiol.* **13**.
60. Roy J, Larroquet L, Surget A, et al. (2020) Impact on cerebral function in rainbow trout fed with plant based omega-3 long chain polyunsaturated fatty acids enriched with DHA and EPA. *Fish & Shellfish Immunology* **103**, 409–420.




Figure 1. Zootechnical parameters of rainbow trout (RT) fed a control (C) or vegetable (V) diet from first feeding to eight months.

Comparison of (A) body weight, (B) feed intake (FI, g feed/kg^{0.8} d⁻¹), and (C) feed efficiency (FE, g of BW/g of feed) per 3-week period for RT fed the C or V diet from first feeding to eight months (days 238). An asterisk (B and C) indicates a significant difference between the dietary treatments as determined by one-way ANOVA (repeated measures) ($p < 0.05$). Results are expressed in g as means \pm SEM ($n = 5$ tanks).




Figure 2. mRNA level of *ffar* at the end of the 8-month feeding period, 20 min and 4 h after the last meal.

The mRNA levels of genes of the *ffar* family (*ffar1*, *ffar2a1a*, *ffar2a2*, *ffar2b1.1*, *ffar2b1.2*, *ffar2b2a*, *ffar2b2b1*, *ffar2b2b2*) in tongue tissue of RT fed either the C or V diet were examined after a duration of feeding of eight months. Gene expression analyses were performed at two time points, measured by RT-qPCR 20 min and 4 h after the last meal. Data are shown as means \pm SEM; fold change $2^{-\Delta\Delta C_t}$ vs. C diet 20 min after feeding for all genes. Two-way ANOVA followed by one-way ANOVA when the interaction was significant, followed by Tukey's post-hoc test; different letters indicate a significant difference ($p < 0.05$) between diet and time ($n = 7$). ns: not significant.




Figure 3. mRNA levels and protein levels of markers of the calcium-signaling pathway evaluated at the end of the 8-month feeding period, 20 min and 4 h after the last meal.

(A–C, D, E, G, H and J) mRNA levels of genes involved in the calcium-signaling pathway (*plcb2*, *plcb3*, *plcb4*, *stim1a*, *stim1b*, *orai1a*, *orai1b* and *serca3a*) in tongue tissue of RT fed the C or V diet for eight months and measured by RT-qPCR 20 min and 4 h after the last meal. Relative gene expression measured using RT-qPCR. Data are shown as means \pm SEM; fold change $2^{-\Delta\Delta C_t}$ vs. the C diet 20 min after feeding for all genes. Two-way ANOVA followed by one-way ANOVA when the interaction was significant, followed by Tukey's post-hoc test; different letters indicate a significant difference ($p < 0.05$) between diet and time ($n = 7$).

(F and I) Protein abundance 4 h after feeding of STIM1 and ORAI1 measured by western blotting in tongue tissue of RT fed the C or V diet for eight months. Data are shown as means \pm SEM; fold change $2^{-\Delta\Delta C_t}$ vs. the C diet. *t*-test was performed ($n = 7$). ns: not significant.




Figure 4. Influence of C and V diets on serotonin-signaling pathways in tongue tissue.

(A) Brain concentrations 20 min after feeding of indolamines (serotonin, 5-HIAA and 5HIAA:serotonin turnover ratio in tongue tissue of RT fed the C or V diet for 8 months. Data are shown as means \pm SEM; fold change compared to the C diet (n=9). An asterisk indicates a significant difference between dietary treatments determined by *t*-test ($p < 0.05$).

(B) mRNA levels of genes involved in metabolic pathways of indolamines (*tph1a*, *tph1b*, *tph2*, *5ht1aa*, *5ht1ab*, *sert*, *vmat2*, *mao*) in myelencephalon tissue of RT fed the C or V diet for eight months (fold change compared to the C diet for each gene) and measured by RT-PCR 20 min and 4 h after the last meal. Data are shown as means \pm SEM; fold change compared to the C diet (n=7). An asterisk indicates a significant difference between dietary treatments determined by *t*-test ($p < 0.05$).

(C-D) Protein abundance 4 h after feeding of TPH2 and VMAT2 measured by western blotting of tongue tissue of RT fed the C or V diet for eight months. Data are shown as means \pm SEM; fold change compared to the C diet (n=7). An asterisk indicates a significant difference between dietary treatments determined by *t*-test ($p < 0.05$).

(D) Enzymatic activity 20 min after feeding of MAO A and B measured via luminescence of tongue tissue of RT fed the C or V diet for eight months. Data are shown as means \pm SEM; fold change compared to the C diet (n=9). An asterisk indicates a significant difference between dietary treatments determined by *t*-test ($p < 0.05$).




Figure 5. mRNA levels of FI markers in hypothalamus at the end of the 8-month feeding period, and 4 h after the last meal.

mRNA levels of genes involved in FI markers (*npya*, *npyb*, *agrp1*, *agrp2*, *pomca*, *pomcb*, *cartpt*, and *mc4ra*) in the hypothalamus of RT fed the C or V diet and measured by RT-qPCR 20 min and 4 h after the last meal. Data are shown as means \pm SEM; fold change $2^{-\Delta\Delta Ct}$ vs. the C diet. *t*-test was performed ($n = 7$).

Table 1. Ingredients and composition of the experimental diets as a function of fish mass/age.

Diet	0 g - 20 g		20 g - 8 months	
	Control diet	Vegetable diet	Control diet	Vegetable diet
Fish meal	27.0	0.0	27.0	0.0
Extruded whole wheat	8.0	3.8	22.5	17.0
Corn gluten	13.0	27.2	8.7	17.0
Wheat gluten	17.4	29.0	7.0	17.0
Peas meal	3.0	2.46	3.0	6.0
Extruded Peas	7.0	6.23	7.0	9.5
White lupin seed meal	2.5	2.0	2.5	4.5
Rapeseed meal	0.6	3.5	0.8	3.0
Soy lecithin	0.0	2.0	0.0	2.0
L-Lysine	0.5	1.2	0.5	1.7
L-methionine	0.0	0.4	0.0	0.4
CaHPO ₄ .2H ₂ O	0.0	1.7	0.0	1.4
Mineral premixa	1.0	1.0	1.0	1.0
Vitamin premixb	1.0	1.0	1.0	1.0
Fish oil	4.8	0.0	4.8	0.0
Palm oil	2.0	5.0	2.0	5.0
Sunflower oil	4.9	2.7	4.9	2.7
Linseed oil	0.0	10.8	0.0	10.8
DHA marine oil (from Polaris) ^c	7.3	0.0	7.3	0.0
Composition (% of DM)				
Dry matter (%)	96.15	94.25	97.30	96.55
Crude protein	53.76	54.40	42.62	41.44
Crude lipid	21.8	21.90	22.64	22.2
Starch	10.84	10.70	17.36	17.74
Ash	7.40	5.02	7.04	4.47
Energy (kJg ⁻¹ DM)	24.75	25.29	24.29	24.81

^aMineral premix: (g or mg kg⁻¹ diet): calcium carbonate (40% Ca), 2.15 g; magnesium oxide (60% Mg), 1.24 g; ferric citrate, 0.2 g; potassium iodide (75% I), 0.4 mg; zinc sulfate (36% Zn), 0.4 g; copper sulfate (25% Cu), 0.3 g; manganese sulfate (33% Mn), 0.3 g; dibasic calcium phosphate (20% Ca, 18% P), 5 g; cobalt sulfate, 2 mg; sodium selenite (30% Se), 3 mg; KCl, 0.9 g; NaCl, 0.4 g (UPAE, INRAE). ^b Vitamin premix: (IU or mg kg⁻¹ diet): DL- α -tocopherol acetate, 60 IU; sodium menadione bisulfate, 5 mg; retinyl acetate, 15,000 IU; DL-cholecalciferol, 3000 IU; thiamin, 15 mg; riboflavin, 30 mg; pyridoxine, 15 mg; B12, 0.05 mg; nicotinic acid, 175 mg; folic acid, 500 mg; inositol, 1000 mg; biotin, 2.5 mg; calcium pantothenate, 50 mg; choline chloride, 2000 mg (UPAE, INRAE). ^c Omegavie® DHA 70 (Polaris, Quimper, France): concentrated anchovy and sardine oil that contains DHA omega-3 fatty acids in triglyceride form.

Table 2. Fatty acid composition of experimental diets (% of total FA).

	0 g - 20 g		20 g - 8 months	
	Control diet	Vegetable diet	Control diet	Vegetable diet
C12:0	0.10	0.18	0.13	0.22
C14:0	1.70	0.32	1.81	0.37
C15:0	0.03	0.04	0.23	0.03
C16:0	12.20	15.37	12.71	15.93
C17:0	0.22	0.07	0.23	0.07
C18:0	2.84	3.61	2.86	3.69
C20:0	0.31	0.23	0.30	0.24
C22:0	0.30	0.23	0.28	0.24
C24:0	0.10	0.14	0.04	0.12
Sum of saturated fatty acids	18.07	20.20	18.60	20.88
C16:1 ω -7	2.07	0.17	2.08	0.16
C18:1 ω -9	16.72	24.74	17.05	25.24
C20:1 ω -9	1.69	0.24	1.41	0.22
C22:1 ω -9	1.77	0.0	1.38	0.03
Sum of MUFAs	22.30	25.15	21.92	25.65
C18:2 ω -6	20.30	26.34	20.45	24.84
C18:3 ω -6	0.13	0.0	0.08	0.0
C20:2 ω -6	0.17	0.0	0.17	0.0
C20:3 ω -6	0.11	0.0	0.12	0.0
C20:4 ω -6	0.85	0.0	0.83	0.0
C22:2 ω -6	0.05	0.10	0.12	0.05
C22:4 ω -6	0.32	0.0	0.30	0.0
C22:5 ω -6	1.56	0.0	1.55	0.0
Sum of ω -6 LC-PUFAs	24.09	26.61	23.62	24.90
C18:3 ω -3	0.80	27.62	0.71	28.24
C18:4 ω -3	0.43	0.0	0.42	0.0
C20:3 ω -3	0.12	0.0	0.12	0.0
C20:4 ω -3	0.34	0.0	0.31	0.0
C20:5 ω -3	4.08	0.0	3.96	0.0
C21:5 ω -3	0.24	0.0	0.25	0.0
C22:4 ω -3	0.14	0.0	0.07	0.0
C22:5 ω -3	1.96	0.0	1.90	0.0
C22:6 ω -3 (DHA)	26.52	0.19	26.53	0.11
Sum of ω -3 LC-PUFAs	34.67	27.86	34.25	28.36
Sum of ω -3 (EPA + DHA)	30.60	0.19	30.50	0.11
ω -3 (DHA + EPA) / ω -6	1.27	0.0	1.29	0.0

Table 3. Nucleotide sequences of the PCR primers used to evaluate mRNA abundance of transcripts by real-time quantitative PCR.

Transcript	Forward Primer	Reverse Primer	Accession Number
Reference			
<i>eef1a1</i>	TCCTCTTGGTCGTTTCGCTG	ACCCGAGGGACATCCTGTG	ENSOMYG00000038328
<i>krt8b</i>	TGGCTACTCCAGTGGTTTCG	CCGCTACCGGAGCTGTAGTT	ENSOMYG00000057091
FFAR receptor			
<i>ffar1</i>	ACTGTTGCACCTGAGTCTGG	GCTGGTCCTGGGTGAAGTTC	ENSOMYG00000041396
<i>ffar2a1a</i>	CCGAGTTCCTCTGCTCCATC	TAGGTGATGGGGAAGGCAAC	ENSOMYG00000004986
<i>ffar2a2</i>	GACAACTTCACCCAGGAGCA	AGCAGAACACACAGGCCAG	ENSOMYG00000030315
<i>ffar2b1.1</i>	TTTTCCACACACAGTTGGCC	AGGTAGTGTTGTCGGCATCT	ENSOMYG00000041393
<i>ffar2b1.2</i>	GTGTGGCCTTCCCTATCAGA	GCAGGGCACAATGTACACAA	ENSOMYG00000041387
<i>ffar2b2a</i>	CCCATCCAACACTCGCTGAA	TGATGACGACGATGCTCAGG	ENSOMYG00000030493
<i>ffar2b2b1</i>	TGACCGCAATCAGTGTGCGAA	CCCAGAAGAAGACGCTAGCC	ENSOMYG00000030500
<i>ffar2b2b2</i>	GTCCAGTACCATCAACGCCA	CTGCACACTCTCCAACAGGGT	ENSOMYG00000005604
Calcium-signaling pathways			
<i>plcb2</i>	TTATCACCCACGGCTTCACC	GGAGAGGATGACGGGGTACT	XM_021584705.1
<i>plcb3</i>	ATAGTGGACGGCATCGTAGC	TGTGTCAGCAGGAAGTCCAA	XM_021577635.1
<i>plcb4</i>	CCGAGGCGTCTGAGGATAAC	GTAGGGGTGGATGTTGGTGG	XM_021600840.1
<i>stim1a</i>	GAACCTTGACCCAGCATCA	GGAGATTGCAGAGTACCGGG	XM_021588396.1
<i>stim1b</i>	CGGACGTGTACAACCTGGACA	CCTTCCGTCCAGATTGTGCT	XM_021563300.1
<i>orai1a</i>	CTATGGTGGAGGTGCAGCTG	TCATCAGGGCGAAAAGGTGC	XM_021621327.1
<i>orai1b</i>	TGAAGGCTTCCAGTCGAACG	GGCACTGAAGGCAATGAGGA	XM_021602671.1
<i>atp2a3a (serca3a)</i>	GCTGTTATCACCACTGCCT	AGATGACAGAAGTGCAGCCC	XM_021624243.1
Indolamine metabolism			
<i>tph1a</i>	ACACCAGAGCCAGACACATG	TCATCTGAAGCTCCGAGGGA	XM_021598622.1
<i>tph1b</i>	AGCGTCCGTTTACAGTGAGG	GCCCACGATGTCCAGTTCAT	XM_021598622.1

<i>tph2</i>	AGCACCTCAAAGACCACGTC	ACTGGTCGAGCTCTGCAATC	MG015698
<i>5ht1aa</i>	CCCAACACTCCACAGTCCTC	ACCGAGCGTCTTTACCGTTT	XM_021622104
<i>5ht1ab</i>	GAGGACCAACGGGGACCCGA	AATCGCCGTGCTTGACCGCA	CCAF0100015582
<i>sert</i>	CCTGCTGCCCTACATGTTGA	GGGGCAGATGTGTTTCCAGA	M_021582096.1
<i>vmat2</i>	TCGCCTTGCTGCTGGATAAT	GAGGCCGTGGTGTGATTCTT	XM_021609975.1
<i>mao</i>	GGTACGTACAAGCAGTGTGGA	GCTCCTTGGCCATACTCA	XP_021434550.1
Neuropeptides			
<i>npyl1a</i>	AAGGCAGAGGTGAGTGCTGT	AGCCTGTGGCTCACTAATCAA	
<i>npyl1b</i>	CGTATTTGTTAGCGCCTTTC	TCATTCGATTTCTTCACTTCCA	NM_001124266
<i>agrp1</i>	ACCAGCAGTCCTGTCTGGGTAA	AGTAGCAGATGGAGCCGAACA	NM_001124266
<i>agrp2</i>	CCAGGAGACGGATTTTGCCA	GAGGGGCACAGCTATCCATC	genbank CR376289
<i>pomca</i>	ACAAGATGAACCACTTCCGCT	TGAACAACGTGAGCAGTGGT	genbank CA343080
<i>pomcb</i>	AGAGCTCCCACCATCTCGT	GCCGCCCATCAAAGTCTAT	NM_001124718.1
<i>cartpt</i>	ACCATGGAGAGCTCCAG	GCGCACTGCTCTCCAA	NM_001124719.1
<i>mc4ra</i>	CCACAACATCATGACCACGC	AGGCAGATGAGGAGCGGTAGT	DQ836925
	CCACAACATCATGACCACG	AGGCAGATGAGGAGCGGTAGT	XM_021613317.1
