



HAL
open science

Characterization of enological oak tannin extracts by multi-analytical methods approach

Kevin Pascotto, H el ene Hall e, Aude A. Watrelot, Aurelie Roland, Emmanuelle Meudec, Pascale Williams, St ephanie Carrillo, Bertrand Robillard, Nicolas Sommerer, C eline Poncet-Legrand, et al.

► To cite this version:

Kevin Pascotto, H el ene Hall e, Aude A. Watrelot, Aurelie Roland, Emmanuelle Meudec, et al.. Characterization of enological oak tannin extracts by multi-analytical methods approach. Congress of the American Society of Enology and Viticulture 2023, Jun 2023, Napa, Micronesia. hal-04167956

HAL Id: hal-04167956

<https://hal.inrae.fr/hal-04167956>

Submitted on 21 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.

Title :

Characterization of enological oak tannin extracts by multi-analytical methods approach

Oak tannin extracts are commonly used to improve wine properties. The main polyphenols found in oak wood extracts are ellagitannins that release ellagic acid upon hydrolysis and comprise numerous structures. Moreover, oak tannin extracts contain other compounds giving a complex mixture. Consequently, the official OIV method based on gravimetric analysis of the tannin fraction adsorbed on polyvinylpyrrolidone is not sufficient to describe their composition and highlight their chemical diversity.

Eight commercial oak tannins were characterized by a combination of analytical approaches, Polyphenols were analyzed using the official OIV method, UV spectrophotometry, UPLC-UV-MS analysis before and after acidic methanolysis, and HPLC-SEC-UV. Neutral sugars and polyols were determined as alditol acetates by GC-FID analysis, before and after hydrolysis. Finally, samples were compared by a non-targeted metabolomic approach based on UHPLC-HRMS/MS.

Gravimetric analysis, absorbance values at 280 nm, and the quantities of ellagic acid released by methanolysis revealed some differences between samples, indicating variations in their tannin composition. This was confirmed by SEC analysis evidencing differences in tannin size distribution, particularly in larger polymer content.

All samples contained significant quantities of sugars, and in particular xylose, mostly found in the linked form, and of quercitol, a polyol marker of oak origin. These compounds contributed to up to 25% of the whole extract composition, the proportions of free and combined sugars and polyols also showing large variations between tannins.

Non targeted UPLC-HRMS analysis detected major ellagitannins such as vescalagin, castalagin, and roburins A-E, but also a large number of derivatives as well as other molecules such as lignans and quercotriterpenosides, and highlighted large differences between samples.

This work demonstrates the variability in the composition of commercial oak tannin extracts, likely to impact their properties, and emphasizes the need for detailed multi-method characterization in the frame of quality control and selection of tannins for specific applications.

Authors and affiliations :

Kevin Pascotto

Institut Œnologique de Champagne, Epernay, France

H l ne Halle

SPO, INRAE, Univ Montpellier, Montpellier SupAgro, Montpellier, France

Aude Watrelot

SPO, INRAE, Univ Montpellier, Montpellier SupAgro, Montpellier, France

Iowa State University, Department of Food Science and Human Nutrition, Ames, USA

Aur lie Roland

SPO, INRAE, Univ Montpellier, Montpellier SupAgro, Montpellier, France

Emmanuelle Meudec

SPO, INRAE, Univ Montpellier, Montpellier SupAgro, Montpellier, France

Pascale Williams

SPO, INRAE, Univ Montpellier, Montpellier SupAgro, Montpellier, France

St phanie Carrillo

SPO, INRAE, Univ Montpellier, Montpellier SupAgro, Montpellier, France

Bertand Robillard

Institut Œnologique de Champagne, Epernay, France

Nicolas Sommerer

SPO, INRAE, Univ Montpellier, Montpellier SupAgro, Montpellier, France

Céline Poncet-Legrand

SPO, INRAE, Univ Montpellier, Montpellier SupAgro, Montpellier, France

Véronique Cheynier

SPO, INRAE, Univ Montpellier, Montpellier SupAgro, Montpellier, France