



The role of cell wall polysaccharides in the extraction of anthocyanins and tannins: results, perspectives of a more positive contribution

Jean-Claude Boulet, Elissa Abi Habib, Stéphanie Carrillo, Stéphanie Roi, Frédéric Veran, Arnaud Verbaere, Emmanuelle Meudec, Anais Rattier, Marie-Agnès Ducasse, Bodil Jørgensen, et al.

► To cite this version:

Jean-Claude Boulet, Elissa Abi Habib, Stéphanie Carrillo, Stéphanie Roi, Frédéric Veran, et al.. The role of cell wall polysaccharides in the extraction of anthocyanins and tannins: results, perspectives of a more positive contribution. OENO Macrowine 2023, Jul 2023, Bordeaux, France. hal-04168521

HAL Id: hal-04168521

<https://hal.inrae.fr/hal-04168521>

Submitted on 21 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Topic: Wine chemistry, wine components

Title: The role of cell wall polysaccharides in the extraction of anthocyanins and tannins: results, perspectives of a more positive contribution.

Authors:

Jean-Claude Boulet^{a,b}, Elissa Abi-Habib^a, Stéphanie Carrillo^a, Stéphanie Roi^a, Frédéric Veran^a, Arnaud Verbaere^{a,b}, Emmanuelle Meudec^{a,b}, Anais Rattier^{a,b}, Marie-Agnès Ducasse^c, Bodil Jorgensen^d, Jeanett Hansen^d, Sophie Le Gall^{e,f}, Céline Poncet-Legrand^a, Véronique Cheynier^{a,b}, Thierry Doco^a, Aude Vernhet^a.

^a Univ. Montpellier, SPO, INRAE, Institut Agro Montpellier Supagro, 34070 Montpellier, France

^b INRAE, PROBE research infrastructure, PFP polyphenols analysis facility, 34070 Montpellier, France

^c IFV, experimental unit of Pech Rouge, 11430 Gruissan, France

^d Department of Plant and Environmental Sciences, University of Copenhagen, 1870 Frederiksberg, Denmark

^e INRAE, UR BIA, 44316 Nantes, France

^f INRAE, PROBE research infrastructure, BIBS biopolymers analysis facility, 44316 Nantes, France

Keywords: polyphenols, cell wall material, binding, CoMPPs

Abstract.

The composition of grape berry cell walls was studied on two grape varieties, two years and two maturation levels at the same time as the extraction of anthocyanins and tannins. The chemical composition of skins, seeds, and pulps, focused on polyphenols and polysaccharides, was compared to the chemical composition in polyphenols after extraction from the skins in model solutions or after wine making of the berries. Polyphenols were mainly characterized by UPLC-MS and HPLC-SEC. Polysaccharides were characterized by analysis of the neutral sugar compositions, and also by the CoMPP (comprehensive micropolymer profiling) analysis, a new method which targets the functional groups of cell wall polysaccharides.

The extractions rates showed huge differences between the non acylated and the para-coumaroylated anthocyanins. The former were much easier to extract than the latter. Particularly in model solutions, the extraction of p-coumaroylated anthocyanins was almost negligible. The extraction rate of tannins was between those of the two anthocyanin families. Moreover, in wines as in model solutions, the final concentrations in tannins, non acylated and p-coumaroylated anthocyanins showed correlations that did not exist in the berry compositions, suggesting a similar mechanism of extraction associating those three families of polyphenols. According to the CoMPPs, these mechanisms would mainly rely on polysaccharidic families, namely hemicelluloses, homogalacturonans, rhamnogalacturonans, and extensins.

The major role of the cell wall polysaccharides in the extraction of tannins and anthocyanins was confirmed. CoMPPs revealed a much more complex mechanism than expected, e.g. homogalacturonans in skins and pulps associated to an increase and a decrease of the polyphenols extractibilities, respectively. Moreover, our study changed the standpoint on cell wall polysaccharides. Up to now, they were considered as detrimental since they bind polyphenols, and were thus expected to increase losses. But they also release soluble polysaccharides (PRAGs) which contribute positively to the colloidal stability of wines.