



**HAL**  
open science

# Importance of DNA-Methylation in Entomopathogenic Bacteria with similar lifestyle

Julien Brillard

► **To cite this version:**

Julien Brillard. Importance of DNA-Methylation in Entomopathogenic Bacteria with similar lifestyle. 10th FEMS congress of European Microbiologists, Jul 2023, Hamburg (D), Germany. hal-04171736

**HAL Id: hal-04171736**

**<https://hal.inrae.fr/hal-04171736>**

Submitted on 26 Jul 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Importance of DNA-Methylation in Entomopathogenic Bacteria with similar lifestyle

### Abstract:

DNA methylation plays various roles in bacteria and can contribute to bacterial interactions with their hosts, through diverse mechanisms such as genome maintenance or epigenetic regulations. The bacteria *Xenorhabdus* and *Photorhabdus* are pathogenic for insects and are found in the gut of soil nematodes with whom they are symbiotically associated. The roles of a conserved DNA-Methyltransferase (MTase) were investigated in these bacteria using deregulated strains and methylomic approaches.

Great diversity was observed in the distribution of MTase-encoding genes, and a single MTase (Dam, for DNA-adenine MTase) was identified in all the species of these two genera. Methylome analysis showed that the GATC motifs recognized by Dam reached a high methylation rate (>99%) in the studied strains. Enrichment of unmethylated motifs in promoter regions was observed and may reveal mechanisms of epigenetic regulation. The Dam overexpression was associated with a reduced swimming motility, caused by a downregulation of flagellar genes, in both *Xenorhabdus* and *Photorhabdus*. In contrast, alteration of various major phenotypes such as hemolysis or virulence depended on the species studied.

Our work revealed the diverse roles played by a conserved DNA MTase during the life cycle of entomopathogenic bacteria.