



**HAL**  
open science

## On a potential involvement of RG-I pectins in the generation of maturation stress in the secondary cell wall of poplar tension wood fibres

Françoise Laurans, Véronique Lainé-Prade, Marie Capron, Olivier Arnould, Amra Šećerović, Tancrede Alméras, Bruno Clair, Annabelle Déjardin, Gilles Pilate

### ► To cite this version:

Françoise Laurans, Véronique Lainé-Prade, Marie Capron, Olivier Arnould, Amra Šećerović, et al.. On a potential involvement of RG-I pectins in the generation of maturation stress in the secondary cell wall of poplar tension wood fibres. 13èmes journées du Réseau Français des Parois, May 2022, Versailles (FR), France. pp.1-2. hal-04171845


**HAL Id: hal-04171845**

**<https://hal.inrae.fr/hal-04171845>**

Submitted on 26 Jul 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# 13èmes journées du Réseau Français des Parois

18 au 20 mai 2022 - Versailles

## Session 1- Biosynthèse, architecture, composition des parois

### On a potential involvement of RG-I pectins in the generation of maturation stress in the secondary cell wall of poplar tension wood fibres

Françoise Laurans<sup>1</sup>, Véronique Lainé-Prade<sup>1</sup>, Marie Capron<sup>2,3</sup>, Olivier Arnould<sup>2</sup>, Miyuki Takeuchi<sup>1,4</sup>, Amra Secerovic<sup>1,5</sup>, Tancrede Almèras<sup>2</sup>, Bruno Clair<sup>2</sup>, Annabelle Déjardin<sup>1</sup>, Gilles Pilate<sup>1</sup>

<sup>1</sup> BioForA, INRA ONF, 45000 Orléans, France ; <sup>2</sup> LMGC, CNRS, Montpellier University, 34000 Montpellier, France; <sup>3</sup> Michelin Campus RDI, Rue Bleue, 63118 Cébazat, France; <sup>4</sup> The Graduate School of Engineering, The University of Tokyo, Tokyo, Japan; <sup>5</sup> AO Research Institute Davos, AO Center Davos, 7270 Davos, Switzerland

Tension wood (TW) produced by temperate hardwood trees is known for its outstanding mechanical properties. The mechanism that generates the tensile force responsible for TW properties is still not elucidated: it may originate from a high porosity hydrogel identified in the G-layer, an extra cell wall layer present in TW fibres. Further, it has been proposed that the hydrolysis of pectin rhamnogalacturonan-I (RG-I) side chains by a  $\beta$ -galactosidase was responsible for the formation of this gel. To test this hypothesis, we used as a proxy of TW fibre maturation, the files of fibres located from the cambium toward the pith to compare both the kinetics of RG-I side chain hydrolysis by immunolocalisation and the evolution of G-layer indentation modulus, a proxy of its stiffness, by Atomic Force Microscopy (Arnould et al., 2021). We found a good correlation between the kinetics of global  $\beta$ -galactosidase activities and the labeling profiles of LM5 (specific to RG-I side chains) and RU1 (specific to RG-I backbone) antibodies that may reflect RG-I side-chain hydrolysis. However, it appeared that this occurred long after the main G-layer stiffening that took place earlier when G-layers were still thin, but it correlated with a second minor increase in G-layer stiffness. The labeling profiles of several other antibodies led to identifying other candidates potentially involved in the building of TW mechanical properties. G-layer stiffening is a complex process that probably involves several molecular actors,  $\beta$ -galactosidases being only one of them.

Arnould O, Capron M, Ramonda M, Laurans F, Almèras T, Pilate G, Clair B. Mechanical characterisation of the developing cell wall layers of tension wood fibres by Atomic Force Microscopy. bioRxiv 2021.09.23.461481; doi: <https://doi.org/10.1101/2021.09.23.461481>

This work was performed in the framework of the project "StressInTrees" (ANR-12-BS09-0004) funded by the French National Research Agency (ANR)