



HAL
open science

Complementary feeding practices are related to the risk of food allergy in the ELFE cohort

T. Adam, A Divaret-chauveau, C. Roduit, Karine Adel-Patient, A. Deschildre, C. Raheison, Marie-Aline Charles, S. Nicklaus, B de Lauzon-guillain

► To cite this version:

T. Adam, A Divaret-chauveau, C. Roduit, Karine Adel-Patient, A. Deschildre, et al.. Complementary feeding practices are related to the risk of food allergy in the ELFE cohort. *Allergy*, 2023, 79 (9), pp.2456-2466. 10.1111/all.15828 . hal-04174304

HAL Id: hal-04174304

<https://hal.inrae.fr/hal-04174304>

Submitted on 31 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Complementary feeding practices are related to the risk of food allergy in the ELFE cohort

T. Adam^{1,2}  | A. Divaret-Chauveau^{2,3}  | C. Roduit^{4,5,6}  | K. Adel-Patient⁷  |
 A. Deschildre⁸ | C. Raheison⁹ | M.-A. Charles^{10,11}  | S. Nicklaus¹²  |
 B. de Lauzon-Guillain¹⁰ 

¹Department of Dermatology and Allergology, University Hospital of Nancy, Vandoeuvre-Lès-Nancy, France

²Department of Pediatric Allergology, University Hospital of Nancy, Vandoeuvre-Lès-Nancy, France

³EA3450 Développement Adaptation et Handicap (DevAH), University of Lorraine, Nancy, France

⁴Division of Respiratory Medicine and Allergology, Department of Paediatrics, Inselspital, University of Bern, Bern, Switzerland

⁵Children's Hospital of Eastern Switzerland, St. Gallen, Switzerland

⁶Christine Kühne-Center for Allergy Research and Education (CK-CARE), Davos, Switzerland

⁷Université Paris-Saclay, CEA, INRAE, Département Médicaments et Technologies pour la Santé (DMTS), Laboratoire d'Immuno-Allergie Alimentaire, Gif-sur-Yvette, France

⁸University of Lille, CHU Lille, Pediatric Pulmonology and Allergy Department, Hôpital Jeanne de Flandre, Lille, France

⁹Department of Pulmonology, CHU of Guadeloupe, French West Indies University, INSERM UMR 1219 EpiCene Team, Bordeaux, France

¹⁰Université Paris Cité and Université Sorbonne Paris Nord, Inserm, INRAE, CRESS, Paris, France

¹¹Unité Mixte Inserm-Ined-EFS Elfe, Ined, Aubervilliers, France

¹²Centre des Sciences du Goût et de l'Alimentation, CNRS, INRAE, Institut Agro, Université Bourgogne Franche-Comté, Dijon, France

Correspondence

T. Adam, CHRU Brabois, Rue du Morvan, Vandoeuvre-Lès-Nancy 54500, France.
 Email: t.adam@chru-nancy.fr

Funding information

Agence Nationale de la Recherche, Grant/Award Number: ANR-19-CE36-0008

Abstract

Background: Conflicting results have been obtained when analyzing the relationship between complementary feeding (CF) practices and allergic diseases in childhood. This study aims to further explore the association between allergic diseases in early childhood ([10.1016/j.jaci.2012.02.036](https://doi.org/10.1016/j.jaci.2012.02.036)) and the age at CF introduction ([10.1016/S0140-6736\(15\)00149-X](https://doi.org/10.1016/S0140-6736(15)00149-X)), food diversity in the first year of life ([10.1016/j.ijporl.2019.109759](https://doi.org/10.1016/j.ijporl.2019.109759)) and the delayed introduction of major allergenic foods.

Methods: This analysis focused on 6662 children from the French nationwide ELFE cohort. Data on feeding practices were collected monthly from 3 to 10 months old. Their age at CF introduction was calculated alongside a diversity score, and the number of major allergenic foods (out of eggs, fish, wheat, and dairy products) not introduced at 8 and 10 months. Their associations with parent-reported allergy-related health events between 1 and 5.5 years were assessed using logistic regressions adjusted for confounding factors. A sensitivity analysis excluding early allergic cases (occurring between 2 months and 1 or 2 years) was conducted.

Abbreviations: CF, complementary feeding; ELFE, Étude Longitudinale Française depuis l'Enfance; PASTURE, protection against allergy; STUdy in rural environment.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.


© 2023 The Authors. *Allergy* published by European Academy of Allergy and Clinical Immunology and John Wiley & Sons Ltd.

Results: Late CF (>6 months) was related to a higher risk of food allergy (OR [95% CI] = 1.35 [1.02; 1.78]), a low diversity score at 8 months to a higher risk of asthma (OR [95% CI] = 1.22 [1.01; 1.48]), and two allergenic foods or more not being introduced at 10 months to a higher risk of rhinoconjunctivitis (OR [95% CI] = 1.20 [1.00; 1.44]) and food allergy (OR [95% CI] = 2.46 [1.77; 3.42]). Only this last association remained significant after the exclusion of early cases.

Conclusion: The delayed introduction of major allergenic foods is related to a higher risk of food allergy, which supports the updated guidelines for allergy prevention.

KEYWORDS

allergic rhinitis, asthma, childhood, complementary feeding, food allergy



GRAPHICAL ABSTRACT

The study aimed to explore the association between complementary feeding practices (age, food diversity and allergenic food) and the risk of food allergy, eczema, asthma, and rhinoconjunctivitis in early childhood. More than 50% children were not introduced with at least one major allergenic food (egg, wheat, fish, dairy products) at 10 months. This delayed introduction (after 10 months) of major allergenic food is associated with a higher risk of food allergy up to the age of 5.5 years.

1 | INTRODUCTION

Many risk factors for allergic diseases have been identified, but some of them, such as gender, a family history of atopy, and pre-existing atopic diseases, are not modifiable.¹⁻³ Nevertheless, other risk factors are modifiable and they are far more interesting to those looking to bolster primary prevention; these include active or passive smoking, feeding practices¹⁻³ [particularly the age at which complementary feeding (CF) is initiated] and food diversity during this period. Non-modifiable factors such as, a family history of atopy^{4,5} or early signs of allergic diseases,^{6,7} further influence allergic disease through modifications to CF practices.

Current European guidelines define early CF as the introduction of CF occurring before the age of 4 months whereas late CF is considered to be when the introduction of CF occurred after the age of 6 months.⁸ They recommend offering a varied diet to infant and the introduction of allergenic foods when CF is commenced any time after 4 months.⁸ In most studies, early CF (<4 months) is not related to a higher risk of atopic dermatitis, food allergy, asthma, or

rhinoconjunctivitis.^{4-6,9} In contrast, late CF (after 6-7 months) is associated with a higher risk of food allergy after 5 years of age¹⁰ but it is not associated with asthma, atopic dermatitis, or allergic rhinitis in the majority of studies.¹¹⁻¹³

Some studies suggest that increased food diversity in the first year of life is linked to a lower risk of having food allergy,^{11,13} asthma,^{13,14} atopic dermatitis,^{6,14} or allergic rhinitis.¹⁴ However, other research reports that there is no association with atopic dermatitis,^{9,12} asthma,⁹ or allergic rhinitis.^{9,14}

When considering the introduction of specific food allergens, the early introduction of peanuts in high-risk infants reduces the risk of a peanut allergy at 5 years of age but it does not affect other food allergies or allergic diseases.¹⁵ Results are more conflicting when it comes to other food allergens, such as eggs, dairy products, and wheat, with some studies showing that there is a protective effect when introducing allergens at the beginning of CF on the risk of food allergy,^{14,16} while other analysis found no association.^{10,17-19} There is no link to the other allergic diseases.^{6,9,12,18-20} A recent study highlighted the benefits of the early

introduction (between 3 and 4 months) and regular consumption up to 36 months, of four allergenic foods (peanut, cow's milk, egg and wheat), on the development of an allergy to these foods.²¹ Hence, evidence concerning the relationship between CF practices and the risk of allergic diseases remains limited, and findings need to be confirmed in larger studies that provide more detailed information on the influence of CF.

In this context, our study aims to explore the link between several dimensions of CF practices (age at initiation, food diversity, introduction of main food allergens) and the risk of allergic diseases (food allergy, eczema, asthma, and rhinoconjunctivitis) up to 5.5 years of age among a large birth cohort in France.

2 | METHODS

2.1 | Study population

The Étude Longitudinale Française depuis l'Enfance (ELFE) is a multi-disciplinary nationwide birth cohort that comprises 18,329 children born in 2011 in a random sample of 349 maternity wards in mainland France.²² The inclusion criteria were single children or twins born after 33 weeks of gestation to mothers aged 18 years or older who were not planning to move outside of metropolitan France in the next 3 years.

2.2 | Ethical statement

Participating mothers had to provide written consent for their own and their child's participation. Fathers signed the consent form for the child's participation when present at inclusion, or they were informed about their rights to oppose it. The ELFE study is approved by the Advisory Committee for Treatment of Health Research Information (Comité Consultatif sur le Traitement des Informations pour la Recherche en Santé), the National Data Protection Authority (Commission Nationale Informatique et Libertés), and the National Statistics Council.

2.3 | Maternal and newborn characteristics

Mothers were interviewed in the maternity ward to obtain medical information about their pregnancy, their newborn, their demographic, their socioeconomic, and lifestyle-related characteristics and their eating habits during pregnancy. This was supplemented with data collection from obstetric and pediatric medical records. At 2 months postpartum, mothers and fathers were interviewed by phone, and more details on demographic and socioeconomic characteristics were collected. Meanwhile, the family's region of residence was determined using their postal code.

Parental and sibling history of allergy was also gathered during the 2-month interview. Children were considered to have a family history of allergy if at least one parent or sibling had experienced an allergic disease.

2.4 | Infant feeding practices

During each interview (monthly from 2 to 10 months), the following feeding practices were recorded: the consumption of solely breast milk, solely infant formula, both breast and formula milk (including plant-based infant formula), animal milk, or a plant-based beverage. The exact age of the child at breastfeeding cessation was recorded, alongside their age when formula milk was introduced, allowing us to calculate the predominant breastfeeding duration.²³

Complementary feeding practices were also collected monthly from 3 to 10 months using a 26-item questionnaire.²⁴ The age at CF initiation was defined as the age in months when foods or beverages other than water, breast milk or infant formula were consumed for the first time more than once a month.²⁵ This was categorized into early (before 4 months), recommended (between 4 and 6 months), and late (after 6 months) strata.

A diversity score was derived from the one used in the PASTURE cohort,¹⁹ and it consisted of the number of food groups (fruit or vegetables, meat, fish, eggs, grains, dairy products, bread and biscuits) that were consumed by the child at a given age. Data regarding consumption of some food items (chocolate, soy, and nuts) collected in the PASTURE study were not available in the ELFE study, so they were not considered in the diversity score. For each month, the diversity score was calculated as a total count of the number of different food groups consumed at least once in a given month, and it varied between 0 (no food diversity) and 8 (the highest food-diversity score). For the present study, the diversity score was considered at 8 and 10 months because of the low variability in food diversity at earlier ages (Figure S1). As we hypothesized that very low diversity was associated with a higher risk of allergy, we defined low diversity as the consumption of 0–2 food groups, intermediate diversity as the consumption of 3–5 food groups, and high diversity as the consumption of 6–8 food groups.

To study the impact of the late introduction of major allergenic foods on allergic diseases, we calculated the number of major allergenic foods (dairy products, hen's eggs, wheat, and fish) that were yet to be introduced at 8 and 10 months. Dairy products were considered to have been introduced if the infant consumed at least one item out of whole, skimmed, or semi-skimmed cow's milk, cheese, or yogurt. Egg was considered introduced if the infant ate the whites or the yolks of eggs. Meanwhile, wheat was considered introduced if the infant consumed at least one item out of pasta, rice, infant cereals, bread, or biscuits.

2.5 | Allergic diseases

Parents reported their children's health data during phone interviews at the ages of 2 months, 1, 2, 3.5, and 5.5 years and each interview covered the period since the last follow-up. Children were considered as prevalent cases and then excluded from all analysis if

parents reported an itchy rash, wheezing, or a medical diagnosis of cow's milk allergy at the 2-month interview.

Children were considered to have wheezing/asthma/eczema, respectively, between 2 months and 5.5 years if the parents reported that the child had at least one wheezing episode(s)/a medical diagnosis of asthma/eczema symptoms during the 1-, 2-, 3.5-, or 5.5-year interviews. As food allergy data were not collected at the 1-year interview, children were considered to have had a food allergy between 2 months and 5.5 years of age if their parents reported that they have been given a medical advice at least once to avoid any type of food due to an allergy at the 2-, 3.5-, or 5.5-year interviews.²⁶ Data about rhinoconjunctivitis were not collected at the 1- and 2-year interviews, so the children were considered to have had rhinoconjunctivitis between the ages of 3.5 and 5.5 years if parents reported at least one episode of allergic conjunctivitis or rhinitis at the 3.5- or 5.5-year interview.

The variable "any allergic disease" was defined as having at least one disease among food allergy, eczema, asthma, or rhinoconjunctivitis between 2 months and 5.5 years.

The children were considered as early cases if eczema or wheezing was reported at the 1-year interview or if a food allergy was reported at the 2-year interview.

2.6 | Sample selection

Analyses were based on the complete case sample, and the selection steps are presented in Figure 1. All of the analyses were conducted after the exclusion of prevalent cases at the age of 2 months.




FIGURE 1 Flow chart of the study's sample selection.

2.7 | Statistical analysis

To compare the included families with their excluded ELFE counterparts, we used chi-squared tests for categorical variables and Student's *t*-tests for continuous variables.

Both unadjusted and adjusted associations between CF characteristics and atopic diseases were assessed using logistic regression models. For the adjusted models, potential confounding factors were identified from the literature and then selected using the direct acyclic graph method.²⁷ These included maternal characteristics (age at delivery, education level, employment status, migration, smoking status during pregnancy, region of residence, and urban/rural area), household characteristics (income per consumption unit, the number of older children at home), infant characteristics (cesarean-section delivery, infant's gender, birth weight, gestational age), predominant breastfeeding duration, family history of allergy, and variables related to the study's design (recruitment wave and maternity unit size) (Figure S2).

To explore the potential moderating effect of some factors, such as family history of allergy, the models also included the interaction term between the family history of allergy and the variables describing the CF practices that were tested. As a significant interaction between a family history of allergy and the child's age at the beginning of CF was only found for eczema, stratified analyses on the family history of allergy were only performed for this particular association.

To explore the issues relating to reverse causation, a sensitivity analysis was conducted after the exclusion of early cases (defined as cases of allergy occurring between the 2 month interview and the

1- or 2-year interviews) as we cannot guarantee that these cases did not occur between the 2-month interview and the beginning of CF. However, this sensitivity analysis also excludes potential incident cases occurring a few months after CF initiation.

To deal with selection and attrition bias, a sensitivity analysis was conducted with weighted data. The weighting was calculated to take the inclusion procedure and biases with regard to non-consent or attrition into account. It also included calibration on the margins from the state register's statistical data and the 2010 French National Perinatal study²⁸ on the following variables: age, region, marital status, migration status, level of education, and primiparity (<https://www.elfe-france.fr/fichier/rte/178/Cote%20recherche/Weighting-Elfe-surveys-general-document.pdf>). A specific weighting was calculated for each subsample.

All analyses were carried out with SAS software version 9.4 (SAS Institute), and the statistical significance was defined as $p < .05$.

3 | RESULTS

3.1 | Descriptive statistics

In comparison with the 6662 children who were included in the study, the excluded children were more likely to be born to a younger mother (30.4 vs 31.4 years) with a lower educational level (16.8% vs 24.8% with a Master's degree) and a lower income (€ 1514 ± 1049 vs € 1790 ± 936 per consumption unit) who smoked throughout their pregnancy (19.2% vs 11.3%). They were also less likely to have a family history of allergy (47.3% vs 52.5%).

The characteristics of the study's population are described in Table 1. The average age at CF initiation was 5.4 months (SD 1.1 months).

Food diversity was low before 6 months of age and increased thereafter up to 10 months (Figure S1). At 8 months, 18.8% of infants had a low diversity score, while 17.0% exhibited a high diversity score. Meanwhile, at 10 months, 2.4% had a low food-diversity score, and 55.6% demonstrated a high diversity score.

At least two major allergenic foods (out of dairy products, fish, wheat and eggs) were not introduced in 43.5% and 10.6% of children at 8 and 10 months, respectively (Figure S3). Additionally, more than half of the children had not been introduced to at least one major allergenic food by the age of 10 months.

3.2 | Age at complementary feeding introduction

As a moderating effect with regard to a family history of allergy (p for interaction = .03) was found on the associations between the age at CF introduction and the risk of eczema, these associations were stratified in accordance with this variable. Compared to the recommended CF initiation (4–6 months), a late CF initiation was associated to a lower risk of eczema among children without a family history of allergy (OR [95% CI] = 0.74, IC [0.57; 0.95]), but this was

not the case among those with a family history of allergy (OR [95% CI] = 1.15, [0.94; 1.40]). Meanwhile, an early CF introduction was related to the risk of eczema neither in children without a family history of allergy (OR [95% CI] = 0.94, [0.69; 1.29]), nor in children with a family history of allergy (OR [95% CI] = 1.16, [0.88; 1.58]). Furthermore, the negative association between a late CF initiation and the risk of eczema in children without a family history of allergy was not anymore significant after the exclusion of early cases (OR [95% CI] = 0.75, [0.54; 1.04]).

TABLE 1 Sample description (n = 6662).

Maternal characteristics	
Age at delivery (years)	31.4 (4.5)
Master's degree	24.8% (1655)
Employed during pregnancy	80.5% (5363)
Household income (€/month/consumption unit)	1790 (936)
Born in country other than France	5.6% (370)
Smoking during pregnancy	
Non smoker	59.2% (3941)
Smoker only before pregnancy	26.2% (1747)
Smoker only in early pregnancy	3.3% (223)
Smoker throughout pregnancy	11.3% (751)
Pre-pregnancy BMI (kg/m ²)	23.3 (4.6)
Rural area	24.4% (1625)
Number of older children	
Firstborn	45.8% (3054)
Secondborn	37.8% (2517)
At least thirdborn	16.4% (1091)
Child's characteristics	
Birthweight (kg)	3.3 (0.5)
Gestational age (weeks)	39.3 (1.4)
Boys	50.8% (3387)
C-section delivery	17.0% (1131)
Predominant breastfeeding duration (months)	2.3 (3.6)
Family history of allergy	52.5% (3499)
Age at complementary feeding introduction	
<4 months	7.5% (499)
4 to 6 months	77.6% (5169)
>6 months	14.9% (994)
Allergic diseases between 2 months and 5.5 years	
Any allergy	64.2% (4279)
Eczema	33.5% (2158)
Food allergy	5.8% (373)
Asthma	26.6% (1770)
Rhinoconjunctivitis	39% (2394)

Note: Values are % (n) or mean (SD).

TABLE 2 Adjusted associations between complementary feeding characteristics and allergic diseases up to 5.5 years.

	Any allergy	Eczema	Food allergy	Asthma	Rhinoconjunctivitis
N	6662	6445	6433	6662	6140
Age at complementary feeding introduction					
<4 months	1.03 [0.84; 1.25]	1.06 [0.86; 1.30]	0.90 [0.58; 1.41]	1.17 [0.95; 1.45]	0.99 [0.81; 1.22]
4–6 months	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]
>6 months	1.03 [0.89; 1.20]	0.95 [0.82; 1.11]	1.35 [1.02; 1.78]	0.96 [0.82; 1.13]	1.06 [0.91; 1.23]
Diversity score at 8 months					
Low diversity (0–2 food groups)	1.09 [0.91; 1.30]	1.01 [0.84; 1.21]	1.50 [1.02; 2.20]	1.22 [1.01; 1.48]	1.09 [0.91; 1.31]
Intermediate diversity (3–5 food groups)	1.09 [0.95; 1.25]	1.02 [0.88; 1.18]	1.32 [0.95; 1.83]	1.09 [0.93; 1.28]	1.03 [0.89; 1.19]
High diversity (6–8 food groups)	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]
Diversity score at 10 months					
Low diversity (0–2 food groups)	1.05 [0.75; 1.46]	1.03 [0.73; 1.45]	1.57 [0.87; 2.85]	1.01 [0.70; 1.45]	1.43 [1.02; 2.01]
Intermediate diversity (3–5 food groups)	1.05 [0.95; 1.17]	1.10 [0.99; 1.23]	1.21 [0.97; 1.51]	1.02 [0.91; 1.15]	1.15 [1.03; 1.28]
High diversity (6–8 food groups)	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]
Allergenic foods ^a at 8 months					
All 4 allergenic foods introduced	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]
1 allergenic food not introduced	1.13 [0.95; 1.34]	0.93 [0.78; 1.12]	1.42 [0.92; 2.20]	1.17 [0.96; 1.42]	1.05 [0.88; 1.25]
At least 2 allergenic foods not introduced	1.18 [0.99; 1.40]	0.99 [0.82; 1.18]	2.03 [1.32; 3.13]	1.13 [0.93; 1.38]	1.16 [0.97; 1.39]
Allergenic foods ^a at 10 months					
All 4 allergenic foods introduced	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]
1 allergenic food not introduced	1.17 [1.04; 1.31]	1.04 [0.92; 1.17]	1.29 [0.99; 1.67]	1.25 [1.10; 1.42]	1.07 [0.95; 1.21]
At least 2 allergenic foods not introduced	1.20 [1.003; 1.44]	1.11 [0.92; 1.34]	2.46 [1.77; 3.42]	1.13 [0.92; 1.38]	1.20 [1.002; 1.44]

Note: Values are OR [95% CI] from logistic regressions adjusted for maternal characteristics: age at delivery (18–24, 25–29, 30–34, ≥35 years), education level (up to lower secondary school, upper secondary school, intermediate, 3-year university degree, at least 5-year university degree), employment status (employed, unemployed, not in the labor force), migration (majority population, descendants of migrants, born in a foreign country under a foreign citizenship), smoking status during pregnancy (never smoker, smoker only before pregnancy, smoker in early pregnancy, smoker throughout pregnancy), region of residence (Paris region, North, East, Paris Basin–East, Paris Basin–West, West, Southwest, Southeast, Mediterranean), urban/rural area; household characteristics: monthly family income per consumption unit (≤€750; €751–1111; €1112–1500; €1501–1944; €1945–2500; >€2500), number of older children at home (no sibling, one sibling, at least two siblings); infant characteristics: cesarean section delivery, infant's sex, birthweight, gestational age; predominant breastfeeding duration family history of allergy, and variables related to study design (recruitment wave, maternity unit size). One model was run separately for each complementary feeding variable and each allergic outcome.

^aFour allergenic foods were considered in the present study: dairy products, eggs, fish and wheat.

When compared with recommended CF initiation, late CF initiation (>6 months) was associated with a higher risk of food allergy up to 5.5 years (Table 2). Nevertheless, this association was attenuated

and nonsignificant after the exclusion of early cases (occurring between the 2-month interview and the 1- or 2-year interview; Table 3). Early CF initiation was not related to the risk of food allergy.

TABLE 3 Adjusted associations between complementary feeding characteristics and allergic diseases up to 5.5 years, after exclusion of early allergic cases (any allergy occurring between the 2-month follow-up and the 1- or 2- year follow-up).

	Any allergy	Eczema	Food allergy	Asthma	Rhinoconjunctivitis
N	4359	4217	4208	4359	4011
Age at complementary feeding introduction					
<4 months	1.03 [0.81; 1.31]	1.08 [0.82; 1.41]	0.59 [0.21; 1.65]	1.11 [0.78; 1.58]	1.10 [0.84; 1.43]
4–6 months	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]
>6 months	0.98 [0.82; 1.17]	0.92 [0.75; 1.12]	1.10 [0.62; 1.96]	0.92 [0.71; 1.21]	1.07 [0.88; 1.30]
Diversity score at 8 months					
Low diversity (0–2 food groups)	1.11 [0.90; 1.37]	1.01 [0.79; 1.29]	1.17 [0.56; 2.46]	1.37 [0.998; 1.87]	1.03 [0.82; 1.29]
Intermediate diversity (3–5 food groups)	1.12 [0.95; 1.33]	1.16 [0.95; 1.40]	1.12 [0.60; 2.09]	1.13 [0.87; 1.47]	1.01 [0.84; 1.21]
High diversity (6–8 food groups)	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]
Diversity score at 10 months					
Low diversity (0–2 food groups)	0.93 [0.63; 1.37]	0.88 [0.55; 1.41]	1.57 [0.54; 4.61]	0.92 [0.51; 1.69]	1.47 [0.96; 2.24]
Intermediate diversity (3–5 food groups)	1.02 [0.90; 1.16]	1.07 [0.93; 1.24]	1.01 [0.66; 1.56]	0.99 [0.82; 1.19]	1.07 [0.93; 1.22]
High diversity (6–8 food groups)	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]
Allergenic foods^a at 8 months					
All four allergenic foods introduced	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]
One allergenic food not introduced	1.10 [0.90; 1.35]	0.98 [0.78; 1.23]	1.86 [0.71; 4.82]	1.18 [0.86; 1.62]	0.95 [0.76; 1.19]
At least two allergenic foods not introduced	1.16 [0.94; 1.43]	0.99 [0.78; 1.25]	2.41 [0.93; 6.21]	1.09 [0.79; 1.50]	1.10 [0.88; 1.38]
Allergenic foods^a at 10 months					
All four allergenic foods introduced	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]	1 [Ref]
One allergenic food not introduced	1.12 [0.98; 1.28]	1.13 [0.97; 1.32]	1.55 [0.92; 2.60]	1.10 [0.90; 1.35]	0.98 [0.85; 1.14]
At least 2 allergenic foods not introduced	1.02 [0.82; 1.28]	1.06 [0.82; 1.36]	2.32 [1.18; 4.59]	0.99 [0.71; 1.38]	1.06 [0.84; 1.35]

Note: Values are OR [95%CI] from logistic regressions adjusted for maternal characteristics (age at delivery, education level, employment status, migration, smoking status during pregnancy, region of residence, urban/rural area), household characteristics (income per consumption unit, number of older children at home), infant characteristics (cesarean section delivery, infant's sex, birthweight, gestational age), predominant breastfeeding duration, family history of allergy, and variables related to study design (recruitment wave, maternity unit size). One model was run separately for each complementary feeding variable and each allergic outcome.

^aFour allergenic foods were considered in the present study: dairy products, eggs, fish, and wheat.

Moreover, the age at CF initiation was not associated with the risk of asthma and rhinoconjunctivitis.

Similar results were observed in unadjusted or weighted analyses (Tables S1 and S2).

3.3 | Food diversity during complementary feeding

In relation to a high food-diversity score, a low food-diversity score (0–2 food groups consumed) at 8 months was associated with a

higher risk of asthma and food allergy (Table 2), and a low or intermediate food-diversity score at 10 months was associated with a greater risk of rhinoconjunctivitis. These associations were attenuated and nonsignificant after the exclusion of early cases (occurring between the 2-month interview and the 1- or 2-year interview; Table 3). In addition, the food-diversity scores were not related to the risk of any allergy or eczema.

Similar results were observed in unadjusted or weighted analyses (Tables S1 and S2).

3.4 | Delayed introduction of major allergenic foods (out of dairy products, hen's eggs, wheat, and fish)

When compared to children that were introduced to all four allergenic foods, children with at least two allergenic foods not introduced at 8 months old were at a higher risk of food allergy (Table 2). Moreover, children with at least two allergenic foods not introduced at 10 months of age were at a higher risk of food allergy and rhinoconjunctivitis. The association between the number of allergenic foods not introduced at 10 months and food allergy remained significant after the exclusion of early cases (occurring between the 2-month interview and the 2-year interview; Table 3).

Similar results were observed in unadjusted or weighted analyses (Tables S1 and S2).

4 | DISCUSSION

In this nationwide birth cohort, food diversity increased regularly during the CF period, but at 10 months old, at least one major allergenic food out of dairy products, egg, fish, and wheat had still not been introduced to more than half of the children (most frequently egg). Late CF introduction, low food-diversity by the age of 10 months and the delayed introduction of major allergenic foods appeared to be associated with a higher risk of allergic diseases, especially food allergy and rhinoconjunctivitis. However, only the association between the delayed introduction of allergenic foods and the risk of food allergy remained significant after the exclusion of early allergic cases (occurring between the 2-month interview and the 1-year interview or, for the food allergy only, the 2-year interview).

In this study, CF introduction after 6 months was associated with a higher risk of food allergy. Hicke-Roberts et al.^{11,12} uncovered similar results with regard to a higher risk of food allergy among infants that were introduced to CF after 7 months, but most studies found no significant association between late CF initiation and food allergy.^{4,5} As the association was attenuated and nonsignificant after the exclusion of early cases, we cannot rule out the possibility that it was due to a reverse causation bias where parents of children with early allergic diseases delayed the introduction of complementary food.

A previous study reported a positive association between the food-diversity score at 6 months old and the risk of food allergy at 1 year.¹² During our research, even if late CF was shown to be associated with a higher risk of food allergy, we did not find a clear association between food diversity and the risk of food allergy. As was the case in our study, previous studies underlined an association between a low food-diversity score in infancy and a higher risk of asthma^{17,18} or rhinoconjunctivitis.¹⁸ However, the associations in our study were not uniform across the ages, which therefore limits the generalizability of our findings. Moreover, in the present paper, the diversity score was based on the food groups consumed during a given month, whereas previous papers have mostly examined the food groups introduced during a set period.^{12,17,18} The frequency of consumption might play a role, but research exploring the association between the frequency of consuming several food items and allergic diseases is very scarce.²⁹ It would consequently be of interest to conduct further studies exploring the potential influence of both the frequency of consumption of each food group and the impact of food diversity on allergic diseases.

Regarding the delayed introduction of allergenic foods, several studies reported that it was associated with a higher risk of food allergy, but most of them focused on a given allergenic food and the risk of allergy in relation to this allergenic food.^{20,30,31} In this study, we showed that the delayed (>10 months) introduction of at least two major allergenic foods (out of dairy products, wheat, egg, and fish) is associated with a higher risk of food allergy or rhinoconjunctivitis. Furthermore, the association with food allergy persisted after the exclusion of early allergic cases (any allergy occurring between the 2-month follow-up and the 2-year follow-up), indicating that this association was not explained by reverse causation. Egg was the most frequent allergenic food not introduced at 10 months of age; nevertheless, it is worth noting here that we were unable to take the introduction of baked eggs (in processed foods like cakes) into account because of the questionnaire's design. This might lead to an overestimate in terms of how delayed the introduction of egg was, but our results suggest that the introduction of a whole egg into infant diets is an important part of food allergy prevention. Indeed, earlier research demonstrate that a weekly protein intake of more than 2 g is important to prevent allergies,^{29,32} so the consumption of lower levels of egg protein through processed foods might not be sufficient to prevent egg allergy.³³ Although several studies found a higher risk of atopic dermatitis or asthma among children with late or scarce amounts of fish being introduced into their diet,^{19,29,32} the delayed introduction of food allergens (including fish) was not associated with the risk of eczema in our study. Meanwhile, some research found an association between the delayed introduction of fish after the age of 11 months and an increased risk of asthma and rhinoconjunctivitis,^{17,34} yet most studies found no link between the introduction of allergenic foods and asthma or rhinoconjunctivitis.^{5,6,9,13,17,19} When the ELFE cohort was launched, French nutritional guidelines recommended delaying the introduction of allergenic food to children with a family history of allergy, and European guidelines were unclear whether parents should delay their introduction or not.³⁵

Our findings support the updated guidelines, indicating that the introduction of food allergens should not be delayed and should occur “when CF is initiated any time after 4 months”⁸ and “as early as the CF is initiated that is to say between 4 and 6 months” (updated French guidelines from the Public Health France website, September 2021).³⁶ Recent studies underline the need for public health communications to disseminate and explain this change in the guidelines to parents and professionals who work with children.^{37,38}

The very large sample and the collection of detailed socio-demographic and economic data ensure the presence of more statistical power and a favorable level of control over potential confounders. As in most observational studies, the selection/attrition bias was quite important, but the weighted analyses suggest that this bias had a limited impact on our findings. Indeed, the prospective design allowed for limited memory bias with regard to both exposure and outcome variables. However, the health data were reported by parents and not validated by medical records, which could lead to a potential measurement bias, even if the items used were derived from international questionnaires.³⁹ In particular, parental reporting could lead to an overestimation of allergic diseases. As children were considered to have food allergies when medical advice was given to avoid one specific food, due to a food allergy, then it was not possible to distinguish non-IgE-mediated food allergies from IgE-mediated food allergies. As allergy outcomes were not reported between 2 months and 1 year or even 2 years for food allergies, allergies reported at the 1 or 2-year interview could have occurred at any time between 2 months and 1 or 2-year of age. It was subsequently not possible to distinguish those occurring before CF introduction (prevalent cases) from those occurring after CF introduction (real incident cases). By comparing our main analysis and our sensitivity analysis (excluding all cases reported at 1 or 2 years for food allergies), we concluded that for some associations (those not significant in the sensitivity analyses), the reverse causation could not be excluded, whereas the association between the delayed introduction of allergenic foods and food allergy is unlikely to be due to this reverse causation.

In this nationwide cohort study, some aspects of CF practices were shown to be related to the risk of allergic diseases up to 5.5 years; this was mainly the case in relation to food allergy. In fact, delayed introduction (after 8 or 10 months) of major allergenic foods (dairy products, wheat, fish, and egg) was linked to a higher risk of food allergies up to 5.5 years (this more than doubled if there was a delayed introduction of at least two of them), even after excluding cases that occurred during the first 2 years of life. Our results support the updated guidelines for CF practices on early CF practices for food allergy prevention.

AUTHOR CONTRIBUTIONS

T. Adam: conceptualization (lead); formal analysis (lead); writing – original draft (lead). A. Divaret-Chauveau: conceptualization (lead); methodology (lead); supervision (lead); writing—review and editing (lead). C. Roduit: methodology (equal); writing—review and editing (equal). K. Adel-Patient: methodology (equal); resources (equal); writing—review

and editing (equal). A. Deschildre: writing—review and editing (supporting). C. Raheison: resources (equal); writing—review and editing (supporting). M.-A. Charles: resources (equal); writing—review and editing (supporting). S. Nicklaus: methodology (equal); resources (equal); writing—review and editing (equal). B. de Lauzon-Guillain: conceptualization (lead); resources (lead); methodology (lead); supervision (lead); formal analysis (lead); writing—review and editing (lead).

ACKNOWLEDGMENTS

We thank the scientific coordinators (B. Geay, H. Léridon, C. Bois, J. L. Lanoé, X. Thierry, C. Zaros), IT and data managers, statisticians (M. Cheminat, C. Ricourt, A. Candea, S. de Visme), administrative and family communication staff, and study technicians (C. Guevel, M. Zoubiri, L. G. L. Gravier, I. Milan, R. Popa) from the ELFE coordination team as well as the families that gave their time for the study. We would also like to thank researchers at the French Institute for Demographic Studies (INED) for the socio-demographic variables created as part of the ANR-funded Veniromond project.

FUNDING INFORMATION

This study was funded by a grant from the French National Research Agency (ANR-19-CE36-0008). The ELFE survey is a joint project between the French Institute for Demographic Studies (INED) and the National Institute of Health and Medical Research (INSERM), in partnership with the French blood transfusion service (Etablissement Français du Sang, EFS), Public Health France, the National Institute for Statistics and Economic Studies (INSEE), the Direction Générale de la Santé (DGS, part of the Ministry of Health and Social Affairs), the Direction Générale de la Prévention des Risques (DGPR, Ministry for the Environment), the Direction de la Recherche, des Études, de l'Évaluation et des Statistiques (DREES, Ministry of Health and Social Affairs), the Département des Études, de la Prospective et des Statistiques (DEPS, Ministry of Culture), and the Caisse Nationale des Allocations Familiales (CNAF), with the support of the Ministry of Higher Education and Research and the Institut National de la Jeunesse et de l'Éducation populaire (INJEP). Via the RECONAI platform, the study received a government grant managed by the National Research Agency under the “Investissements d'avenir” program (ANR-11-EQPX-0038, ANR-19-COHO-0001). The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

CONFLICT OF INTEREST STATEMENT

T. Adam reports support for attending meetings from Stallergens, ALK, and Sanofi. A. Divaret-Chauveau reports receiving support for this manuscript from the French National Research Agency (ANR-19-CE36-0008) and the University of Lorraine, grants for other studies from Don du Souffle, Fondation du Souffle, Novartis, and the Association Régionale d'Aide aux Insuffisants Respiratoires de Lorraine (ARAIRLOR). The author also had a contract with the French public agency ANSES as an expert in allergies and pediatrics for the Human Nutrition Committee, received consulting fees from Stallergens, ALK,

and Aimmune Therapeutics, received payment as a speaker from Aimmune Therapeutics and Novartis, and received support for attending meetings from Aimmune Therapeutics, Novartis, Mead Johnson, and Nutricia. C. Roduit received consulting fees from Aimmune Therapeutics, payment as a speaker and support for attending meetings from ALK. K. Adel-Patient's lab received grants for lectures from Stallergens, and research grants from Mead Johnson, and she is an expert for the Human Nutrition Committee and Biotechnology Group in the French public agency ANSES. A. Deschildre received speaker/consulting fees from Nestlé Health Science, Sanofi, Regeneron Pharmaceuticals Inc, ALK, Novartis, GSK, Stallergenes Greer, Astra Zeneca, Aimmune Therapeutics, and DBV Technologies, and support for attending meetings from MEDA, Nutricia, Astra Zeneca, Sanofi, and Aimmune Therapeutics. C. Raheison has no conflict of interest to declare. M.-A. Charles received research grants from the Centre National Interprofessionnel de l'Économie Laitière (CNIEL) that were not related to this manuscript. S. Nicklaus has no conflict of interest to declare. B. de Lauzon-Guillain received support for this manuscript from the French National Research Agency and, outside the present project, grants from the Fondation pour la Recherche Médicale. She is also an expert for the Human Nutrition Committee from the French public agency ANSES.

DATA AVAILABILITY STATEMENT

The data underlying the findings in this study cannot be made freely available for ethical and legal reasons because they include a substantial number of variables that, together, could be used to identify the participants based on a few key characteristics and could then be used to access other personal data. Therefore, the French ethics authority strictly forbids making these data freely available. However, they can be obtained upon request from the ELFE principal investigator. Readers may contact marie-aline.charles@inserm.fr to request the data.

ORCID

T. Adam  <https://orcid.org/0000-0003-4481-7787>
 A. Divaret-Chauveau  <https://orcid.org/0000-0002-2492-9864>
 C. Roduit  <https://orcid.org/0000-0002-5988-0570>
 K. Adel-Patient  <https://orcid.org/0000-0002-2242-0626>
 M.-A. Charles  <https://orcid.org/0000-0003-4025-4390>
 S. Nicklaus  <https://orcid.org/0000-0002-0659-2823>
 B. de Lauzon-Guillain  <https://orcid.org/0000-0001-5887-8842>

REFERENCES

- Lack G. Update on risk factors for food allergy. *J Allergy Clin Immunol.* 2012;129(5):1187-1197. doi:10.1016/j.jaci.2012.02.036
- Weidinger S, Novak N. Atopic dermatitis. *The Lancet.* 2016;387(10023):1109-1122. doi:10.1016/S0140-6736(15)00149-X
- Testa D, Bari DI, Nunziata M, et al. Allergic rhinitis and asthma assessment of risk factors in pediatric patients: a systematic review. *Int J Pediatr Otorhinolaryngol.* 2020;129:109759. doi:10.1016/j.ijporl.2019.109759
- Luccioli S, Zhang Y, Verrill L, Ramos-Valle M, Kwegyir-Afful E. Infant feeding practices and reported food allergies at 6 years of age. *Pediatrics.* 2014;134(Suppl 1):S21-S28. doi:10.1542/peds.2014-0646E
- Zutavern A, Brockow I, Schaaf B, et al. Timing of solid food introduction in relation to atopic dermatitis and atopic sensitization: results from a prospective birth cohort study. *Pediatrics.* 2006;117(2):401-411. doi:10.1542/peds.2004-2521
- Zutavern A, Brockow I, Schaaf B, et al. Timing of solid food introduction in relation to eczema, asthma, allergic rhinitis, and food and inhalant sensitization at the age of 6 years: results from the prospective birth cohort study LISA. *Pediatrics.* 2008;121(1):e44-e52. doi:10.1542/peds.2006-3553
- Nwaru BI, Takkinen HM, Niemelä O, et al. Timing of infant feeding in relation to childhood asthma and allergic diseases. *J Allergy Clin Immunol.* 2013;131(1):78-86. doi:10.1016/j.jaci.2012.10.028
- Fewtrell M, Bronsky J, Campoy C, et al. Complementary feeding: a position paper by the European Society for Paediatric Gastroenterology, hepatology, and nutrition (ESPGHAN) committee on nutrition. *J Pediatr Gastroenterol Nutr.* 2017;64(1):119-132. doi:10.1097/MPG.0000000000001454
- Nwaru BI, Takkinen HM, Niemelä O, et al. Introduction of complementary foods in infancy and atopic sensitization at the age of 5 years: timing and food diversity in a Finnish birth cohort. *Allergy.* 2013;68(4):507-516. doi:10.1111/all.12118
- Venter C, Maslin K, Dean T, Arshad SH. Does concurrent breastfeeding alongside the introduction of solid food prevent the development of food allergy? *J Nutr Sci.* 2016;5:e40. doi:10.1017/jns.2016.31
- Hicke-Roberts A, Wennergren G, Hesselmar B. Late introduction of solids into infants' diets may increase the risk of food allergy development. *BMC Pediatr.* 2020;20(1):273. doi:10.1186/s12887-020-02158-x
- Venter C, Maslin K, Holloway JW, et al. Different measures of diet diversity during infancy and the association with childhood food allergy in a UK birth cohort study. *J Allergy Clin Immunol Pract.* 2020;8(6):2017-2026. doi:10.1016/j.jaip.2020.01.029
- Filipiak B, Zutavern A, Koletzko S, et al. Solid food introduction in relation to eczema: results from a four-year prospective birth cohort study. *J Pediatr.* 2007;151(4):352-358. doi:10.1016/j.jpeds.2007.05.018
- Joseph CLM, Ownby DR, Havstad SL, et al. Early complementary feeding and risk of food sensitization in a birth cohort. *J Allergy Clin Immunol.* 2011;127(5):1203-1210.e5. doi:10.1016/j.jaci.2011.02.018
- Du Toit G, Roberts G, Sayre PH, et al. Randomized trial of peanut consumption in infants at risk for peanut allergy. *N Engl J Med.* 2015;372(9):803-813. doi:10.1056/NEJMoa1414850
- Ierodiakonou D, Garcia-Larsen V, Logan A, et al. Timing of allergenic food introduction to the infant diet and risk of allergic or autoimmune disease: a systematic review and meta-analysis. *Jama.* 2016;316(11):1181-1192. doi:10.1001/jama.2016.12623
- Roduit C, Frei R, Depner M, et al. Increased food diversity in the first year of life is inversely associated with allergic diseases. *J Allergy Clin Immunol.* 2014;133(4):1056-1064.e7. doi:10.1016/j.jaci.2013.12.1044
- Nwaru BI, Takkinen HM, Kaila M, et al. Food diversity in infancy and the risk of childhood asthma and allergies. *J Allergy Clin Immunol.* 2014;133(4):1084-1091. doi:10.1016/j.jaci.2013.12.1069
- Roduit C, Frei R, Loss G, et al. Development of atopic dermatitis according to age of onset and association with early-life exposures. *J Allergy Clin Immunol.* 2012;130(1):130-136.e5. doi:10.1016/j.jaci.2012.02.043
- Perkin MR, Logan K, Bahnson HT, et al. Efficacy of the enquiring about tolerance (EAT) study among infants at high risk of developing food allergy. *J Allergy Clin Immunol.* 2019;144(6):1606-1614.e2. doi:10.1016/j.jaci.2019.06.045

21. Skjerven HO, Lie A, Vettukattil R, et al. Early food intervention and skin emollients to prevent food allergy in young children (PreventADALL): a factorial, multicentre, cluster-randomised trial. *Lancet*. 2022;399(10344):2398-2411. doi:10.1016/S0140-6736(22)00687-0
22. Charles MA, Thierry X, Lanoe JL, et al. Cohort profile: the French national cohort of children (ELFE): birth to 5 years. *Int J Epidemiol*. 2020;49(2):368-369j. doi:10.1093/ije/dyz227
23. Wagner S, Kersuzan C, Gojard S, et al. Breastfeeding initiation and duration in France: the importance of intergenerational and previous maternal breastfeeding experiences – results from the nationwide ELFE study. *Midwifery*. 2019;69:67-75. doi:10.1016/j.midw.2018.10.020
24. Accès aux données et questionnaires. Ined - Institut national d'études démographiques. Accessed August 3, 2022. [https://www.elfe-france.fr/fr/cote-recherche/acces-aux-donnees-et-questions/](https://www.elfe-france.fr/fr/cote-recherche/acces-aux-donnees-et-questions)
25. Bournez M, Ksiazek E, Wagner S, et al. Factors associated with the introduction of complementary feeding in the French ELFE cohort study. *Matern Child Nutr*. 2018;14(2):e12536. doi:10.1111/mcn.12536
26. Payet D, Adjibade M, Baudry J, et al. Organic food consumption during the complementary feeding period and respiratory or allergic diseases up to age 5.5 years in the ELFE cohort. *Front Nutr*. 2021;8:791430. doi:10.3389/fnut.2021.791430
27. Shrier I, Platt RW. Reducing bias through directed acyclic graphs. *BMC Med Res Methodol*. 2008;8:70. doi:10.1186/1471-2288-8-70
28. Blondel B, Lelong N, Kermarrec M, Goffinet F. National Coordination Group of the National Perinatal Surveys. Trends in perinatal health in France from 1995 to 2010. Results from the French National Perinatal Surveys. *J Gynecol Obstet Biol Reprod (Paris)*. 2012;41(4):e1-e15. doi:10.1016/j.jgyn.2012.04.014
29. Oien T, Storrø O, Johnsen R. Do early intake of fish and fish oil protect against eczema and doctor-diagnosed asthma at 2 years of age? A cohort study. *J Epidemiol Community Health*. 2010;64(2):124-129. doi:10.1136/jech.2008.084921
30. Natsume O, Kabashima S, Nakazato J, et al. Two-step egg introduction for prevention of egg allergy in high-risk infants with eczema (PETIT): a randomised, double-blind, placebo-controlled trial. *Lancet*. 2017;389(10066):276-286. doi:10.1016/S0140-6736(16)31418-0
31. Poole JA, Barriga K, Leung DYM, et al. Timing of initial exposure to cereal grains and the risk of wheat allergy. *Pediatrics*. 2006;117(6):2175-2182. doi:10.1542/peds.2005-1803
32. Kull I, Bergström A, Lilja G, Pershagen G, Wickman M. Fish consumption during the first year of life and development of allergic diseases during childhood. *Allergy*. 2006;61(8):1009-1015. doi:10.1111/j.1398-9995.2006.01115.x
33. Perkin MR, Logan K, Tseng A, et al. Randomized trial of introduction of allergenic foods in breast-fed infants. *N Engl J Med*. 2016;374(18):1733-1743. doi:10.1056/NEJMoa1514210
34. Klingberg S, Brekke HK, Ludvigsson J. Introduction of fish and other foods during infancy and risk of asthma in the all babies In Southeast Sweden cohort study. *Eur J Pediatr*. 2019;178(3):395-402. doi:10.1007/s00431-018-03312-5
35. Agostoni C, Decsi T, Fewtrell M, et al. ESPGHAN committee on nutrition: complementary feeding: a commentary by the ESPGHAN committee on nutrition. *J Pediatr Gastroenterol Nutr*. 2008;46(1):99-110. doi:10.1097/01.mpg.0000304464.60788.bd
36. Santé publique France accompagne les parents pour prendre en main les nouvelles recommandations sur la diversification alimentaire des tout-petits. Accessed August 1, 2022. <https://www.sante-publiquefrance.fr/presse/2021/sante-publique-france-accompagne-les-parents-pour-prendre-en-main-les-nouvelles-recommandations-sur-la-diversification-alimentaire-des-tout-petits>
37. De Rosso S, Schwartz C, Ducrot P, Nicklaus S. The perceptions and needs of French parents and pediatricians concerning information on complementary feeding. *Nutrients*. 2021;13(7):2142. doi:10.3390/nu13072142
38. DeRosso S, Ducrot P, Chabanet C, Nicklaus S, Schwartz C. Increasing parental knowledge about child feeding: evaluation of the effect of public health policy communication media in France. *Front Public Health*. 2022;10:782620. doi:10.3389/fpubh.2022.782620
39. Asher MI, Montefort S, Björkstén B, et al. Worldwide time trends in the prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and eczema in childhood: ISAAC phases one and three repeat multicountry cross-sectional surveys. *Lancet Lond Engl*. 2006;368(9537):733-743. doi:10.1016/S0140-6736(06)69283-0

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Adam T, Divaret-Chauveau A, Roduit C, et al. Complementary feeding practices are related to the risk of food allergy in the ELFE cohort. *Allergy*. 2023;00:1-11. doi:10.1111/all.15828