

Enrichment strategy of chemiolithotrophic biofilm for nitrogen fixation in MES

Axel Rous, Eric Trably, Elie Le-Quemener, Nicolas Bernet

▶ To cite this version:

Axel Rous, Eric Trably, Elie Le-Quemener, Nicolas Bernet. Enrichment strategy of chemiolithotrophic biofilm for nitrogen fixation in MES. ISMET8 – 2022 Global Conference, Sep 2022, Chania, Greece. hal-04176863

HAL Id: hal-04176863 https://hal.inrae.fr/hal-04176863

Submitted on 3 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Enrichment strategy of chemiolithotrophic biofilm for nitrogen fixation in MES

Axel Rous, EricTrably, Elie Le-Quemener and Nicolas Bernet

INRAE, Univ Montpellier, LBE, 102 avenue des Etangs, 11100, Narbonne, France

Corresponding author email: axel.rous@inrae.fr

ABSTRACT

Nitrogen fixation (from N_2 to NH_4) is the first step in the transformation of atmospheric nitrogen into nitrogen that can be assimilated by living organisms. This reaction is catalysed by so-called diazotrophic microorganisms that reduce atmospheric nitrogen into ammonia to fuel their growth. Biomass production by N_2 and CO_2 fixation via electrostimulation in a microbial electrochemical system has been shown recently (Rago et al. 2019). We propose to investigate enrichment strategies allowing the selection of a cathodic biofilm capable of growing from the present gases and a mineral energy supply.

Two enrichment methods were tested. The first one focused on enriching nitrogen-fixing bacteria in C-organic supplied medium. Then a MES was use to enrich chemolithoautotrophic bacteria capable of using CO₂ and electrons supplied at the cathode. The second strategy focused on the enrichment of chemolithoautotrophic bacteria before enriching nitrogen-fixing bacteria without using MES and with H₂ as energy source.

We used dual-chamber reactors separated by AEM with a three electrodes set-up. The carbon felt working electrode was polarized at -0.7 V vs SHE (pH 6,5-7) when OCV were used as control.

The rate of nitrogen fixation measured after 200 days in the MES without organic-C nor assimilable N was higher than in the second enrichment method ($60.8 \pm 11.5 \, \mu mol_{C2H4}/L_{gaz}.d$) versus 12.7 $\pm 2.6 \, \mu mol_{C2H4}/L_{gaz}.d$). About 7 times more 16S rRNA copies/mL were observed in the medium of the MES than in the medium of the second method with H₂-supply. Enrichment led to a biofilm accumulation during 87 days in MES with N₂ and CO₂ as sole nitrogen and carbon sources respectively. Acetate was also measured up to 0.75 g/L after 15 days suggesting that bacteria are able to produce acetate with CO₂ as carbon source in our first enrichment strategy. 16S rRNA sequencing showed the presence of *Hydrogenophilaceae*, *Chitinophagaceae* and *Rhodospirillaceae* (*Azospirillum*) in MES, known as nitrogen-fixing bacteria.

Finally, these results confirm that our first enrichment method had successfully enrich a microbial community able to fix N_2 and CO_2 in a cathodic biofilm.




Fig 1. Comparison of Acetylene Reduction Assay (nitrogen fixation rate) between enriched biofilm in MES and H₂ enriched community

Fig 2. Bacterial community of the two enrichments for chemiolithoautotrophics nitrogen fixing bacteria

Acknowledgements: The research was supported by the CATHOMIX project financed by National Research Agency (n° ANR-19-CE43-0013) in France .

References:

Rago, Laura et al. 2019. « Bioelectrochemical Nitrogen Fixation (e-BNF): Electro-Stimulation of Enriched Biofilm Communities Drives Autotrophic Nitrogen and Carbon Fixation ». *Bioelectrochemistry* 125: 105-15.

