



HAL
open science

A dynamic mechanistic model of microbial fermentation and methane production in the cow rumen

Rafael Munoz Tamayo, Seppo Ahvenjärvi, Ali Reza Bayat, Ilma Tapio

► **To cite this version:**

Rafael Munoz Tamayo, Seppo Ahvenjärvi, Ali Reza Bayat, Ilma Tapio. A dynamic mechanistic model of microbial fermentation and methane production in the cow rumen. ADSA Annual Meeting, ADSA, Jun 2023, Ottawa, Canada. hal-04181482

HAL Id: hal-04181482

<https://hal.inrae.fr/hal-04181482v1>

Submitted on 16 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A dynamic mechanistic model of microbial fermentation and methane production in the cow rumen

R. Muñoz-Tamayo ^a, S. Ahvenjärvi ^b, A. R. Bayat ^b, I. Tapio ^c

^a Université Paris-Saclay, INRAE, AgroParisTech, UMR Modélisation Systémique Appliquée aux Ruminants, 91120, Palaiseau, France

^b Animal Nutrition, Production Systems, Natural Resources Institute Finland (Luke), Jokioinen, Finland

^c Genomics and Breeding, Production Systems, Natural Resources Institute Finland (Luke), Jokioinen, Finland

Dynamic mathematical models have been developed to enhance understanding of rumen function. These models include Molly¹, the Dijkstra model², Karoline³ and their extensions. Model evaluations showed that there is still room to enhance their accuracy for predicting volatile fatty acids concentration and methane production. Previously, Muñoz-Tamayo et al. (2016) developed an alternative model of rumen fermentation under *in vitro* conditions aimed at better representing the rumen microbiota and hydrogen dynamics. In the present work, we extended the previous model to account for *in vivo* conditions. We performed an experiment with four Nordic-Red dairy cows equipped with rumen fistulas to provide dynamic data to support model construction. The experiment determined animal daily patterns of feed intake, rumen fermentation and enteric methane production. Feed intake and methane emissions were measured in respiration chambers during two days after an adaptation period. Rumen liquid was collected every three hours for determination of volatile fatty acids concentration. In the model, the rumen microbiota is represented by three functional groups namely sugar-, amino acids- and hydrogen-utilisers. The feed is expressed in three pools as neutral detergent fiber, non-structural carbohydrates and proteins. The model consists of 18 compartments described by ordinary differential equations. The model performance was satisfactory as evaluated by the coefficient of variation of the root mean squared error, which was 9% for acetate, 11% for butyrate, 13% for propionate and 21% for methane. Our model has the potential to be used as virtual platform to simulate the dynamics of rumen fermentation and methane production.

Acknowledgements. The authors acknowledge funding from the MASTER project, an Innovation Action funded by the European Union's Horizon 2020 research and innovation programme under grant agreement No 818368.

References

- 1 Baldwin, R. L. *et al. J. Dairy Res.* 54, 107–131 (1987)
- 2 Dijkstra, J. *et al. J. Nutr.* 122, 2239–2256 (1992)
- 3 Danfær, A. *et al.* (CABI Publishing, 2006)
- 4 Muñoz-Tamayo, R. *et al. Anim. Feed Sci. Technol.* 220, 1–21 (2016)