



HAL
open science

Chemical Profile, Antioxidant and Anti-Inflammatory Potency of Extracts of *Vitex madiensis* Oliv. and *Crossopteryx febrifuga* (Afzel ex G. Don)

Ghislaine Bounvou-Tsona, Maël Gainche, Caroline Decombat, Isabelle Ripoche, Kevin Bikindou, Laetitia Delort, Florence Caldefie-Chézet, Aubin Loumouamou, Pierre Chalard

► To cite this version:

Ghislaine Bounvou-Tsona, Maël Gainche, Caroline Decombat, Isabelle Ripoche, Kevin Bikindou, et al.. Chemical Profile, Antioxidant and Anti-Inflammatory Potency of Extracts of *Vitex madiensis* Oliv. and *Crossopteryx febrifuga* (Afzel ex G. Don). *Plants*, 2023, 12 (2), pp.386. 10.3390/plants12020386 . hal-04181723

HAL Id: hal-04181723

<https://hal.inrae.fr/hal-04181723>

Submitted on 16 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Article

Chemical Profile, Antioxidant and Anti-Inflammatory Potency of Extracts of *Vitex madiensis* Oliv. and *Crossopteryx febrifuga* (Afzel ex G. Don)

Ghislaine Boungou-Tsona ^{1,2,3}, Maël Gainche ³ , Caroline Decombat ⁴ , Isabelle Ripoche ^{3,*}, Kevin Bikindou ^{1,2}, Laetitia Delort ⁴ , Florence Caldefie-Chézet ⁴ , Aubin Loumouamou ^{1,2} and Pierre Chalard ^{3,*} 

- ¹ Equipe Pluridisciplinaire de Recherche en Alimentation et Nutrition (EPRAN), Faculté des Sciences et Techniques, Université Marien Ngouabi, Brazzaville BP 389, Congo
- ² Département des Sciences Chimiques, Institut National de Recherche en Sciences Exactes et Naturelles (IRSEN), UR Chimie des Substances Naturelles, Cité Scientifique de Brazzaville, Brazzaville BP 2400, Congo
- ³ Institut de Chimie de Clermont-Ferrand, Université Clermont Auvergne, Clermont Auvergne INP, Centre National de la Recherche Scientifique, F-63000 Clermont-Ferrand, France
- ⁴ Unité de Nutrition Humaine, l'Alimentation et l'Environnement, Institut National de Recherche pour l'Agriculture, Université Clermont-Auvergne, CRNH Auvergne, F-63000 Clermont-Ferrand, France
- * Correspondence: isabelle.ripoche@sigma-clermont.fr (I.R.); pierre.chalard@sigma-clermont.fr (P.C.)

Abstract: *Vitex madiensis* Oliv. (Lamiaceae) and *Crossopteryx febrifuga* (Rubiaceae), two plants commonly used in traditional African medicines to treat malaria and pain, were studied either to determine their chemical profiles or to evaluate their antioxidant and anti-inflammatory activities. In this study, we investigated leaves, trunk bark, root bark and fruits methanolic extracts of both plants in order to find out which part of the plant is responsible for the activity. The analyses of the chemical profiles allowed us to confirm the presence of several ecdysteroids, especially 20-hydroxyecdysone in some parts of *V. madiensis* and to highlight the presence of organic acids and phenol derivatives in *C. febrifuga*. Among the four parts of the plants studied, only the fruits extract of *C. febrifuga* could present anti-inflammatory activity by decreasing ROS production. The leaves and trunk bark extracts of *V. madiensis* showed significant free radical scavenging activity compared to ascorbic acid, and the same extracts decrease ROS production significantly. The activity of these two extracts could be explained by the presence of ecdysteroids and flavonoids. The ROS production inhibition of *V. madiensis* is particularly interesting to investigate with further analyses.

Keywords: chemical profiles; African plants; antioxidant and anti-inflammatory activities



Citation: Boungou-Tsona, G.; Gainche, M.; Decombat, C.; Ripoche, I.; Bikindou, K.; Delort, L.; Caldefie-Chézet, F.; Loumouamou, A.; Chalard, P. Chemical Profile, Antioxidant and Anti-Inflammatory Potency of Extracts of *Vitex madiensis* Oliv. and *Crossopteryx febrifuga* (Afzel ex G. Don). *Plants* **2023**, *12*, 386. <https://doi.org/10.3390/plants12020386>

Academic Editors: Seok-Geun Lee and In Jin Ha

Received: 2 December 2022

Revised: 27 December 2022

Accepted: 9 January 2023

Published: 13 January 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Crossopteryx febrifuga (Rubiaceae) and *Vitex madiensis* Oliv. (Lamiaceae) are two species growing in tropical and subtropical regions throughout the world. These plants are widely present in the savannah areas of West, East and Central Africa [1–3]. In Congo-Brazzaville, these two species grow particularly in the area of the “Plateau des cataractes” (Pool department) in the southern part of the country. These two plants are commonly advised for the treatment of pain by traditional healers.

C. febrifuga is a shrub whose various organs have multiple uses in traditional African medicine. Indeed, this plant has been used for many years for the treatment of pain and malaria, and its effectiveness is widely acclaimed, particularly by the Hausa communities of Nigeria [4]. The fruits are used in Mali for treatment of respiratory tract infections, as well as an antitussive and as a febrifuge [5]. The bark is used to treat fever and diarrhea in Guinea [6]. The roots, leaves and trunk bark are used to relieve stomach aches, chest pain and as an analgesic [7]. In Congo-Brazzaville, the fruit is used to treat diarrhea and fever. In the same country, this plant is used to treat headaches, migraine, bacterial infections

and epilepsy [8]. In addition, several studies have demonstrated that methanolic extracts of different parts of the plant have antimalarial, antipyretic, analgesic, antioxidant and anti-inflammatory properties [1,4,9]. Despite these uses in traditional medicine and the evaluation of biological properties of crude extracts, the phytochemical profile of this plant and its relationship with therapeutic uses have not been extensively investigated. Two bisdesmoside saponins were isolated [10] from *C. febrifuga* roots and two alkaloids, crossopterin and crossoptin, have been isolated from bark extracts [11].

V. madiensis is a shrub that is about 5 m tall with a massive underground woody rootstock. The leaves, fruits, stem bark and roots have traditional medicinal uses in several African countries [12]. The plant is used for the treatment of headaches, toothaches, aches and pains and more generally as an analgesic. In Gabon and Angola, it is used to treat the symptoms of malaria [13–15]. Unlike other species of the genus *V. madiensis*, there are few phytochemical studies on *V. madiensis*. These studies mentioned the identification of phytoecdysteroids from the root bark [16], particularly 20-hydroxyecdysone and ajugasterone C. Phytoecdysteroids have important physiological effects on insects, particularly in the defense of plants against them [17]. In general, species of the *Vitex* genus contain a variety of potentially bioactive molecules, such as iridoids, flavonoids, diterpenoids and phytosteroids [18,19], thus explaining the origin of the observed biological properties.

The objective of this work was to carry out a comparative study of the chemical profiles of methanolic extracts of the different parts (leaves, fruits, trunk bark and root bark) of *C. febrifuga* and *V. madiensis* and to determine their antioxidant properties by evaluating their free radical scavenging activity and their anti-inflammatory activity with tests on reactive oxygen species (ROS) production by leukocytes.

2. Results

2.1. Chemical Profiles

A methanolic maceration of each part of plant was carried out to extract most of the secondary metabolites. By comparing the yields obtained (Table 1), we could conclude that the yields were quite similar for all parts of *V. madiensis* (between 3 and 6%). For *C. febrifuga*, the yields were quite high for the leaves, trunk bark and roots (between 10 and 19%), but the yield is lower for fruits (3%).

The extract of each part of plants was analyzed by HPLC in order to determine the best conditions to separate most of the compounds present in the extracts. For both plants, the analyses were performed using a method developed in our group to study complex matrices and for which a database of around one hundred compounds was established using commercially available standards.

Table 1. Ecdysteroids and other compounds identified in extracts of leaves, trunk bark, root bark and fruits of *Vitex madiensis*.

N°	Compound	Tr (min)	Formula	M-H (m/z)	M + (m/z)	MS ² (m/z)	Le	TB	RB	F	Reference
1	Quinic acid	3.96	C ₇ H ₁₂ O ₆	191.0553		191 /85/127/93/85	++	+	+++	+	Standard
2	Citric acid	6.91	C ₆ H ₈ O ₇	191.0189		191 /129/11/87/85	+	+	+	+	Standard
3	1-oxo-eucommiol	7.76	C ₉ H ₁₄ O ₅	201.0757		201 /109/139	++	+	+	+	[20]
4	5-Ethylidene-2-hydroxy-2-hydroxymethyl-3-methylhexanedioic acid	9.40	C ₁₀ H ₁₆ O ₆	231.0866		231 /213/169/125/157/187/143	+	-	-	+	[21]
5	Protocatechuic acid	11.49	C ₇ H ₆ O ₄	153.0179		109/ 153 /110	+	-	-	-	[22]
6	Vicenin-1	14.72	C ₂₆ H ₂₈ O ₁₄	563.1413		563 /353/383/297/473/443/297/325/503/545	+	-	-	-	[23]
7	Homoorientin	15.15	C ₂₁ H ₂₀ O ₁₁	447.0928		429/357/327	+	+	+	-	Standard
8	Orientin	15.91	C ₂₁ H ₂₀ O ₁₁	447.0928		369/357/327/299	+	+	-	-	Standard
9	Vitexin	17.90	C ₂₁ H ₂₀ O ₁₀	431.0981			+	+	-	+	Standard
10	20-Hydroxyecdysone	19.15	C ₂₇ H ₄₄ O ₇	525.3065 *	481.3152	445/371/165/427/125/69/ 481 [#]	+++	+	+++	+++	[24–26]
11	Luteolin-7-O-glucuronide	19.57	C ₂₁ H ₁₈ O ₁₂	461.0734		285	+	+	-	-	[27]
12	3,4-dicaffeoylquinic acid	22.22	C ₂₅ H ₂₄ O ₁₂	515.1201		353/191/179	+	+	+	+	Standard
13	Luteolin-4'-O-glucoside	23.84	C ₂₁ H ₂₀ O ₁₁	447.0939		285/ 447	+	+	-	-	Standard
14	Isovitexirone	24.41	C ₂₇ H ₄₂ O ₇	523.2917 *	479.2998	425/443/373/123/145/219/303/407/461/ 479 [#]	+	+	-	+	[27]
15	Vitexirone	26.77	C ₂₇ H ₄₂ O ₇	523.2908 *	479.2998	69/443/425/299/109/407/ 479 /461/281/311 [#]	+	+	+++	+	[28]
16	Ajugasterone C	32.70	C ₂₇ H ₄₄ O ₇	525.3061 *	481.3152	427/445/81/409/299/311/ 481 /463/189 [#]	+	+	++	-	[29]
17	Luteolin	38.57	C ₁₅ H ₁₀ O ₆	285.0405			+	-	-	+	Standard
18	3,7-Dimethylquercetin	40.99	C ₁₇ H ₁₄ O ₇	329.0664		314/299/271/243	+	-	-	-	Standard
19	Eupatorin	41.82	C ₁₈ H ₁₆ O ₇	343.0824		328/298/270	+	-	-	-	[30]
20	Phytuberin	42.33	C ₁₇ H ₂₆ O ₄	293.1759		236/221/ 293	-	+	-	-	[31]

noindent+: presence of the compound in the extract; ++: abundant compound; +++: very abundant compound; -: absence of the compound in the extract; L: leaves; TB: trunk bark; R: root bark; Fr: fruit. * (M+HCOO⁻); fragmented ions are highlighted in bold; # The MS² analyses were performed in positive mode.

2.1.1. Extracts of *V. madiensis*

To determine the chemical composition of each extract, we carried out LC/MS analyses either in positive and negative mode to identify the compounds present in the extract using the method developed by HPLC. As the positive mode did not give more information about the chemical profile, we only investigated the negative mode. The chromatographic profiles of the extracts of leaves, trunk bark, root bark and fruits of *V. madiensis* are presented in Figure 1.




Figure 1. LC-MS chromatograms in negative mode of methanolic extracts of *V. madiensis* (A) leaves; (B) trunk bark; (C) root bark; (D) fruits.

These results showed that the main compounds present in both extracts were ecdysteroids, considered as chemotaxonomic markers of the genus *Vitex* [32]. Among the family of ecdysteroids, we identified four major compounds: 20-hydroxyecdysone 10, isovitexirone 14, vitexirone 15 and ajugasterone C 16 (Figure 2).




Figure 2. Structures of some ecdysteroids identified in *V. madiensis*.

Other families of compounds have also been identified such as organic acid, flavonoids and fatty acids. Indeed, leaves and trunk bark extracts contain flavone glycosides such as homoorientin, orientin and vitexin; and flavonoids such as luteolin, 3,7-dimethylquercetin. The presence of caffeic acid derivatives such as dicaffeoylquinic acid were detected in both extracts (Table 1).

Comparison of the four chromatograms allowed us to conclude that the leaves extract contain more families of compounds (ecdysteroids, flavonoids, fatty acids) than the other parts. Root bark and fruits contain mainly ecdysteroids and the major compound of both extracts seemed to be 20-hydroxyecdysone. As the four LC/MS analyses were performed with the same concentration of extract, we could also compare the relative quantity of ecdysteroids, potentially responsible for the biological activities of each extract. We could estimate that the root bark extract is more concentrated in ecdysteroids than the other parts of the plant, and the leaves and fruits extracts seem to be rich in 20-hydroxyecdysone.

2.1.2. Extracts of *C. febrifuga*

As previously described, chemical profiles of extracts from leaves, trunk bark, root and fruit of *C. febrifuga* were performed by LC/MS analysis (Figure 3).

The chromatograms showed that the extracts of the different parts of *C. febrifuga* were quite similar events if some compounds were not present in the same concentration. The extracts contained especially organic acids, phenols, particularly chlorogenic acid derivatives, and flavonoids (quercetin derivatives). Organic acids such as citric acid, protocatechic acid and vanillic acid are present in the fruits extract. The other extracts contain mainly iridoid glycosides such as geniposidic acid, loganin, 11-methylxoside and glycosylated flavonoides such as hyperoside (Figure 4). The trunk barks contain geniposidic acid and quinic acid esters such as chlorogenic acid; the root bark extract contains caffeic acid diester, 3,4-dicaffeoylquinic acid and shanzhiside. The whole extracts contain the same unidentified compound possessing a retention time of 12.2 min. The structure of this compound has not been yet identified as it has not been described in *Crossopteryx* species. The identified compounds and their masses are presented in Table 2.

2.2. Antioxidant Activity of Extracts

The antioxidant potency of the methanolic extracts was evaluated by measuring their free radical scavenging activity using a DPPH assay. The IC_{50} (concentration providing 50% inhibition) was calculated graphically using a calibration curve in the linear range by plotting the extract concentration vs. the corresponding scavenging effect. The extracts from the leaves, stem barks and fruits of *V. madiensis* significantly inhibit the DPPH radical. The inhibitory concentrations (IC_{50}) are 110, 125 and 210 $\mu\text{g/mL}$, respectively. The leaves and trunk bark showed promising antioxidant potential with IC_{50} close to that of ascorbic

acid, taken as the reference antioxidant ($IC_{50} = 100 \mu\text{g}/\text{mL}$) (Table 3). These results are comparable with those obtained with other *Vitex species*, in particular *Vitex doniana*, *Vitex fischeri* and *Vitex keniensis* [33,34].




Figure 3. LC-MS chromatograms of *C. febrifuga* methanolic extracts: (A) leaves; (B) trunk bark; (C) root bark; (D) fruits.




Figure 4. Structures of some compounds identified in *C. febrifuga*.

Table 2. Compounds identified in extracts of leaves, trunk bark, root bark and fruits of *Crossopteryx febrifuga*.

N°	Composés	Tr (min)	Formula	M-H	MS ² (m/z)	L	TB	RB	F	Reference
1	Uronic acid	3.34	C ₆ H ₁₂ O ₇	195.0499	75/ 195 /129/87/99/89/85	+	+	++	+	Standard
2	Quinic acid	3.87	C ₇ H ₁₂ O ₆	191.0547	191 /85/127/93/85	+++	+++	+	+++	Standard
3	Sucrose	4.11	C ₁₂ H ₂₂ O ₁₁	341.1087		-	+	+	+	Standard
4	Citric acid	6.80	C ₆ H ₈ O ₇	191.0186	191 /129/111/87/85	+	-	+	+	Standard
5	Fumaric acid	7.34	C ₄ H ₄ O ₄	115.0032	71/ 115 /72/51	-	-	-	+	[35]
6	Shanzhiside	9.75	C ₁₆ H ₂₄ O ₁₁	391.1241	183/165/139/99/89/71/101/ 391	-	-	+	-	[36]
7	Geniposidic acid	10.03	C ₁₆ H ₂₂ O ₁₀	373.1135	123/149/211/167/ 373 /193	+	+	-	-	[37]
8	Ixoside	10.40	C ₁₆ H ₂₀ O ₁₁	387.0927	181/93/343/89/59/137/163/119/71/101/205/ 387	-	+	++	-	[38]
9	Protocatechuic acid	11.43	C ₇ H ₆ O ₄	153.0177	109/ 153 /110	-	-	-	+	[22]
10	Shanzhiside methyl ester	12.23	C ₁₈ H ₂₈ O ₁₃	451.1454 *	243/101/405/ 451	+	+++	+++	-	[39]
11	Chlorogenic acid	12.98	C ₁₆ H ₁₈ O ₉	353.0878		-	+	-	+	Standard
12	Rehmannioside A	13.47	C ₂₁ H ₃₂ O ₁₅	523.1664	293/89/233/71/125/477	-	-	+	-	[37]
13	4-Hydroxybenzoic acid	13.80	C ₇ H ₆ O ₃	137.0238	137 /138/136/109/108/81/119/93	-	-	-	+	[22]
14	Loganine	14.69	C ₁₈ H ₂₈ O ₁₂	435.1504 *	227/101/139/ 435	+	-	-	-	[40]
15	11-Methylxoside	15.03	C ₁₇ H ₂₂ O ₁₁	401.1086	101/137/ 401 /195/93/239/221	-	+	++	-	[41]
16	Hyperoside	18.10	C ₂₁ H ₂₀ O ₁₂	463.0881	300/ 463 /271/255/179	+	-	-	+	Standard
17	Isoquercetin	18.51	C ₂₁ H ₂₀ O ₁₂	463.0881	300/ 463 /271/255/179/151	+	-	-	-	Standard
18	Vanillic acid	19.78	C ₈ H ₈ O ₄	167.0338	167/152/111	-	-	-	+	Standard
19	3,4-dicaffeoylquinic acid	24.89	C ₂₅ H ₂₄ O ₁₂	515.1189		-	-	+	-	Standard
20	Azelaic acid	26.71	C ₉ H ₁₆ O ₄	187.0965	125/187/169/126/97/143	-	-	-	+	Standard

+: presence of the compound in the extract; -: absence of the compound in the extract; L: leaves, TB: trunk bark; R: root bark; F: fruit; * (M+HCOO⁻), fragmented ions are highlighted in bold.

Table 3. Inhibitory concentrations (IC₅₀s) of extracts.

	<i>V. madiensis</i>				<i>C. febrifuga</i>				Ascorbic Acid
	L	TB	R	F	L	TB	R	F	
IC ₅₀ (µg/mL)	110	125	980	210	100	110	200	710	100

Similarly for *C. febrifuga*, extracts from the leaves, trunk bark and root bark significantly inhibited the DPPH radical (Table 3), with IC₅₀s of 100, 110 and 200 µg/mL, respectively. Fruit extract did not have a significant effect (IC₅₀ = 710 µg/mL).

2.3. ROS Production Inhibition

In order to investigate further the anti-inflammatory and antioxidant effects of both plants, we examined the ability of the extracts to decrease cellular ROS production by incubating blood leukocytes from healthy donors stimulated by PMA. Stimulation with PMA, a direct PKC activator, resulted in a significant increase in ROS production after 2 h of incubation. Different concentrations of each extract of *V. madiensis* and *C. febrifuga* were tested (0 (Control), 10, 25, 50, and 100 µg/mL). Concerning the *V. madiensis* extracts, the leaf extract seemed most active reducing ROS production by 27 to 58% with increasing concentration, and this inhibition was statistically significant from a concentration of 25 µg/mL (Figure 5). The trunk bark and fruit extracts also reduced the production of ROS induced by PMA stimulation of leukocytes to become significant at concentrations of 50 and 100 µg/mL, respectively. The effect of the bark extract appeared to be dose-dependent at the lowest concentrations ($R^2 = 0.9979$). For root bark extract, there was a large inter-individual variability; nevertheless, ROS production appeared to decrease in a dose-dependent manner ($R^2 = 0.9915$).




Figure 5. Effect of *Vitex madiensis* extracts on ROS production of blood leukocytes stimulated with PMA for 2 h. Values are expressed as percentage of the Control (cells incubated with PMA and without extract). * $p < 0.05$, compared with Control normalized as 100% (dotted line).

With regard to *C. febrifuga*, results showed that none of the extracts had a real effect on the production of ROS of leukocytes stimulated by PMA, except for the fruit extract, which significantly inhibited this production compared to the control from 50 µg/mL until an effect of around 42% was reached at 100 µg/mL (Figure 6).




Figure 6. Effect of *Crossopteryx febrifuga* extracts on ROS production of blood leukocytes stimulated with PMA for 2 h. Values are expressed as percentage of the Control (cells incubated with PMA and without extract). * $p < 0.05$, compared with Control normalized as 100% (dotted line).

2.4. Effect of Extracts on Leukocyte Viability

We then examined the impact of the extracts on leukocyte viability in order to address the issue that decreased ROS production was related to decreased viability. This study was carried out under the same conditions as above, but this time over a period of 24 h using the resazurin test. It was shown that the leaves and trunk bark extracts of *Vitex madiensis* (Figure 7) as well as the fruit and root bark extracts of *Crossopteryx febrifuga* (Figure 8) did not show any effect on cell viability. No significant difference was observed for these extracts ($p > 0.05$) at all the concentrations tested compared to the 100% normalized Control (cells incubated with PMA and without extract). The other extracts showed effects on cell viability at 50 and 100 µg/mL at 24 h ($p < 0.05$).




Figure 7. Effect of *Vitex madiensis* extracts on leukocyte viability. Blood cells were treated with the indicated concentrations of extracts, in the presence of PMA for 24 h. Data were shown as means \pm SEM (Control = 100%), * $p < 0.05$ compared with Control.




Figure 8. Effect of *Crossopteryx febrifuga* extracts on leukocyte viability. Blood cells were treated with the indicated concentrations of extracts, in the presence of PMA for 24 h. Data were shown as means \pm SEM (Control = 100%), * $p < 0.05$ compared with Control.

3. Discussion

The chemical profiles of the methanolic extracts of various parts of *V. madiensis* were investigated and allowed us to identify four phytosterols: 20-hydroxyecdysone, ajugasterone C, isovitexirone and vitexirone. These ecdysteroids, found in some species of the *Vitex* genus, were identified in leaves, trunk bark and root bark, and fruits, confirming the hypothesis that ecdysteroids are taxonomic markers of the *Vitex* genus [32]. As previously described, 20-hydroxyecdysone is the most present ecdysteroids in *Vitex* species [32] and has been isolated from *V. madiensis* [16]. Ajugasterone C and vitexirone have been identified in *Vitex fisherii* [28], *Vitex leptobotrys* [42], *Vitex polygama* [43], *Vitex scabra* [44] and *V. madiensis*. Moreover, ecdysteroids, and especially 20-hydroxyecdysone, possess various biological activities such as anti-diabetic, antioxidant, anti-inflammatory, angiogenic, cardioprotective, neuroprotective, lung, kidney and gastric proactive activities. A recent study demonstrated that 20-hydroxyecdysone could prevent the appearance of severe forms of COVID-19 [45].

The extracts of *V. madiensis* showed a significant free radical scavenging activity compared to ascorbic acid. The leaves and trunk bark extracts possess the highest activity, confirming the antioxidant power of *V. madiensis*, as already observed in previous studies [19]. It should also be noted that many species of the *Vitex* genus, especially leaf and fruit extracts, show significant scavenging activity [16,19,33,46].

To our knowledge, the activity of *V. madiensis* extracts on the ROS production induced by blood leukocytes has not been described, and we found it relevant to explore this field. First, it was established that leaf extract at 25 $\mu\text{g}/\text{mL}$ reduced ROS production significantly after 2 h of incubation with stimulated leukocytes by PMA, without affecting the viability of cells. Our study also showed that the extracts of trunk bark and fruits of *V. madiensis* had an effect on ROS production by leukocytes, and it became significant at concentrations of 50 and 100 $\mu\text{g}/\text{mL}$, respectively. Even then, the viability study of leukocytes confirmed the antioxidant effect of *Vitex madiensis* extracts as it was not the consequence of a decrease in leukocyte viability. These results confer to the extracts both an anti-radical effect and a potential anti-inflammatory activity. Oxidative species are produced not only under pathological situations (inflammatory diseases, cancers, autoimmune diseases, etc . . .) but also during physiological situations such as cellular metabolism [47–49]. Indeed, in the body, ROS are chemical mediators that are involved in cellular communication and are part of the defence mechanisms in an inflammatory context [50–52]. On the other hand, if the

ROS production exceeds the defence capacity, they are able to induce direct cellular injury via oxidative stress to nucleic acids, cellular proteins and by activating lipid peroxidation, which destroys the cell membrane [31,53,54]. Thus, the neutralization of free radicals or reactive oxygen species by a naturally antioxidant extract is valuable for cell protection from oxidative stress.

The antioxidant activity and the ROS production inhibition of *V. madiensis* extracts could be due to the presence of 20-hydroxyecdysone and probably the ecdysteroids in high concentration in the plant. Indeed, ecdysteroids have already been described for their antioxidant capacity. In particular, ecdysterone exerts protective effects against lipid peroxidation from free radicals, obtaining a status of antioxidant [55,56].

The methanolic extracts of *C. febrifuga* contained mostly iridoids derivatives and flavonoids. The trunk bark and the root bark extracts were particularly rich in glycosylated iridoids such as ixoside, shanziside and methyl ester shanziside, which seem to be the major compounds of these two extracts. The fruits extract contained widely flavonoids and phenolic derivatives such as hyperoside and chlorogenic acid. The extracts of leaves, trunk bark and root bark of *C. febrifuga* showed significant antioxidant effects due to their free radical scavenging activity. The extracts of leaves and trunk bark present a higher antioxidant capacity compared to the root extract with an IC_{50} close to ascorbic acid. In the literature, several authors state that the antioxidant activity of plant extracts depends on its content of phenolic compounds [33,57,58]. In addition, flavonoids are the most important class of polyphenols that have shown greater potential for biological activities such as antioxidant, anti-inflammatory, antibacterial, anticancer and anti-allergic activity [59]. This partly coincides with our study, whose LC-MS profiles of the different extracts of *C. febrifuga*, and particularly the fruit extract, are characterized by a few phenolic and flavonoid compounds, which would be responsible for the antioxidant potential observed with DPPH.

The fruit extract did not show a significant effect on the DPPH radical, unlike previous studies carried out on the fruits of this plant whose results showed significant effects on the DPPH radical [60]. However, this fruit extract inhibited ROS production induced by blood leukocytes stimulated with PMA and this, significantly at concentrations above 50 $\mu\text{g}/\text{mL}$. It would be interesting to investigate which constituent could be involved, at least in part, in the protective effect against ROS toxicity.

4. Materials and Methods

4.1. Plant Material

The plant material (leaves, fruits, trunk bark and root bark) of *Vitex madiensis* Oliv. and *Crossopteryx febrifuga* (Afzel ex G. Don) BENTH was collected in the savannah zone, 25 km South of Brazzaville. After collection of the samples, one specimen of each plant species was authenticated and preserved in the herbarium of the Institut National de Recherche en Sciences Exactes et Naturelles (IRSEN) of Brazzaville under the number T7041 for *Vitex madiensis* Oliv. and B434 for *Crossopteryx febrifuga* (Afzel ex G. Don) BENTH. The plant material was dried in the shade for 7 days and then ground with a grinder.

4.2. Cell Material

Blood (collected from healthy volunteers, $n = 3$) was provided by the Etablissement Français du Sang (EFS) of the city of Clermont-Ferrand according to the contract n°16-21-62 (in accordance with the following articles L1222-1, L1222-8, L1243-4 and R1243-61 of the public health code). Whole blood leukocytes were obtained as previously described by Cholet et al. [61].

4.3. Methanolic Extracts

All the parts of the plants were air-dried grinded and then were macerated in methanol for 24 h (plant/solvent ratio: 1/10). In each case, the mixture was then filtered and evaporated under vacuo to dryness. The yields obtained for each part of the plant are

summarized in Table 4. The crude extract of each plant part is obtained and then stored at 4 °C before analysis.

Table 4. Yield of methanolic extraction of each part of plants (L: leaves, TB: trunk bark, RB: root bark, F: fruits).

Parts of plants	<i>Vitex madiensis</i>				<i>Crossopteryx febrifuga</i>			
	L	TB	RB	F	L	TB	RB	F
Mass of plants (g)	200	200	115	300	200	200	300	112
Mass of extract (g)	12	6	5.8	12	38	20	42	3.4
Yield (%)	6	3	5	4	19	10	14	3

4.4. Determination of the Chemical Profiles

The chemical profile of each crude extract was determined by chromatographic analyses using Ultra-High Performance Liquid Chromatography (UHPLC). HPLC-MS analyses were performed on an Ultimate 3000 RSLC UHPLC system (Thermo Fisher Scientific Inc., Waltham, MA, USA) coupled to a quaternary rapid separation pump (Ultimate autosampler) and a rapid separation diode array detector. Compounds were separated on an Uptisphere Strategy C18 column (250 × 4.6 mm, 5 µm, Interchim, Montluçon, France), which was controlled at 30 °C. The mobile phase was a mixture of 0.1% (*v/v*) formic acid in water (phase A) and 0.1% (*v/v*) formic acid in acetonitrile (phase B). The gradient of phase A was 100% (0 min), 80% (10 min), 73% (35 min), 0% (40–50 min) and 100% (51–60 min). The flow rate was 0.8 mL/min, and the injection volume was 5 µL. The UHPLC system was connected to an Orbitrap (Thermo Fisher Scientific Inc., Waltham, MA, USA) mass spectrometer, operated in the positive and negative electrospray ionization mode. Source operating conditions were: 3 kV spray voltage; 320 °C heated capillary temperature; 400 °C auxiliary gas temperature; sheath, sweep and auxiliary gas (nitrogen) flow rate 50, 10 and 2 arbitrary units, respectively; and collision cell voltage between 10 and 50 eV. Full scan data were obtained at a resolution of 70,000, whereas MS² data were obtained at a resolution of 17,500. Data were processed using Xcalibur software (Thermo Fisher Scientific Inc., City, MA, USA).

The sample to be analysed was prepared under the following operating conditions: 5 mg of the methanolic extract was diluted in 5 mL of MeOH of HPLC grade quality. The solution is then filtered with a 0.45 µm PTFE filter. Part of the filtrate is placed in a vial for analysis.

4.5. Antioxidant Properties

The in vitro antiradical activity was evaluated by the 2,2 diphenyl-1-picrylhydrazyl radical (DPPH) scavenging method. The protocol, previously described in literature, was used with some modifications [62,63].

Extract and ascorbic acid solutions in concentrations ranging from 200 to 1200 µg/mL, and 0.18 mM DPPH were prepared in methanol. In addition, 300 µL of each concentration of extracts and ascorbic acid were mixed with 3 mL of 0.18 mM DPPH solution. The kinetics were monitored for 45 min at 517 nm using a visible Jasco UV spectrophotometer (a methanolic solution was used as blank). The experiments were repeated three times, and the results were presented as a mean ± SEM. The inhibitory concentration 50 (IC₅₀) of each extract was determined. The lower the IC₅₀ value, the better the antiradical activity [64].

4.6. ROS Production Inhibition

The ROS production inhibition was determined by the method of inhibiting the production of reactive oxygen species (ROS) induced by human blood leukocytes.

The leukocyte preparations (*n* = 3) were obtained as previously described. The cells were placed in a 96 well polystyrene plate (Cell Wells, Corning, NY, USA), incubated

with methanolic extracts added at different concentrations (0, 10, 25, 50 and 100 µg/mL) and dihydrorhodamine 123 (DHR, 1 µM, Cayman Chemical Company, Ann Arbor, MI, USA) and finally stimulated or not by phorbol 12-myristate 13-acetate (PMA, 1 µM, Sigma-Aldrich, Saint Louis, USA). Fluorescence readings (excitation/emission: 485/535 nm) in kinetics of rhodamine 123, which is the product of DHR 123 oxidation by ROS, were taken over 2 h every 10 min using a Fluoroskan Ascent FL[®] apparatus (ThermoFisher Scientific, Illkirch, France) [65]. These experiments were repeated three times and the percentage of ROS production in the presence of the extracts was calculated. ROS production was calculated relative to 100% normalized control.

4.7. Effect of Extracts on Leukocyte Viability

Leukocytes preparations, obtained from the same donors, were placed in 96 well polystyrene plates incubated with methanolic extracts added at different concentrations (0, 10, 25, 50 and 100 µg/mL) and finally stimulated or not by PMA (1 µM). After 24 h of incubation at 37 °C under a 5% CO₂ atmosphere, resazurin (Sigma-Aldrich) was added to each well at the concentration 25 µg/mL. Leukocytes viability and then cytotoxicity of the different extracts were monitored by fluorescence reading (excitation/emission: 530/590 nm) using a Fluoroskan Ascent FL[®] instrument. The percentage of cell viability was calculated relative to 100% normalized control.

4.8. Data Analysis

Data were expressed as the mean ± SEM and represented at least three independent experiments. Statistical analysis and significance were measured using the paired, bilateral Student's *t*-test. Differences at $p \leq 0.05$ (flagged as *) were considered statistically significant.

5. Conclusions

V. madiensis is one of the species of the *Vitex* genus which synthesizes ecdysteroids, steroid hormones from invertebrates which are involved in the regulation of moulting, development and reproduction. The biosynthesis of these compounds is carried out, in a variable way, at the level of root bark, barks and fruits. In addition, 20-hydroxyecdysone, vitexirone, and ajugasterone C are the major compounds found in *Vitex* genus among the ecdysteroids family. The extracts of leaves, trunk bark and fruits of *V. madiensis* were studied and showed a free radical scavenging and anti-inflammatory activities; moreover, both extracts did not show any cytotoxic effect on blood cells. The extracts of *C. febrifuga*, mainly contain organic acids, sugars and phenolic compounds, but the chemical composition is very dependent on the organ of the plant. Despite a significant cytotoxic effect on blood cells, the extracts of leaves, trunk bark and root bark of *C. febrifuga* showed a significant inhibition of DPPH.

This study allowed us to demonstrate that leaves and trunk bark extracts of *V. madiensis* present an anti-inflammatory activity by decreasing ROS production. As previously described, we supposed that the activity is due to the presence of ecdysteroids, but we will isolate all the major compounds of both extracts to evaluate the potential anti-inflammatory activity.

Author Contributions: Conceptualization, A.L. and P.C.; Funding acquisition, A.L. and P.C.; Investigation, G.B.-T. and M.G.; Supervision, I.R., C.D., P.C., A.L., K.B., L.D. and F.C.-C.; Writing—original draft, G.B.-T., I.R., P.C. and C.D.; Writing—review and editing, I.R., P.C. and C.D. All authors have read and agreed to the published version of the manuscript.

Funding: This work was financially supported by the SCAC (“Service de Coopération et d’Action Culturelle”) French Embassy in Brazzaville, Congo. Funding number: Campus France 862970F.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki, and approved by French Public Health Code in accordance with the following articles L1222-1, L1222-8, L1243-4 and R1243-61.

Informed Consent Statement: Blood was collected from healthy human volunteers (Etablissement Français du Sang, EFS, Clermont-Ferrand, France). Donors gave their written informed consent for the use of blood samples for research purposes under EFS contract n°16-21-62.

Acknowledgments: We are grateful to Martin Leremboure from Clermont Auvergne University for mass analysis.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Salawu, O.A.; Chindo, A.Y.; Tijani, A.B.; Adzu, B. Analgesic, anti-inflammatory, antipyretic and antiplasmodial effects of the methanolic extract of *Crossopteryx febrifuga*. *J. Med. Plants Res.* **2008**, *2*, 213–218.
2. Chantaranothai, P. A revision of the genus *Vitex* (Lamiaceae) in Thailand. *Trop. Nat. Hist.* **2011**, *11*, 91–118.
3. Adjanohoum, E.J. *Contribution aux Etudes Ethnobotaniques et Floristiques en République Populaire du Congo*; ACCT: Paris, France, 1988; p. 605.
4. Audu, J.A. Medicinal Plants and their use in Bauchi State. *Big. Field* **1989**, *54*, 157–168.
5. Bridson, D.; Verdcourt, B.; Rubiaceae, I. *Flora of Tropical East Africa*; aa Balkema: Rotterdam, The Netherlands, 1988; p. 2.
6. Tomas-Barberan, F.A.; Hostettmann, K.A. Cytotoxic triterpenoid and flavonoids from *Crossopteryx febrifuga*. *Planta Med.* **1988**, *54*, 266–267. [[CrossRef](#)] [[PubMed](#)]
7. Danton, O. Extraction des Substances Naturelles Antalgiques à Partir des Plantes Utilisées Dans la Pharmacopée Traditionnelle au Mali. Ph.D. Thesis, Université Clermont Auvergne, Clermont-Ferrand, France, 2017.
8. Bouquet, A. *Féticheurs et Médecines Traditionnelles du Congo (Brazzaville)*; Travaux et Documents de l'ORSTOM: Paris, France, 1972; Volume 36, p. 282.
9. Elufioye, T.O.; Agbedahunsi, J.M. Antimalarial activities of *Tithonia diversifolia* (Asteraceae) and *Crossopteryx febrifuga* (Rubiaceae) on mice in vivo. *J. Ethnopharmacol.* **2004**, *93*, 167–171. [[CrossRef](#)] [[PubMed](#)]
10. Gariboldi, P.; Verotta, L.; Gabetta, B. Saponins from *Crossopteryx febrifuga*. *Phytochemistry* **1990**, *29*, 2629–2635. [[CrossRef](#)]
11. Tona, L.; Kambu, K.; Ngimbi, N.; Mesia, K.; Penge, O.; Lusakibanza, M.; Cimanga, K.; De Bruyne, T.; Apers, S.; Totte, J.; et al. Antiamoebic and spasmolytic activity of extract of some antidiarrhoeal traditional preparations used in Kinshasa Congo. *Phytomedicine* **2000**, *7*, 31–38. [[CrossRef](#)]
12. Latham, P. *Useful Plants of Bas-Congo Province, Democratic Republic of the Congo*; DFIG: London, UK, 2004; p. 320.
13. Arbonnier, M. *Arbres, Arbustes et Lianes des Zones Sèches d'Afrique de L'ouest*, 3rd ed.; Quae: Versailles, France, 2009.
14. Kerharo, J.; Adams, J.G. *La Pharmacopée Sénégalaise Traditionnelle, Plantes Médicinales et Toxiques*; Vigot Frères: Paris, France, 1974.
15. Ondo, J.P.; Lekana-Douki, J.P.; Bongui, J.B.; Zang Edou, E.S.; Zatra, R.; Toure-Ndoua, F.S.; Elombi, A.; Lebib, J.; Seguin, E. In vitro antiplasmodial activity and cytotoxicity of extracts and fractions of *Vitex madiensis*, medicinal plant of Gabon. *Trop. Med. Int. Health* **2012**, *17*, 316–321. [[CrossRef](#)] [[PubMed](#)]
16. Kubo, I.; Matsumoto, A.; Ayafor, J.F. Efficient Isolation of a Large Amount of 20-Hydroxyecdysone from *Vitex madiensis* (Verbenaceae) by Droplet Counter-Current Chromatography. *Agr. Biol. Chem.* **1984**, *48*, 1683–1684. [[CrossRef](#)]
17. Nguyen, T.K.T.; Ninh, K.B.; Do Thi, T.; Tran, M.L.; Vu Huong, G.; Nguyen, X.N.; Phan, V.K. Ecdysteroids from leaves of *Vitex trifolia*. *Vietnam. J. Chem.* **2018**, *56*, 162–166.
18. Meena, A.K.; Niranjan, U.S.; Rao, M.M.; Padhi, M.M.; Ramesh, B. A review of the important chemical constituents and medicinal uses of *Vitex* genus. *Asian J. Tradit. Med.* **2011**, *6*, 54–60.
19. Lengbiye, E.M.; Koto-te-Nyiwa, N. *Vitex madiensis* Oliv. (Lamiaceae): Phytochemistry, pharmacology and future directions, a mini-review. *J. Pharmacogn. Phytochem.* **2018**, *7*, 244–251.
20. Masateru, O.; Yasuyuki, I.; Satoko, K.; Toshihiro, N. Two new iridoids from *Vitex trifolia* Fructus (Fruit of *Vitex rotundifolia* L.). *Chem. Pharm. Bull.* **1997**, *45*, 1094–1096.
21. Chen, P.; Yi, W.; Lulin, C.; Wei, J.; Yan, N.; Qing, S.; Lu, G.; Quancheng, Z.; Licheng, Y.; Shufang, W. Comparison of the anti-inflammatory active constituents and hepatotoxic pyrrolizidine alkaloids in two *Senecio* plants and their preparations by LC–UV and LC–MS. *J. Pharmaceut. Biomed.* **2015**, *115*, 260–271. [[CrossRef](#)]
22. Vallverdú-Queralt, A.; Boix, N.; Piqué, E.; Gómez-Catalan, J.; Medina-Rejon, A.; Sasot, G.; Mercader-Martí, M.; Llobet, M.J.; Lamuela-Raventos, M.R. Identification of phenolic compounds in red wine extract samples and zebrafish embryos by HPLC-ESI-LTQ-Orbitrap-MS. *Food Chem.* **2015**, *181*, 146–151. [[CrossRef](#)]
23. Elhawary, S.S.; Younis, I.Y.; El Bishbishy, M.H.; Khatib, A.R. LC–MS/MS-based chemometric analysis of phytochemical diversity in 13 *Ficus* spp. (Moraceae): Correlation to their in vitro antimicrobial and in silico quorum sensing inhibitory activities. *Ind. Crops Prod.* **2018**, *126*, 261–271. [[CrossRef](#)]
24. Wang, Y.-H.; Avula, B.; Jadhav, B.-A.; Smillie, T.-J.; Khan, I.A. Structural characterization and identification of ecdysteroids from *Sida rhombifolia* L. in positive electrospray ionization by tandem mass spectrometry. *Rapid Commun. Mass Spectrom.* **2008**, *22*, 2413–2422. [[CrossRef](#)]
25. Zhou, J.; Qi, Y.; Hou, Y.; Zhao, J.; Li, Y.; Xue, X.; Wu, L.; Zhang, J.; Chen, F. Quantitative determination of juvenile hormone III and 20-hydroxyecdysone in queen larvae and drone pupae of *Apis mellifera* by ultrasonic-assisted extraction and liquid chromatography with electrospray ionization tandem mass spectrometry. *J. Chromatogr. B* **2011**, *879*, 2533–2541. [[CrossRef](#)]

26. Available online: <https://mona.fiehnlab.ucdavis.edu/spectra/display/CCMSLIB00005723108> (accessed on 22 December 2022).
27. Olenikova, D.N.; Kashchenkoa, N.I. Phytoecdysteroids of *Serratula centauroides* Herb from Cisbaikalia. *Russ. J. Bioorg. Chem.* **2019**, *45*, 913–919. [[CrossRef](#)]
28. Kubo, I.; Asaka, Y.; Stout, M.J.; Nakatsu, T. Structure of a novel phytoecdysteroid, vitexirone, and efficient isolation of phytoecdysteroids from *Vitex fisherii*. *J. Chem. Ecol.* **1990**, *16*, 2581–2588. [[CrossRef](#)] [[PubMed](#)]
29. Le Bizec, B.; Antignac, J.-P.; Monteau, F.; Andre, F. Ecdysteroids: One potential new anabolic family in breeding animals. *Anal. Chim. Acta* **2002**, *47*, 389–397. [[CrossRef](#)]
30. Feng, R.; Luya, L.; Zhang, X.; Zhang, Y.; Chen, Y.; Feng, X.; Zhang, L.; Zhang, G. Assessment of a developed HPLC-MS/MS approach for determining plasma eupatorin in rats and its application in pharmacokinetics analysis. *RSC Adv.* **2020**, *10*, 32020. [[CrossRef](#)] [[PubMed](#)]
31. Mittal, M.; Siddiqui, M.R.; Tran, K.; Reddy, S.P.; Malik, A.B. Reactive Oxygen Species in Inflammation and Tissue Injury. *Antioxid. Redox. Signal.* **2014**, *20*, 1126–1167. [[CrossRef](#)]
32. Sena Filho, J.G.; Duringerc, J.; Maiaa, G.L.A.; Tavaresa, J.F.; Haroudo, X.H.S.; Sobral da Silvaa, M.; Em.dio da-Cunhaa, V.L.; Barbosa-Filhoa, J.M. Ecdysteroids from *Vitex* Species: Distribution and Compilation of Their ¹³C-NMR Spectral Data. *Chem. Biodivers.* **2008**, *5*, 707–713. [[CrossRef](#)] [[PubMed](#)]
33. Sarr, S.O.; Fall, A.D.; Ngueye, R.; Diop, A.; Diatta, K.; Diop, N.; Ndiaye, B.; Diop, Y.M. Etude de l'activité antioxydante des extraits des feuilles de *Vitex doniana* (Verbenaceae). *Int. J. Biol. Chem. Sci.* **2015**, *9*, 1263–1269. [[CrossRef](#)]
34. Ochieng, C.O.; Nandwa, B.O. Proximate composition, phenolic content and antioxidant activities of three black plum (*Vitex* sp.) fruits: Preliminary results. *J. Food Technol.* **2010**, *8*, 118–125. [[CrossRef](#)]
35. Said, A.A.H.; Abuotabl, E.A.; Raoof, G.F.A. Identification of Constituents from *Pleio gynium timorens* (Dc.) Leenh Pericarp and Seeds Using High-Performance Liquid Chromatography with Electrospray Ionization Mass Spectrometry. *AASCIT J. Chem.* **2017**, *3*, 30–36.
36. Serralheiro, M.L.; Guedes, R.; Fadel, R.S.; Bendif, H. Data on identification of primary and secondary metabolites in aqueous extract of *Verbascum betonicifolium*. *Data Brief* **2020**, *32*, 106146. [[CrossRef](#)]
37. Maoyuan, W.; Qinglong, W.; Qing, Y.; Xiaoxia, Y.; Shixiu, F.; Zhunian, W. Comparison of Anthraquinones, Iridoid Glycosides and Triterpenoids in *Morinda officinalis* and *Morinda citrifolia* Using UPLC/Q-TOF-MS and Multivariate Statistical Analysis. *Molecules* **2020**, *25*, 160.
38. Ibekwe, N.N.; Adesomoju, A.A.; Igoli, J.O.; Barry, C.E.; Okogun, J.I. An iridoid glucoside from the leaves of *Pavetta crassipes*. *J. Chem. Soc. Niger.* **2019**, *44*, 629–632.
39. Available online: https://mona.fiehnlab.ucdavis.edu/spectra/display/Bruker_HCD_library000649 (accessed on 22 December 2022).
40. Xue, G.; Zhu, M.; Meng, N.; Guan, J.; Zhang, J.; Yang, J.; Wang, Y.; Cui, Y.; Chai, X. Integrating Study on Qualitative and Quantitative Characterization of the Major Constituents in Shuanghuanglian Injection with UHPLC/Q-Orbitrap-MS and UPLC-PDA. *J. Anal. Methods Chem.* **2021**, *22*, 1–11. [[CrossRef](#)] [[PubMed](#)]
41. Bealem, A.; Sakava, P.; Favier, L.; Hauchard, D.; Guegan, J.-P.; Nyemb, J.N.; Talla, E.; Henoumont, C.; Laurent, S.; Atchade De Theodore, A.; et al. Chemical Constituents of *Crossopteryx febrifuga* (*Afzel ex G. Don*) Benth (Rubiaceae), their Phytotoxicity and Antioxidant activities. *Nat. Prod. Chem. Res.* **2021**, *9*, 1000416.
42. Thuy, T.T.; Porzel, A.; Ripperger, H.; Sung, T.V.; Adam, G. Chalcones and ecdysteroids from *Vitex leptobotrys*. *Phytochemistry* **1998**, *49*, 2603–2605. [[CrossRef](#)]
43. Dos Santos, T.C.; Delle Monache, F.G.; Leita, S.G. Ecdysteroids from two Brazilian *Vitex* species. *Fitoterapia* **2001**, *72*, 215–220. [[CrossRef](#)]
44. Suksamrarn, A.; Kumpun, S.; Yingyongnarongkul, B. Ecdysteroids of *Vitex scabra* Stem Bark. *J. Nat. Prod.* **2002**, *65*, 1690–1692. [[CrossRef](#)]
45. Dinan, L.; Dioh, W.; Veillet, S.; Lafont, R. 20-Hydroxyecdysnoe, from Plant extracts to Clinical Use: Therapeutic Potential for the Treatment of Neuromuscular, Cardio-Metabolic and Respiratory Diseases. *Biomedicines* **2021**, *9*, 492. [[CrossRef](#)]
46. Ganapaty, S.; Vidyddhar, K.N. Phytoconstituents and biological activities of *Vitex*—A review. *J. Nat. Remedies* **2005**, *5*, 75–95.
47. Guzik, T.J.; Korbust, R.; Adamek-Guzik, T. Nitric oxide and superoxide in inflammation and immune regulation. *J. Physiol. Pharmacol.* **2003**, *54*, 469–487.
48. Reczek, C.R.; Chandel, N.S. The Two Faces of Reactive Oxygen Species in Cancer. *Annu. Rev. Cancer Biol.* **2017**, *1*, 79–98. [[CrossRef](#)]
49. Perl, A.; Gergely, P.; Banki, K. Mitochondrial dysfunction in T cells of patients with systemic lupus erythematosus. *Int. Rev. Immunol.* **2004**, *23*, 293–313. [[CrossRef](#)]
50. Herb, M.; Schramm, M. Functions of ROS in Macrophages and Antimicrobial Immunity. *Antioxidants* **2021**, *10*, 313. [[CrossRef](#)] [[PubMed](#)]
51. Novo, E.; Parola, M. Redox mechanisms in hepatic chronic wound healing and fibrogenesis. *Fibrogenesis Tissue Repair.* **2008**, *1*, 5–8. [[CrossRef](#)] [[PubMed](#)]
52. Santanam, N.; Ramachandran, S.; Parthasarathy, S. Oxygen Radicals, Antioxidants, and Lipid Peroxidation. *Semin. Reprod. Med.* **1998**, *16*, 275–280. [[CrossRef](#)] [[PubMed](#)]
53. Bulger, E.M.; Maier, R.V. Antioxidants in critical illness. *Arch. Surg.* **2001**, *136*, 1201–1207. [[CrossRef](#)]

54. Li, X.; Fang, P.; Mai, J.; Choi, E.T.; Wang, H.; Yang, X.-F. Targeting mitochondrial reactive oxygen species as novel therapy for inflammatory diseases and cancers. *J. Hematol. Oncol.* **2013**, *6*, 19. [[CrossRef](#)]
55. Cai, Y.-J.; Dai, J.-Q.; Fang, J.-G.; Ma, L.-P.; Hou, L.-F.; Yang, L.; Liu, Z.-L. Antioxidative and free radical scavenging effects of ecdysteroids from *Serratula strangulata*. *Can. J. Physiol. Pharmacol.* **2002**, *80*, 1187–1194. [[CrossRef](#)]
56. Das, N.; Mishra, S.K.; Bishatee, A.; Ali, A.S.; Bishayee, A. The Phytochemical, biological and medicinal attributes of phytoecdysteroids: An updated review. *Acta Pharm. Sin. B* **2021**, *11*, 1740–1766. [[CrossRef](#)]
57. Akanni, O.O.; Owumi, S.E.; Adaramoye, O.A. In vitro studies to assess the antioxidative, radical scavenging and arginase inhibitory potentials of extracts from *Artocarpus altilis*, *Ficus exasperata* and *Kigelia africana*. *Asian Pac. J. Trop. Biomed.* **2014**, *4*, 492–499. [[CrossRef](#)]
58. Fall, A.D.; Sy, A.N.; Fokou, J.B.H.; Fomi, J.O.N.; Dieng, M.; Dieng, S.I.M.; Bassene, E. Phytochemical screening, polyphenol content and antioxidant studies of ethanol leaf extract of *Combretum aculeatum* Vent. *Eur. J. Med. Plants* **2015**, *10*, 1–7. [[CrossRef](#)]
59. Anyasor, G.N.; Ogunwenmi, O.; Oyelana, O.A.; Akpofunre, B.E. Phytochemical constituents and antioxidant activities of aqueous and methanol stem extracts of *Costus afer* Ker Gawl. (Costaceae). *Afr. J. Biotechnol.* **2010**, *9*, 4880–4884.
60. Maiga, A.; Malterud, K.E.; Diallo, D.; Paulsen, B.S. Antioxidant and 15-lipoxygenase inhibitory activities of the malian medicinal plants *Diospyros abyssinica* (Hiern) F. white (Ebenaceae), *Lannea velutina* A. Rich (Anacardiaceae) and *Crossopteryx febrifuga* (Afzer) Benth (Rubiaceae). *J. Ethnopharmacol.* **2006**, *104*, 132–137. [[CrossRef](#)] [[PubMed](#)]
61. Cholet, J.; Decombat, C.; Varelle-Delarbre, M.; Gainche, M.; Berry, A.; Senejoux, F.; Ripoche, I.; Delort, L.; Vermery, M.; Fraisse, D.; et al. In vitro anti-inflammatory and immunomodulatory activities of an extract from roots of *Buplerum Rotundifolium*. *Medicines* **2019**, *6*, 101. [[CrossRef](#)]
62. Brand-Williams, W.; Cuvelier, M.E.; Berset, C. Use of free radical method to evaluate antioxidant activity. *Lebensm. Wiss. Technol.* **1995**, *28*, 25–30. [[CrossRef](#)]
63. Meda, N.R.; Fraisse, D.; Gnoula, C.; Vivier, M.; Felgines, C. Characterization of antioxidants from *Detarium microcarpum* Guill. et Perr. leaves using HPLC-DAD coupled with pre-column DPPH assay. *Eur. Food Res. Technol.* **2017**, *243*, 1659–1666. [[CrossRef](#)]
64. Sahgal, G.; Ramanathan, S.; Sasidharan, S.; Mordi, M.N.; Ismail, S.; Mansor, S.M. In vitro antioxidant and xanthine oxidase inhibitory activities of methanolic *Swietenia mahagoni* seed extracts. *Molecules* **2009**, *14*, 4476–4485. [[CrossRef](#)]
65. Cholet, J.; Decombat, C.; Varelle-Delarbre, M.; Gainche, M.; Berry, A.; Ogeron, C.; Ripoche, I.; Delort, L.; Vermerie, M.; Fraisse, D.; et al. Comparison of The Anti-Inflammatory and Immunomodulatory Mechanisms of Two Medicinal Herbs: Meadowsweet (*Filipendula ulmaria*) and *Harpagophytum* (*Harpagophytum procumbens*). *Int. J. Plant Anim. Environ. Sci.* **2019**, *9*, 145–163.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.