



HAL
open science

Outdoor shelter for grazing sheep: what impact of tree density on sheep behaviour, welfare and performance?

Cécile Ginane, Véronique Deiss, Mickaël Bernard, Camille Béral, Dorothée Bizeray-Filoché

► To cite this version:

Cécile Ginane, Véronique Deiss, Mickaël Bernard, Camille Béral, Dorothée Bizeray-Filoché. Outdoor shelter for grazing sheep: what impact of tree density on sheep behaviour, welfare and performance?. 8. INTERNATIONAL CONFERENCE ON THE ASSESSMENT OF ANIMAL WELFARE AT FARM AND GROUP LEVEL, Teagasc - University College Cork, Aug 2021, Virtual conference, Ireland. hal-04181888

HAL Id: hal-04181888

<https://hal.inrae.fr/hal-04181888>

Submitted on 16 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Outdoor shelter for grazing sheep: what impact of tree density on sheep behaviour, welfare and performance?

Cécile Ginane¹, Véronique Deiss¹, Mickaël Bernard¹, Camille Béral², Dorothée Bizeray-Filoche³

¹INRAE, Centre Clermont Auvergne Rhône Alpes, site de Theix, 63122 Saint-Genès-Champanelle, France, ²AGROOF SCOP, 19 rue du Luxembourg, 30140 Anduze, France, ³UniLaSalle, Rue Pierre Waguet, 60026 Beauvais, France; cecile.ginane@inrae.fr

Ruminants reared outdoors are confronted with various climatic hazards that can affect their welfare, and trees can provide potential shelter. The present study tested the effect of tree density in pasture on the behaviour, welfare and performance of sheep, in different geographical contexts. Six grassland plots in north-west (NWG) and three in a semi-mountainous area of central France (SMG) were investigated. The NWG, consisting of lowland meadows with adult fruit trees, were grazed by groups of minimum 15 ewes. Plots either contained few trees (Control, C) or were treed (T). The SMG, planted with adult cherry wood and maple trees at different densities (C: 1 tree, T+: 60 trees/ha, T++: 150 trees/ha) were continuously grazed by 10 ewes (and their lambs until weaning) over the grazing season. Ewes' activity and position relative to shade were observed over periods of 2 to 4 days (6 to 7.5h / day) in spring and summer of 2016 and 2017, simultaneously on plots C and T, or C, T+ and T++. Reactions to insects and respiratory rate were also carried out, as well as ewes and lambs' performance in SMG. Data were analysed with the SAS Enterprise Guide® software (Proc MIXED), for testing the effects of treatment and date, season or year (repeated statement) depending on the analysis. The proportion of time spent in the shade increased with tree density (35% (C) vs. 43% (T), $p < 0.001$ in NWG; 30% (C) vs. 73% (T+) vs. 88% (T++), $p < 0.0001$ in SMG), as well as with the increase in a heat load index (HLI, calculated in SMG from meteorological data, $p < 0.0001$). Animals actively selected the shade provided by the trees, even outside the hottest summer days, and despite greater reactions to insects under the trees ($p < 0.01$). Ewes mobilized their reserves more strongly on the tree-planted plots ($P < 0.05$), but only in 2016. All of them ensured a correct growth of their lambs. Trees on grassland represent an enrichment that the sheep actively use for shelter.