



HAL
open science

Are vaccine-induced antibodies useful for phagocytosis of mastitis associated *Escherichia coli* ?

Pierre Germon, Florence B Gilbert, Pascal Rainard

► To cite this version:

Pierre Germon, Florence B Gilbert, Pascal Rainard. Are vaccine-induced antibodies useful for phagocytosis of mastitis associated *Escherichia coli*?. 7. European Veterinary Immunology Workshop (EVIW), Aug 2021, On-line conference, France. hal-04182557

HAL Id: hal-04182557

<https://hal.inrae.fr/hal-04182557v1>

Submitted on 17 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

INRAE

➤ Are **vaccine-induced** antibodies useful for phagocytosis of **mastitis** associated *Escherichia coli* ?



➤ Mastitis in numbers...


- Still a disease with significant impact
 - High prevalence
 - 30-35% of dairy cows des animaux ont une mammite au cours d'une lactation
 - High economic cost
 - Cost : 230€ per case (IDELE)
 - Overall cost : 121 to 165 € /cow (Hogeveen 2019)
 - Reduced producing lifetime (2,5 lactations per cow)
 - High antibiotic usage (70% of antibiotics in dairy farms => mastitis)
- Pathogens involved (Poutrel, 2015)
 - Clinical mastitis : *Streptococcus uberis* (22 %), *Escherichia coli* (16 %), *Staphylococcus aureus* (16 %)
 - Sub-clinical mastitis : *Streptococcus uberis* (20-30 %), *S. aureus* (20-30 %)



INRAE

Are vaccine-induced antibodies useful for phagocytosis of mastitis associated *Escherichia coli* ?
August 2021 – 7th European Veterinary Immunology Workshop

➤ Mastitis: **inflammation** of the mammary gland, in general of **bacterial** origin



1. Entry of the pathogen through the teat canal
2. Bacterial multiplication in milk
3. Recognition of bacteria by the host immune system => inflammation
4. Production of anti-microbial peptides + phagocytosis by neutrophils => bacterial clearance

➤ One strategy to improve bacterial clearance is **vaccination to promote antibody-mediated opsonisation** and increase phagocytosis by neutrophils...


➤ This is the rationale behind the use of the *E. coli* J5 vaccine to prevent *E. coli* mastitis



INRAE

Are vaccine-induced antibodies useful for phagocytosis of mastitis associated *Escherichia coli* ?
August 2021 – 7th European Veterinary Immunology Workshop

1. Are J5 vaccine-induced antibodies able to reach their target antigen ?



➔ J5 vaccine-induced antibodies only weakly recognize smooth strains




INRAE

Are vaccine-induced antibodies useful for phagocytosis of mastitis associated *Escherichia coli* ?

August 2021 – 7th European Veterinary Immunology Workshop

2. Are antigens responsible for labeling of rough strains accessible to antibodies in smooth strains ?






➔ Result: Adsorption with smooth strain doesn't remove antibodies recognizing rough strains (case **#2**)

➔ Conclusion: Rough strains surface antigens recognized by J5-antiserum are not accessible on smooth strains





3. Why is a small fraction of smooth strains still labelled with J5 antiserum ?

Hypothesis : labeling is due to expression of type 1 fimbriae



- ➔ **Result:** Adsorption of J5 antiserum with fimbriated strain removes antibodies recognizing smooth P4 strains
- ➔ **Conclusion:** Labeling of strain P4 by J5 antiserum is linked to labeling of type 1 fimbriae

4. Are J5-induced antibodies improving phagocytosis by neutrophils ?



→ J5-induced antibodies do not improve phagocytosis compared to pre-immune serum

5. Summary

- ➔ J5 vaccine-induced antibodies only weakly recognize smooth strains
- ➔ Rough strains surface antigens recognized by J5-antiserum are not accessible on smooth strains (case #2)
- ➔ Labeling of strain P4 by J5 antiserum is linked to labeling of type 1 fimbriae
- ➔ J5-induced antibodies do not improve phagocytosis compared to pre-immune serum

Rainard, P., M. Reperant-Ferter, C. Gitton and P. Germon (2021). Shielding Effect of *Escherichia coli* O-Antigen Polysaccharide on J5-Induced Cross-Reactive Antibodies. *mSphere* 6(1).

 INRAE

Are vaccine-induced antibodies useful for phagocytosis of mastitis associated *Escherichia coli* ?
August 2021 – 7th European Veterinary Immunology Workshop