



**HAL**  
open science

## Dépérissement des chênes et communautés d'insectes des canopées

Elodie Le Souchu, Guilhem Parmain, Stéphanie Bankhead-Dronnet, Mathias Brand, Sébastien Damoiseau, Christian Sallé, Christophe Bouget, Aurélien Sallé

► **To cite this version:**

Elodie Le Souchu, Guilhem Parmain, Stéphanie Bankhead-Dronnet, Mathias Brand, Sébastien Damoiseau, et al.. Dépérissement des chênes et communautés d'insectes des canopées. Réunion annuelle du Groupe des Entomologistes Forestiers Francophones, GEFF, Sep 2022, Lamotte-Beuvron, France. hal-04185484

**HAL Id: hal-04185484**


**<https://hal.inrae.fr/hal-04185484>**

Submitted on 22 Aug 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





# Dépérissement des chênes et communautés d'insectes des canopées


Contrasting responses of trophic guilds to forest decline  
reshape canopy insect community

E. Le Souchu, G. Parmain, S. Bankhead-Dronnet, M. Brand, S. Damoiseau, C. Sallé, C. Bouget & A. Sallé





Percent of total mortality attributable to drought



**Climate change:** increased severity and frequency of droughts & heatwaves (Allen et al. 2010)

- Large-scale forest disturbances
- Large-scale forest declines and diebacks
- **Large-scale degradation of forest canopies**




Declining oaks – © A. Sallé




**Forest decline:** gradual degradation of forest health leading to tree death after several years. Leads to **collapses and pulses of microhabitats and trophic resources.**

- **Collapses:** Rapid crown alteration with loss of foliage & change in foliage quality and microclimates.
- Deleterious effects on foliage-feeding species (phyllophagous, gall-inducer, ...)?



Overview of decline-induced changes in arthropod communities (upper insets), resources and habitats (medium insets), and microclimates (lower insets) in forest canopies.

Sallé et al. 2021 Front. For. Glob. Change



Declining tree (left) & healthy tree (right)

© A. Sallé



→ **Pulses:** Positive effects on saproxylic species (mycophagous, xylophagous, ...) and flower-dependent taxa ?



Quantity and diversity of dead wood

Sap flow

Opportunistic fungi



Trunk cavities

Soil resources

Floral resources



- **Question:**

What are the effects of oak decline on the functional structure of the community of canopy-dwelling insects ?

- **Objectives:**

To characterize the community of canopy insects.

To assess the effects of decline on this community and its trophic guilds.


- **Hypotheses:**

- I. The level of decline influences the species composition, due to species turnover
- II. Contrasting effects of forest decline on abundance and species richness of larval trophic guilds



**Study areas in 2019:** Oak dominated forests.

H: Healthy stand - MD: Moderately declining stand - D: Declining stand



**Dieback surveys in 2019 and 2020:**

Quantification of crown degradation (DEPERIS protocol) on 10 trees/plot and 30 trees/stand



MB = 0  
MR = 0  
DEPERIS = 0/A



MB = 1  
MR = 2  
DEPERIS = 2,6/C



MB = 1  
MR = 3  
DEPERIS = 3,4/D



MB = 2  
MR = 3  
DEPERIS = 3,8/E



**Sampling:** 2 trees/stand, with **one green multi-funnel trap + one flight-interception trap** on each.

→ 1 plot = 1 tree with the two types of traps + 9 trees around the trap tree

Traps hanged in the **oak canopy** (10-15 m).

Sampling performed over the **activity period of insects** (from April to September 2019).

**Identification:** At the **lowest taxonomic level**.

**Analysis:** Data were pooled at the plot level



Oak canopy (left) with green multi-funnel trap (middle) & flight-interception trap (right). © E. Le Souchu




## Community overview

Approx. **132,000 indiv.** for **1,569 species.**

Divided into **7 orders**, **127 families** and **16 larval trophic guilds.**

Species richness of sampled orders



### Larval trophic guilds

#### Plant-based diet

Gall-inducer  
Rhizophagous  
Seminiphagous  
Phyllophagous  
Sap-feeder

#### Flower-dependent

Anthophagous  
Pollinivorous/Nectarivorous

#### Wood-based diet

Saproxylophagous  
Xylophagous

#### Polyphagous

Social polyphagous  
Polyphagous

#### Prey/host-dependent

Parasitoid  
Zoophagous  
Zoophyllophagous

#### Others

Mycophagous


Saprophagous




## Community overview

**Abundance of larval trophic guilds  
(total : 132,000 ind.)**

- Xylophagous
- Phyllophagous
- Zoophagous
- Parasitoid
- Rhizophagous
- Anthophagous
- Saprophagous
- Sap feeder
- Zoophyllophagous
- Gall inducer
- Mycophagous
- Saproxylophagous
- Social Polyphagous
- Pollinivorous
- Seminiphagous
- Polyphagous




**Species richness of larval trophic guilds  
(total : 1,569 sp.)**





## Community richness & composition



The level of plot decline **has an effect on species composition of the community**

PERMANOVA (999 permutations ;  $R^2 = 0,13$  ;  $pval = 0.001$  \*\*\*)

The decline **has no effect on species richness**


GLMER (neg. binom.; best model = null model)

**The decline reshapes the community but does not make it neither richer nor poorer.**

NMDS ordination (k=3, stress=0.13) of the assemblages of species by site (stand and tree), grouped by levels of plot decline



## Community richness & composition




The associations between the spatial scale analysed and each diversity component


Decline levels: Healthy, Moderate decline and Declining

$\gamma = 1,569$

### Additive partitioning of the species diversity components



## Community richness & composition




The associations between the spatial scale analysed and each diversity component

Decline levels: Healthy, Moderate decline and Declining

$\gamma = 1,569$

### Additive partitioning of the species diversity components



Turnover contributes to change species composition, among levels of plot decline




## Larval trophic guilds

	Larval trophic guild	Decline effect on guild abundance	Decline effect on species richness
Plant based diet	Gall-inducer	Negative *	ns
	Rhizophagous	ns	ns
	Seminiphagous	ns	ns
	Phyllophagous	ns	ns
	Sap feeder	ns	ns
Flower dependant	Anthophagous	Positive then negative d1: . ; d2: *	ns
	Pollinivorous	Positive d1: * ; d2: *	ns
Wood based diet	Xylophagous	ns	ns
	Saproxylophagous	ns	ns
Polyphagous	Polyphagous	ns	ns
	Social polyphagous	ns	ns
Prey/host dependent	Parasitoid	ns	ns
	Zoophagous	ns	ns
	Zoophytophagous	ns	ns
Others	Mycophagous	Positive then negative d1: *** ; d2: ***	ns
	Saprophagous	ns	ns
TOTAL		ns	ns


## Larval trophic guilds

	Larval trophic guild	Decline effect on guild abundance	Decline effect on species richness
Plant based diet	Gall-inducer	Negative	*
	Rhizophagous	ns	ns
	Seminiphagous	ns	
	Phyllophagous	ns	
	Sap feeder	ns	
Flower dependant	Anthophagous	Positive then negative	
	Pollinivorous	Positive	
Wood based diet	Xylophagous	ns	
	Saproxylophagous	ns	
Polyphagous	Polyphagous	ns	
	Social polyphagous	ns	
Prey/host dependent	Parasitoid	ns	
	Zoophagous	ns	
	Zoophytophagous	ns	
Others	Mycophagous	Positive then negative	
	Saprophagous	ns	
TOTAL		ns	






## Larval trophic guilds



## Larval trophic guilds

	Larval trophic guild	Decline effect on guild abundance
Plant based diet	Gall-inducer	Negative
	Rhizophagous	ns
	Seminiphagous	ns
	Phyllophagous	ns
	Sap feeder	ns
Flower dependant	Anthophagous	Positive then negative
	Pollinivorous	Positive
Wood based diet	Xylophagous	ns
	Saproxylophagous	ns
Polyphagous	Polyphagous	ns
	Social polyphagous	ns
Prey/host dependent	Parasitoid	ns
	Zoopagous	ns
	Zoophytophagous	ns
Others	Mycophagous	Positive then negative
	Saprophagous	ns
TOTAL		ns





## Larval trophic guilds

	Larval trophic guild	Decline effect on guild abundance	Decline effect on species richness
Plant based diet	Gall-inducer	Negative *	ns
	Rhizophagous	ns	ns
	Seminiphagous	ns	ns
	Phyllophagous	ns	ns
	Sap feeder	ns	ns
Flower dependant	Anthophagous	Positive then negative	d1: . ; d2: *
	Pollinivorous	Positive	d1: * ; d2: *
Wood based diet	Xylophagous	ns	<p>Contradicts previous results of decline and dieback on saproxylic species e.g. Beudert et al. 2015; Kozak et al. 2021; Cours et al. 2021</p> <p>Few dendro-microhabitats left in managed forests ?</p>
	Saproxylophagous	ns	
Polyphagous	Polyphagous	ns	
	Social polyphagous	ns	
Prey/host dependent	Parasitoid	ns	
	Zoophagous	ns	ns
	Zoophytophagous	ns	ns
Others	Mycophagous	Positive then negative	d1: *** ; d2: ***
	Saprophagous	ns	ns
TOTAL		ns	ns



## Conclusion

### Community richness and composition:

- The decline level modifies the community composition but not the species richness.
- Species turnover contributes to community modification.

### Larval trophic guilds:

- Few contrasting responses of trophic guild abundance.
- No effect on the species richness by guild.

- Management buffers changes in microhabitats and resources between declining and healthy stands ?
- Integration of both larval AND adult trophic guilds as response variable ?
- Too many species with different ecological traits within each trophic guild ? Use sub-guilds or taxonomic groups instead ?



*Philaenus spumarius*

© S. Damoiseau

### Improvement of conservation strategies:

Decline reshapes the insect community and acts as a driver of diversity.

Promoting a mosaic of healthy and declining patches within a forest would conserve canopy insect diversity.



# Thank you for your attention !

Mail : [elodie.lesouchu@gmail.com](mailto:elodie.lesouchu@gmail.com)



## Acknowledgements

Région Centre Val-de-Loire (financer)

Insect taxonomists

Carl Moliard, Xavier Pineau & Barbara Joncour (technicians)


Erwann Marhic, Léa Garreau, Manon Durand, Nathan Jolivard, Lisa Desbrée (students)

## Species richness

According to species richness estimators:  
between **1,799 and 2,371** sp. expected (67 – 87% of observed sp.)

Completeness of species composition detection estimated using various indices  
(Chao, Bootstrap and Jackknife 1 & 2)

Area	Expected					Observed
	Chao	Jackknife 1	Jackknife 2	Bootstrap	Range	
<b>Total</b>	2,150	2,086	2,372	1,799	1,799 – 2,372	<b>1,569 (66 -87%)</b>
<b>Orléans</b>	1,480	1,367	1,593	1,154	1,154 – 1,593	<b>989</b>
<b>Vierzon</b>	1,575	1,456	1,688	1,240	1,240 – 1,688	<b>1,071</b>
<b>Marcenat</b>	1,223	1,170	1,333	1,000	1,000 – 1,333	<b>860</b>



## Community composition

	Factors	df	F.Model	R <sup>2</sup>	p-value
	Decline	2	3.98	0.13	0.001 ***
Community composition	Forest	2	6.73	0.22	0.001 ***
	Decline*Forest	3	1.83	0.09	0.01 *

Effect of the degree of stand decline and forest on the entire insect community studied.  
PERMANOVAS made with 999 permutations. p < 0.05: \*; p < 0.01: \*\*; p < 0.001: \*\*\*.

**Decline effect on the overall composition of species** and guild

+ all the guilds except social polyphagous and sap feeder

**Forest effect on the overall composition of species** and guild

+ all the guilds except social polyphagous and zoophagous