



HAL
open science

Chronical heat stress modifies muscle metabolism and improves meat quality traits in pigs

Bénédicte Lebret, Aira Maye Serviento, David Renaudeau

► **To cite this version:**

Bénédicte Lebret, Aira Maye Serviento, David Renaudeau. Chronical heat stress modifies muscle metabolism and improves meat quality traits in pigs. 69. International Congress of Meat Science and Technology (Icomst), Aug 2023, Padova, Italy. pp.48-49. hal-04186010

HAL Id: hal-04186010

<https://hal.inrae.fr/hal-04186010>

Submitted on 18 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

• 2023 PADOVA, ITALY

ICoMST

69th INTERNATIONAL CONGRESS of
MEAT SCIENCE AND TECHNOLOGY

Proceedings of the 69th International Congress of Meat Science and Technology

from Tradition to Green Innovation



August 20-25, 2023
Padova, Italy

Edited by:

Antonella Dalle Zotte

Alberto Brugiapaglia

Marco Cullere

www.icomst2023.com

CHRONICAL HEAT STRESS MODIFIES MUSCLE METABOLISM AND IMPROVES MEAT QUALITY TRAITS IN PIGS

Bénédicte Lebret^{1*}, Aira M. Serviento¹ and David Renaudeau¹

¹PEGASE, INRAE, Institut Agro, 35590 Saint-Gilles, France

*Corresponding author email: benedicte.lebret@inrae.fr

I. INTRODUCTION

Chronic heat stress (HS) during pig growing compromises pork production efficiency due to lower growth performance resulting from reduced feed intake, as adaptive response to limit metabolic heat production [1]. The lower feed intake associated to the disturbed endocrine homeostasis induced by HS can also affect carcass composition, depending on intensity and duration of HS and pig's physiological stage [1, 2]. Effects of HS on muscle properties and pork quality are also controversial [2] and may depend on muscle metabolic type [3]. Moreover, animal physiological responses to HS during growing (GHS) can be affected by a previous HS experienced during prenatal period (PHS), as recently shown by Serviento *et al.* [4]. A better understanding of HS consequences on pork production and quality is of high importance for actors of the pork sector as more frequent and intense heat waves are highly likely to occur in the future due to global warming. Thus, this study aimed to evaluate the effects of PHS and/or GHS on muscle metabolism and pork quality in various muscle metabolic types.

II. MATERIALS AND METHODS

A total of 48 female pigs from 24 litters were used in the present study. From 6 to 109 d of gestation, 12 sows were kept under thermoneutral (TN) conditions (cyclic 18 to 24°C; PTN) and 12 sows under chronic HS (cyclic 28 to 34°C; PHS). Two female offspring per sow were randomly selected and housed from 82 to 140 d of age under cyclic GTN (18 to 24°C; n = 24) or GHS (28 to 34°C; n = 24) environments. There were four experimental treatments based on G and P thermal environments: GTN-PTN, GTN-PHS, GHS-PTN, and GHS-PHS. Pigs were fed *ad libitum* with a growing-finishing diet and were slaughtered at 140 d of age [4]. Pig growth and carcass traits were recorded. Quality traits were measured on *Longissimus thoracis* (LT) and *Semimembranosus* (SM) glycolytic and on *Semispinalis* (SS, neck) oxidative muscles. Metabolic properties of LT and SS muscles were determined as described previously [5]. Data were analysed using a mixed model (SAS software v9.4) with P, G and their interaction (P x G) as fixed effects, and sire and slaughter day as random effects. Final body weight was included as covariate for backfat thickness and carcass lean meat content.

III. RESULTS AND DISCUSSION

Compared with GTN, GHS pigs had lower growth rate and final body weight (Table 1) due to lower feed intake (not shown). GHS pigs had thicker backfat, especially for GHS-PHS pigs whereas carcass lean meat content did not differ according to P or G environments. Increased carcass fatness of pigs under PHS [6] or GHS [1] were already found and present data show that combining P and G heat stress promoted carcass fatness. GHS pigs had higher ultimate pH than GTN pigs in LT and SM, but not in the oxidative SS (Table 1). GHS pigs had lower LT drip loss, especially for GHS-PTN vs GTN-PTN pigs, and lower LT chroma but similar lightness values to GTN pigs (not shown). GHS pigs had lower glycolytic potential and activities of energy metabolism in LT compared with GTN pigs, but similar lipid content. In contrast, SS glycolytic potential and metabolic enzyme activities (not shown) were not modified by GHS. Thus, growing HS induced muscle-specific metabolic responses, with reduced energy metabolism in the glycolytic LT but not in the oxidative SS, leading to improved technological quality and appearance (lower drip) of pork loin. The altered LT energy metabolism of GHS pigs, which

could result from their altered thyroid function [4], is in line with other findings [2, 7]. In contrast, prenatal HS had scarce effects on muscle metabolism and pork quality.

Table 1. Growth and carcass, muscle and meat quality traits according to thermal environment

| Thermal treatment ¹ | GTN-PTN | GTN-PHS | GHS-PTN | GHS-PHS | RMSE | Significance ² | | |
|--|-------------------|--------------------|-------------------|--------------------|------|---------------------------|--------|-------|
| | | | | | | P | G | P x G |
| Final body weight, kg | 104.4 | 105.8 | 98.8 | 95.6 | 7.0 | 0.667 | <0.001 | 0.268 |
| Average daily gain, g/d | 1110 | 1129 | 984 | 966 | 93 | 0.990 | <0.001 | 0.492 |
| Hot carcass weight, kg | 80.8 | 81.7 | 77.0 | 74.2 | 5.5 | 0.548 | 0.001 | 0.254 |
| Backfat thickness, mm | 10.2 ^a | 10.0 ^a | 10.5 ^a | 12.5 ^b | 1.8 | 0.094 | 0.038 | 0.043 |
| Lean meat content, % | 63.5 | 63.4 | 63.2 | 61.8 | 1.4 | 0.077 | 0.074 | 0.013 |
| Longissimus thoracis muscle | | | | | | | | |
| pH 24 h | 5.51 | 5.50 | 5.59 | 5.54 | 0.07 | 0.143 | 0.006 | 0.355 |
| Drip loss 1-3 d p.m., % | 4.03 ^b | 3.15 ^{ab} | 2.62 ^a | 3.16 ^{ab} | 1.19 | 0.611 | 0.048 | 0.046 |
| Intramuscular fat, % | 1.45 | 1.47 | 1.48 | 1.65 | 0.42 | 0.430 | 0.395 | 0.552 |
| Glycolytic potential, $\mu\text{mol/g}$ | 174 | 168 | 144 | 152 | 18 | 0.817 | <0.001 | 0.186 |
| Lactate dehydrogenase, $\mu\text{mol/min/g}$ | 2307 | 2271 | 2148 | 2141 | 162 | 0.653 | 0.004 | 0.752 |
| Citrate synthase, $\mu\text{mol/min/g}$ | 7.54 | 7.77 | 6.80 | 6.92 | 1.08 | 0.582 | 0.014 | 0.860 |
| Semimembranosus: pH 24 h | 5.54 | 5.56 | 5.67 | 5.65 | 0.12 | 0.972 | 0.004 | 0.538 |
| Semispinalis: pH 24 h | 6.07 | 5.99 | 6.14 | 6.01 | 0.29 | 0.214 | 0.596 | 0.779 |

¹ GTN: growing thermoneutral, PTN: prenatal thermoneutral, GHS: growing heat stress, PHS: prenatal heat stress.

² P values of the fixed effects of prenatal (P), growing temperature (G), their interaction (PxG) and Root Mean Square Error obtained from the mixed model. a, b : differences between thermal treatments (P < 0.05).

IV. CONCLUSION

The present study confirms that chronic HS during growing impairs growth and carcass composition especially for pigs previously subjected to prenatal HS. Growing HS induces specific metabolic responses according to muscle type, leading to improved technological quality of pork loin and ham.

ACKNOWLEDGEMENTS

Authors thank the staff of INRAE experimental farm and slaughterhouse (UE 3P, 35590 Saint-Gilles, France) and of INRAE PEGASE laboratories.

REFERENCES

- Ross, J. W., Hale, B. J., Gabler, N. K., Rhoads, R. P., Keating, A. F., & Baumgard, L. H. (2015). Physiological consequences of heat stress in pigs. *Animal Production Science* 55: 1381-1390.
- Gonzalez-Rivas, P. A., Chauhan, S. S., Ha, M., Fegan, N., Dunshea, F. R., & Warner, R. D. (2022). Effects of heat stress on animal physiology, metabolism, and meat quality: A review. *Meat Science* 162: 108025.
- Lefaucheur, L., Le Dividich, J., Mouro, J., Monin, G., Ecolan, P., & Krauss, D. (1991). Influence of environmental temperature on growth, muscle and adipose tissue metabolism, and meat quality in swine. *Journal of Animal Science* 69: 2844-2854.
- Serviento, A. M., Lebret, B., & Renaudeau, D. (2020). Chronic prenatal heat stress alters growth, carcass composition, and physiological response of growing pigs subjected to postnatal heat stress. *Journal of Animal Science* 98: 13 p.
- Lebret B., Lhuisset S., Labussière E., & Louveau I., (2023). Combining pig genetic and feeding strategies improves the sensory, nutritional and technological quality of pork in the context of relocation of feed resources. *Meat Science* 197: 109074
- Johnson, J. S., Sanz Fernandez, M. V., Patience, J. F., Ross, J. W., Gabler, N. K., Lucy, M. C., Safranski, T. J., Rhoads R. P., & Baumgard, L. H. (2015). Effects of in utero heat stress on postnatal body composition in pigs: II. Finishing phase. *Journal of Animal Science* 93: 82-92.
- Ma, X., Wang, L., Shi, Z., Chen, W., Yang, X., Hu, Y., Zheng, C., & Jiang, Z. (2019). Mechanism of continuous high temperature affecting growth performance, meat quality, and muscle biochemical properties of finishing pigs. *Genes & Nutrition* 14: 23.