



HAL
open science

Identifying the most Efficient Detailed Trajectories toward Healthy Diets - A Graph-Based Analysis

Elie Perraud, Juhui Wang, Alison Dussiot, H el ene Fouillet, Fran ois Mariotti

► **To cite this version:**

Elie Perraud, Juhui Wang, Alison Dussiot, H el ene Fouillet, Fran ois Mariotti. Identifying the most Efficient Detailed Trajectories toward Healthy Diets - A Graph-Based Analysis. *Journal of Nutrition*, 2023, 153 (9), pp.2744-2752. 10.1016/j.tjnut.2023.07.007 . hal-04186238

HAL Id: hal-04186238

<https://hal.inrae.fr/hal-04186238>

Submitted on 25 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.

1 **Identifying the most efficient detailed trajectories toward healthy**
2 **diets – a graph-based analysis.**

Elie Perraud, Juhui Wang, Alison Dussiot, H  l  ne Fouillet, Fran  ois Mariotti

Universit   Paris-Saclay, AgroParisTech, INRAE, UMR PNCA, 91120, Palaiseau, France

All authors report no conflicts of interest.

Address correspondence to FM (email: francois.mariotti@agroparistech.fr).

Abbreviations used: CIQUAL: database of the French Centre d'Information sur la Qualit   des Aliments; DALYs: Disability-Adjusted Life Years; GDB: Global Burden of Disease; HR: Health Risk; INCA3: third individual and national study on food consumption survey; TMREL: Theoretical Minimum Risk Exposure Level

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data sharing: Data described in the manuscript, code book, and analytic code will be made available upon request pending application and approval.

Number of tables: 2

Number of figures: 3

Supplemental material: 12

3 **Abstract**

4 **Background:** Much effort has been devoted to defining healthy diets, which could lower the
5 burden of disease and provide targets for populations. However, these target diets are far
6 removed from current diets, so at best, the population is expected to move slowly along a
7 trajectory.

8 **Objective:** Our aim was to characterize the different possible trajectories toward a target diet
9 and identify the most efficient one for health in order to point out the first dietary changes
10 being the most urgent to implement.

11 **Methods:** Using graph theory, we have developed a new method to represent in a graph all
12 stepwise change trajectories toward a target healthy diet, with trajectories all avoiding risk of
13 nutrient deficiency. Then, we have identified and characterized the trajectory with the highest
14 value for long-term health. Observed male and female average diets are from the French
15 representative survey INCA3, and target diets were set using multicriteria optimization. The
16 best trajectories were found using the Dijkstra algorithm with the Health risk criteria based on
17 epidemiological data.

18 **Results:** Within ~2.6M diets in the graphs, we found optimal trajectories that were rather
19 similar for males and females regarding the most efficient changes in the first phase of the
20 pathways. In particular, we found that a one-step increase in the consumption of whole/semi-
21 refined bread (60 g) was the first step in all healthiest trajectories. In males, the subsequent
22 decrease in red meat was immediately preceded by increases in legumes.

23 **Conclusions:** We show simple practical dietary changes that can be prioritized along an
24 integral pathway that is the most efficient overall for health when transiting toward a distant
25 healthy diet. We put forward a new method to analyze dietary strategy for public health
26 transition and highlight the first critical steps to prioritize.

27 **Keywords:** graph theory, trajectory optimization, diet optimization, goal programming, long-
28 term morbidity and mortality

29 **Introduction**

30 Epidemiological studies about the associations between food group consumption or nutrient
31 and chronic diseases have shown that inadequate diets, too poor in healthy foods/nutrients and
32 too rich in unhealthy foods/nutrients intakes are one of the major causes of disability-adjusted
33 life years (DALYs) worldwide (1). A dietary transition is required and has been advocated by
34 all international institutes (2,3). From a dietary viewpoint and based on the expected decrease
35 in the burden of chronic diseases, changes to western diets notably include lower intake of red
36 meat, sugar-sweetened beverages, and refined grains, and higher intakes of whole grains,
37 fruits and vegetables, nuts and seeds, and other healthy plant sources (4). At the nutrient level,
38 it implies lower intakes of sodium, sugar, and SFA and higher intake of PUFA, EPA+DHA,
39 and fiber (5).

40 Much effort has been devoted to defining healthy diets that make the final objective for
41 populations. Food pattern modeling has identified many healthy diets, notably using
42 optimization models (6,7). By construction, such diets are nutritionally adequate and
43 associated with a lower risk of chronic disease. Although often including limitations of the
44 distance to observed diet, to make feasible objectives, these target diets are much departed
45 from current diets. This can be explained partly by current western diets having nutritional
46 adequacy risks for several nutrients. Thus, modifying the diet to reach the adequacy level of
47 each nutrient and adherence to a healthy dietary pattern is indeed a profound modification of
48 the diet.

49 A sudden complete change of the diet is not attainable, particularly at the population level,
50 and at best, the population is expected to transition along a trajectory toward a target diet at a
51 rather slow pace (8,9). In reality, the trajectory of change that a population is starting should
52 greatly impact its nutritional health status and, thereby, the cost for society (10). However,
53 there has been little research to characterize such trajectories, particularly to identify those
54 most efficient for health during the long and gradual process toward the target diet. Yet, such
55 data are needed to specify the pathways of changes and point out the first steps within the
56 pathways that are likely to be the most critical and urgent to implement.

57 Showing the first steps to be taken is also probably an important lever to sustained changes.
58 Significant barriers to change are the lack of information (11) and the efforts needed to
59 change eating routines (12). For example, even if most of the population knows the dietary
60 guidelines, there are applied only by a small percentage (13). The very small changes toward

61 healthy diets in the last years (14,15) call for revising strategies in public health nutrition.
62 Focusing on more precise and effective information may be one of the effective options to be
63 developed. However, for this purpose, we need more information about how small changes,
64 such as an increase or a decrease in portion size, must be coordinated and prioritized to make
65 the most efficient trajectory of changes. Such trajectories should be optimal for long-term
66 health (reducing, at best, the risk of chronic disease) along the trajectory, but also, it should
67 not include a diet with an increased risk of nutrient deficiency. Both the former and the later
68 points are crucial as changes at a population level are slow, and a transitory diet toward the
69 targeted one needs to be safe. Some of the population could stay with this diet for a long time.
70 For example, between 2006 and 2014 in France, daily meat consumption decreased by < 3g,
71 and daily vegetable consumption increased by <9g (14,15).

72 Many works have aimed to analyze dietary changes and transitions. Two types of methods
73 have been used to identify diet transitions. A first type of study used optimization under
74 constraint as a method. This method enables finding the best diets in the context of an
75 increasingly binding constraint. For example, this method has been used to find the best diets
76 while reducing meat intake (16). However, this method does not produce real trajectories but
77 a succession of milestones. Thus, the trajectory that would consist of going through all these
78 stages is not defined as the most efficient one overall. A second type of study used gradual
79 changes (17,18). According to an incremental algorithm, this method leads to the selection of
80 the best change that could be applied to a diet at each step, but this method is agnostic to the
81 overall trajectory. Accordingly, such trajectories include many local optimums and do not
82 reach any target diet. Thus, no methods have considered a series of simple changes and
83 analyzed complete transition pathways. Dissecting all possible changes in a distant dietary
84 transition requires a systematic approach with a conceptual framework and computer-
85 intensive methods. Here, we report on research that used a new method for analyzing diet
86 trajectories using the graph theory conceptualization. In this study, we identified and
87 characterized the best and safe trajectories toward a target diet using a graph theory and
88 highlight critical simple dietary changes to be prioritized.

89

90 **Methods**

91 1. Studied population and dietary data

92 This study used data from the third individual and national study on food consumption survey
93 (INCA3) performed in France between 2014 and 2015 (19,20).

94 The participant selection process for this study involved a three-stage random sampling
95 design conducted by the National Institute of Statistics and Economic Studies (INSEE), based
96 on the 2011 annual population census. Ineligible individuals included those who were
97 institutionalized, those planning to move out of their homes within the next 2 months, and
98 those who were unable to be interviewed. Further description of the method is in (19).

99 Participants identified as under-reporters were excluded using the basal metabolic rate as
100 estimated by the Henry equation (21), using the cut-off values recommended by Black (22).
101 Older adults (above 54 years old for female and 64 years old for male (23)) were excluded as
102 their risk of long-term morbidity and mortality differed from younger adults, and we wanted
103 to keep a homogeneity in our population. The final sample contained 1125 adults
104 (Supplemental Figure 1).

105 Dietary data were collected by professional investigators assisted by dietary software from
106 three unplanned, non-consecutive, 24h dietary recalls spread over a three-week period (two
107 during the week and one at the weekend). Portion sizes were estimated using validated
108 photographs (19). The nutrient content values of the food were extracted from the 2016
109 database of the French Centre d'Information sur la Qualité des Aliments (CIQUAL) (24).
110 Mixed foods were broken down into ingredients and then gathered into 33 food groups. These
111 food groups were derived from the 45 food groups established by Dussiot et al. (6). The
112 "Fruits" group encompasses fresh fruit, dried fruits, and processed fruits. The "Whole meal
113 and semi-refined bread and bakery products" group comprises whole grain and semi-refined
114 grain bread and bakery products. The "Other refined starches" group includes rice, pasta,
115 other starches, potatoes, and other tubers. The "Processed starch-based products" group
116 consists of starch-based products, sweet/fat processed products, and salt/fat processed starch
117 products. The "Red meat" group encompasses beef, veal, pork, and other meats, and offal.
118 The "Other fish" group includes other fish, mollusks, and crustaceans. The "Fresh dairy
119 products" group includes fresh natural dairy products and fresh sweetened dairy products. The
120 "Animal fats and assimilated fats" group comprises animal fats, assimilated fats, butter, and
121 low-fat butter. The "Sugar-sweetened beverages, including fruit juices" group includes
122 sweetened soda-type drinks and fruit juices. The "Condiments" group encompasses
123 condiments, aromatic herbs, and spices except salt. For a detailed description of each food

124 group, please refer to the Supplementary Material in Dussiot et al (6). The nutrient content of
125 each food group was calculated as the mean nutrient content of food items constituting the
126 group weighted by their mean intake, as previously described (6).

127 A sex-specific average diet was calculated using the data from the 564 males and the 561
128 females (Table 1) and used as the initial diet for defining the trajectories.

129 2. Target diet

130 The target diets were identified for each sex, according to a multi-criteria optimization
131 approach previously developed by our group (6,25). Using this approach, by construction,
132 target diets ensure adequate nutrient intakes and are found within current consumption limits
133 (5th and 95th percentiles), and they combine minimal long-term health risks and minimal
134 departure from the observed diet. The method is described in full in Supplemental method 1.
135 Target diets are described in Table 1.

136 3. Graphs theory and construction

137 A graph is a pair $G = (N, E)$ of nodes (N) and edges (E) (26). In our model, nodes represent
138 daily consumed diets, and edges correspond to a modification of one portion step. If two
139 nodes are linked with an edge, it is possible to transition from one diet to the other.

140 As explained above, all foods consumed in the INCA3 survey have been classified into 33
141 food groups according to their nutritional characteristics. A node (diet) is thus a vector with
142 33 variables, one for each food group. To simplify the graph, the quantity eaten for each food
143 group was rounded to the number of portions of this group.

144 For each food group, we decided on portion size; it represents the food quantity of a mean
145 food portion eaten in INCA3. The portion sizes were based on the distribution of food
146 consumption and known portioned food. For example, yogurts are portioned in France and
147 weigh 125g. Therefore, this portion size was used for the group "fresh dairy product."

148 We also adapted the portion sizes, so that food groups with a similar role in the dietary pattern
149 have the same portion size. For example, "poultry" and "red meat" have the same portion
150 size. Supplemental Table 1 presents the portion size for each of the food groups.

151 4. Method of graph creation and use

152 Graphs were constructed from both the initial and objective diet nodes. All nodes were
153 created by proximity; this means adding nodes one portion step away from the previous one;
154 until the graph was connected. A connected graph is a graph with all nodes having a path to

155 all other nodes of the graph. Then another extension was added to ensure the graph had no
156 bottleneck due to the way it was created. It was difficult to increase the graph size again as the
157 number of nodes in the graph, especially the one for male, was above a million nodes. In
158 order to reduce the number of nodes, constraints and limits were added during the extension
159 process. The nodes not following these constraints were not added to the graph.

160 Each node had to respect the following constraints. The number of steps of portion size for
161 each food group should be between 0 and the maximum step of portion size as observed in the
162 INCA3 population. The energy of the diet of the node should be between 80% and 120% of
163 the energy brought by the initial diet. The SecDiet of the node, a score of nutrient security
164 (27) (Supplemental Method 2), should be >99% of the SecDiet of the initial diet. The
165 percentage of macronutrients in each node should be between the minimum and maximum
166 macronutrient intake in the population. The following constraints were added based on French
167 dietary guidelines. In cases where the daily consumption of nuts falls below 40g or the
168 consumption of fruits and vegetables falls below 400g, it is considered below the minimum
169 threshold and cannot be further reduced. Similarly, if the consumption of processed meat
170 exceeds 150g or the consumption of red meat exceeds 500g per day, it is considered above the
171 maximum threshold and cannot be further increased.

172 5. Trajectories analysis

173 The optimized trajectory was found using Dijkstra algorithm (28). The trajectories were
174 optimized to minimize the long-term health risk all along the trajectory using the Health Risk
175 (HR) criterion (Supplemental Method 1). The trajectory is thus optimized, so the sum of the
176 HR of all trajectory nodes reaches its minimum. Suboptimal trajectories were also identified.

177 The HR criterion was set to target the dietary recommendations from the GBD based on
178 epidemiological studies about the associations between the consumption of different food
179 groups and the risk of chronic diseases (1,29). HR represents the combined and normalized
180 distance to the theoretical minimum-risk exposure levels (TMREL) for three unhealthy food
181 groups (red meat, processed meat and sweetened beverages), six healthy food groups (whole
182 grain products, fruits, vegetables, legumes, nuts and seeds, and milk) food groups, four
183 healthy nutrients (fiber, calcium, EPA/DHA, PUFA) and one unhealthy nutrient (sodium),
184 weighted by the importance of reaching each TMREL value in terms of its relative impact on
185 DALYs in France. HR can range from 0 to 1, from the situation where the diet meets all the

186 TMREL values (minimum risk) to the situation where it is farthest from them (maximal risk).
187 A precise description of the HR criterion can be found in Supplemental Method 1.

188 The percentage of the Population Reference Intake for several nutrients was calculated to
189 describe the trajectories.

190 All methods were implemented on python 3.8.3 (30). The networkX package version 2.4 was
191 used to implement the graph and the Dijkstra algorithm (31). The community package version
192 0.15 was used to implement the Louvain algorithm (32).

193

194 **Results**

195 **Table 2** presents the characteristics of the graphs for male and the ones for female. The
196 number of nodes between the initial and target nodes was 24 steps for male and 20 steps for
197 female. Due to this increased number of steps between the initial and target nodes, the graph
198 for male was bigger than for female. The mean number of degrees (*i.e.*, edges per node)
199 representing the density of the graph was slightly higher for male than for female.
200 Supplemental Table 2 describe the lifestyle and sociodemographic characteristics of the
201 included participants and Supplemental Table 3 describe the daily energy and nutrient intake
202 of the sample.

203 **Figure 1** describes the evolution of the health risk (HR) along the trajectory as a percentage
204 of the difference between the worst HR value (of the observed initial diet) and the best HR
205 value (which is that of the final target diet). As the trajectory was optimized to minimize the
206 health risk (HR), the first steps were the most HR-efficient, *i.e.* those with the highest HR
207 yield, which then progressively decreased with each step. The HR criteria decreased strongly
208 (around -80% of its initial, maximal value) in the first eight steps for both sexes.

209 **Figure 2** and **Figure 3** describe the optimal diet trajectory for male and female respectively.
210 For both male and female with quite distinct target diets, the first step, the most efficient one,
211 was the increase of whole and semi-refined bread, with an increase of one step for female to
212 reach the number of portions in their target diet and an increase of two steps for male (step 1
213 and step 2), which also reached the number of portions in their target diet.

214 After these first increases in whole and semi-refined bread, the fruit category was increased.
215 Fruits were increased for woman in step two and for male in step three. The increase in
216 vegetables occurred at step five for female and step seven for male. For male, between the

217 increases in fruit and vegetable there was a substitution of red meat by legumes in three
218 successive steps (two increases in legumes followed by one decrease in red meat). After these
219 substitutions and increases, bread and refined bakery products were decreased for male. For
220 female, between the increases in fruit and vegetable were the milk increase and the bread and
221 refined bakery products decrease. Supplemental Figures 2 and 3 describe the nutrient intakes
222 along the steps of the trajectory.


223 For each sex, fifty suboptimal trajectories were analyzed and were found similar to the
224 optimal one for the first 17 steps for male and the first 12 steps for female (i.e., only the
225 following less HR-effective steps differed). Supplemental Figures 4 and 5 and Supplemental
226 Tables 4 and 5 describe the other choices available for the different steps of the trajectories.

227

228 *Figure 1 Decrease in the HR criterion along the diet trajectories from the observed average diet to the*
 229 *set target diet of male (left panel) or female (right panel).*


230 *Changes in the HR criterion is shown as a percentage of the difference between the worst HR value (of*
 231 *the observed initial diet) and the best HR value (which is that of the final target diet).*

232 *HR: Health Risk*



233
 234


235 *Figure 2 Best trajectory identified in male based on the Health Risk Criterion, which uses estimates*
 236 *from the Global Burden of Disease.*



237

238

239 Figure 3 *Best trajectory identified in female based on the Health Risk Criterion, which uses estimates*
 240 *from the Global Burden of Disease.*



241

242

243 *Table 1 Food group consumptions (g/d) by sex in the initial and target diets*

	Initial diet		Target diet	
	Male	Female	Male	Woman
Vegetables	176	160	400	368
Fruits	142	124	419	316
Nuts and seeds	3	2	11	13
Bread and refined bakery products	168	115	27	11
Whole meal and semi-refined bread and bakery products	11	15	99	60
Other refined starches	177	121	193	139
Other whole and semi-refined starches	4	4	5	8
Processed starch-based products	26	21	0	13
Legumes	13	6	86	19
Poultry	30	31	46	17
Red Meat	79	42	54	31
Processed meat	50	30	0	0
Oily fish	8	6	26	21
Other fish	27	19	1	24
Eggs and egg-based dishes	14	14	11	12
Milk	84	75	190	126
Fresh dairy products	81	81	28	74
Sweet milky desserts	19	16	0	6
Cheese	49	36	48	13
Animal fats and assimilated fats	11	10	4	0

Vegetable fats rich in α -linoleic acid	0	0	16	5
Vegetable fats low in α -linoleic acid	12	10	15	4
Sauces and fresh creams	35	32	35	100
Sweet products	103	83	103	98
Drinking water	1007	929	878	1108
Sugar-sweetened beverages, including fruit juices	221	208	0	75
Hot drinks	494	507	379	507
Salt	1	1	0	0
Condiments	6	4	13	5
Soups and bouillons	76	79	0	53
Substitutes for animal products	3	5	29	28
Other foods	4	2	4	2
Alcoholic drinks	216	59	0	3

244

245

246 *Table 2 Description of the graph and of the initial and target nodes*

Sex		Male	Female
Number of nodes in the graph		2 344 253	281 257
Number of nodes between the initial and target nodes		24	20
mean		18.1	15.7
SD		3.7	3.0
min		2	2
max		24	19

247

248 **Discussion**

249 Using graph analysis as a new method applied to dietary transitions, we could identify and
250 characterize trajectories of dietary changes that are optimal for long-term health while
251 avoiding nutrient deficiency risk.

252 This work was based on the premise that identifying detailed trajectories could help decipher
253 the most important steps in a dietary transition, mainly by analyzing the pattern of changes
254 along the trajectory and highlighting the first steps being the most effective in mitigating
255 health risks. In this regard, our findings point out critical changes to transition toward a
256 healthy diet.

257 1. Trajectories analysis

258 Firstly, it is important to note that the graph generated for males is nearly ten times larger than
259 the one for females. This discrepancy is mostly related to the different number of steps
260 required to transition from the initial to the target diet. As the gap between observed and
261 target diets increases, the number of nodes in the graph expands exponentially. This high gap
262 is a result of the poorer quality of observed diet in males. However, despite this difference in
263 size, the graph structure remains similar, as the average number of neighbors per node
264 remains similar.

265 With the first six steps of the trajectories, males and females HR scores are already improved
266 by at least two third of the maximum improvement. It confirms that this method identifies
267 very efficient steps, especially in the first steps of the trajectories.

268 The first dietary change in all dietary trajectories was an increase in whole/semi-refined
269 bread. This result aligns with the importance of whole grains in health, as shown by the GBD
270 studies (1,33). In France, given that a diet low in whole grain is the highest dietary risk factor
271 in both males and females (34), it is not surprising that an increase in whole grain products is
272 the first step. The importance of the increase in whole grain is clearly shown by the presence
273 of two successive increases in whole bread in the first two steps of the trajectory in males.
274 When whole grains are concerned, attention should be paid to the residual pesticide levels
275 related to agricultural practices. Whole grains products are, on average, higher in pesticides
276 than refined grains products (35), which might be considered as potentially opposing the
277 benefits of whole and semi-refined bread. However, this potential drawback should not be
278 expected to call into question the final benefit of whole and semi-refined bread since, as

279 accounted for in the GBD data, the DALYs gain is based on epidemiological data, with
280 consumption mostly coming from conventional agriculture (36).

281

282 Important steps in the dietary trajectory that we identified were increases in fruit and, to a
283 lesser extent, vegetables, which is again in line with the relative importance of fruit and
284 vegetables in the HR criteria. Importantly, these increases are early and that of fruit
285 immediately follows the increase in whole/semi-refined bread.

286 A decrease in red meat in the first phase of the trajectory was only found in male, at step #6,
287 despite the importance of red meat in the HR function. This result can be explained mainly by
288 the importance of the food groups being increased before, which are consumed in very small
289 amounts compared to the recommended value. It should be noted that red meat could have
290 been decreased at each of the first steps of the trajectories because it was not limited by a
291 constraint related to nutritional security. Thus, its reduction at step #6 only comes from the
292 optimization criteria. For females, no decrease was needed since the target diet was not very
293 low in red meat and, when expressed as portion steps, the initial and the target values had
294 similar amounts. On the other hand, since meat is dense in protein, bioavailable iron, and
295 vitamin B12 (37), it has often been proposed that reducing meat consumption may increase
296 the inadequacy of some nutrients. In a recent work, we showed that the critical nutrients when
297 decreasing meat in healthy diets are bioavailable iron, zinc, and vitamin A (16). However, we
298 found no increase in the risk of inadequacy for all those nutrients when reducing meat within
299 this trajectory (Supplemental Figure 2.N, 2.U and 2.A and Supplemental Figure 3.N, 3.U and
300 3.A). This is in line with previous studies showing that protein quantity is not an important
301 risk in the transition in western countries to a plant-based diet (38,39). The risk of insufficient
302 intake of vitamin B12 also appears to depend on the overall decrease in foods of animal origin
303 rather than meat in isolation (40).

304 Furthermore, as for other trajectory steps, decreasing red meat was done without increasing
305 the risk of deficiency since the graph was trimmed to avoid diets featuring any increase in the
306 risk of overt deficiency, as assessed using the SecDiet metric. It should be noted that for
307 female all along the trajectory and for male after step#18, bioavailable iron is under the
308 reference value (Supplemental Figure 3.N). However, the intake remains above the lower
309 threshold value ensuring <5% deficiency prevalence. Thus, iron-deficiency anemia in the
310 population would stay around today's level. We did not impose that the iron reference value

311 was met during the trajectory and in the target diet. This choice was made as the current diet
312 does not meet this high reference value, in particular for female, and previous work has
313 demonstrated that the current value regarding bioavailable iron is over constraining when
314 intending to model healthier diets (6). It is striking that increases in legumes preceded the
315 decrease in red meat consumption. As we have previously discussed (41), this shows the
316 importance of the transition between animal to plant protein. The importance of legumes in
317 this pathway may mean that in diet changes consisting in lowering red meat consumption,
318 some pathways not featuring legumes are discarded because of nutritional risk; since legumes
319 are rich in nutrients that may be limiting in this case.

320 2. Nutrient-secured healthiest trajectories and possible implication for guidelines

321 In this respect, the trajectory we have identified is the result of a combination of the long-term
322 health value of the portioned changes, the position of the initial and target diets, and the strict
323 avoidance of pathways leading to diets that would be nutritionally unsecured. This latter
324 characteristic is particularly important because while the target diet is fully nutritionally
325 adequate, pathways of changes leading to that diet may include an increase in the risk of
326 nutrient deficiency. Therefore, not all changes were available at all steps and the optimization
327 consisted in choosing best choice at each step (Supplemental Figures 4 and 5 and
328 Supplemental Tables 4 and 5) to make the best trajectory overall, i.e. the most efficient for
329 health.

330 Our findings of critical steps within the best overall pathways toward a healthier diet may
331 have implications for more precise dietary guidelines. Pointing out simple changes to be
332 prioritized should be easier for the general audience and, therefore, may help food-based
333 dietary guidelines to be more effective. For instance, although increasing whole grain
334 products is included in dietary guidelines, it is not the recommendation that is the most
335 widely known and applied (42) compared to the well-known recommendation regarding fruit
336 and vegetables (13). Furthermore, pointing out a more precise change, such as here focusing
337 on whole-meal bread, should give more applicable advice while maintaining a strong
338 scientific rationale. Thus, prioritizing some dietary changes could change how public health
339 nutrition programs are put forward. For whole grain in general, and whole bread in
340 particular, which can be eaten similarly to white bread, the option may be even easier to
341 apply. In this regard, it should be conceded that because the graph-based method consists of a
342 series of increases/decreases, our results identify isolated increases in whole-meal bread, not

343 swaps for white bread. However, an increase in one portion step of whole-meal bread consists
344 in adding 60g of whole-meal bread, and this change can be conceived and implemented apart
345 from a reduction in white bread. Furthermore, some directly successive changes might be
346 translated as swaps in the recommendations, provided it makes sense from a practical
347 viewpoint. We think it is the case for the decrease in red meat, which is immediately preceded
348 by two successive increases in legumes and can be easily operationalized as replacing 40g red
349 meat with 80g legumes. Of course, our results do not mean that these changes do not require a
350 large effort from the population. Indeed, in western countries, a large part of the population
351 does not know how to cook legumes (43), and culturally speaking, meat is still an important
352 part of the meal. Lastly, insisting on the diet changes that appear first in the identified
353 trajectories could stand as a rationale way to prioritize changes, while keeping those changes
354 integrated within a full trajectory. Defining ordered step-by-step changes could concur with
355 setting graded tasks, which is part of the behavioral changes techniques (44,45).

356 3. Graph analysis for identifying optimal dietary transition toward target diets

357 Graph analysis methods are well known for the analysis of road networks (14), social
358 networks (15), or protein structures (16). However, to our knowledge, it has never been used
359 to analyze diet transition. Other methodological approaches have been used to analyze diet
360 transitions. While optimization studies are often used to identify target diets that could back
361 food-based dietary guidelines (46), optimization has also been used to identify intermediate
362 diets. For example, we recently identified a series of diets with progressively proportionally
363 reduced amounts of meat (16) and this could be taken as showing a trajectory of change made
364 of intermediate diets towards the lowest level of meat possible. However, this approach is
365 different since it shows a series of healthy diets after forced scenarios of changes (here,
366 reduction in meat consumption) and not the best overall healthy pathway toward the final
367 target. These diets may not be part of the most efficient pathway. Furthermore, this serial
368 optimization method shows diets that are not directly connected at the finest level, i.e., they
369 are not demonstrated as being connected to one another with changes pathways with no
370 nutritional risks, and diets can be very far from each other. Few methods have considered
371 trajectories as a series of very fine level changes. Verger et al. (18) and De Gavelle et al. (17)
372 have used such an approach made by step-by-step changes at portion size level. However, this
373 method also has some limits. If a local optimum is reached, it is impossible to continue to a
374 better one. Also, the method finds the best next step at each point and not the best path

375 overall. A lesser improvement at one step could unlock a diet path that is in fact much more
376 efficient overall.

377 In contrast, with the present graph-based method, we can optimize all along the trajectory,
378 i.e., find the best series of diets taken together considering their health value. Another central
379 advantage of the method is that we can apply any set of constraints on all nodes included in
380 the graph. Here, the dietary and nutritional constraints that we chose were embedded in the
381 graph, so they applied to all trajectories. For example, in all the graph during its construction,
382 the SecDiet could not be lower than that of the observed diet, and this precluded that the
383 identified trajectories could include a point diet with a nutritional security risk.

384 One major feature of this method is that it identifies a trajectory from the observed diet to a
385 target diet, but it requires that the target diet should be defined with another method. This
386 study used multicriteria optimization under constraints to define the target diet. Another
387 parametrization of the model (in particular for less conservative weighting of diet deviation)
388 or another method that would change the target diet may lead to a different graph and a
389 different optimal trajectory than the one we identified here. One of the limits of this method is
390 the restraint on the target diet in terms of the total distance from the observed diet because the
391 target diet should be close enough to the initial one, given the fine level of resolution of the
392 graph if the target is not close enough, its dimension will be too large. Being close enough
393 means having less than 25 number of portions steps (i.e., nodes) between the node
394 representing the observed diet and the node representing the target diet. Indeed, because of
395 how the graph is created and even with the constraints added during the construction, the
396 number of nodes in the graph is exponential to the number of portions steps between the
397 initial and target diets, and the graph for male is already composed of 2.3 million nodes. Thus,
398 if choosing a target diet that is farther from the initial diet, then the resolution of the
399 representation of the diet with the graph should be lower, which means choosing a lower
400 number of larger food groups or higher steps in portion sizes of each food. It should also be
401 noted that this methodological approach strongly depends on the optimization criteria. In this
402 study, the HR criteria we chose represent the diet proximity to reference values (TMREL) that
403 are those of the lower risk for chronic diseases, weighted by the attributable fraction of the
404 risk factor. Thus, optimizing this score aims to identify the trajectory that would lead to the
405 lower risk of long-term morbidity and mortality, therefore being the integral trajectory that is
406 the more efficient for long-term health. The improvement of health risk criteria along the
407 trajectory is not uniform, and the best trajectory that has been identified is the best overall,

408 accounting together for the 14 components of the criteria and taking into account the whole
409 trajectory till the end. Therefore, some components that align less with the others in the
410 optimization may be only reduced in the second part of the trajectory and may even be
411 transiently deteriorated. This was the case for sodium. Reaching sodium reference value has
412 been found constraining in modelling studies (6), and changes within food categories, which
413 were not addressed here, are helpful to lower sodium intake (18,47). The advantage of the
414 method is to weight all components in the health criteria to find the overall best change for the
415 general population, but it may be less appropriate for some specific populations with different
416 risk factors. However, the method can be adapted to account for different criteria in a more
417 specific population. For instance, in people with hypertension, sodium intake could be the
418 optimization criteria either alone or combined with others. The method presented here is
419 indeed versatile, with the possibility to add constraints directly into the nodes of the graph. In
420 this example, a maximum level of sodium could be added if the goal is to identify trajectories
421 strictly avoiding transient increase in sodium intake. The same method can be used with other
422 target diets and optimization criteria. For example, another study could find optimal pathways
423 to adherence to the dietary guidelines or optimal pathways for reducing an environmental
424 pressure score or a combination of a nutritional and environmental score. This method could
425 also be used to identify individuals' trajectories of diet transition. Indeed, the only change
426 necessary is the modification of the initial diet. Other changes, like adapting the target diet,
427 could also be made depending on the question. However, it should be noted that generating
428 graphs for each individual and identifying the optimal trajectories requires considerable
429 computational time and can, therefore, only be carried out for a restricted number of
430 individuals. Moreover, as they progress, the trajectories of different individuals will probably
431 rapidly converge to a similar path. Therefore, rather than at the individual level, a set of
432 trajectories may easily be drawn for subgroups of the population and give enough information
433 about the variability of the first steps according to the characteristics of these groups, such as
434 their initial dietary pattern, as identified in a population survey. Defining the individual
435 trajectory of a given individual to provide tailored information is also possible on this basis
436 but remains a very different field of application.

437 It should also be noted that this method does not include all the factors that need to be
438 taken into account if the results are to be readily translated into dietary recommendations.
439 Indeed, this work focuses on nutrition/health impact of diet trajectories and does not take
440 into account to what extent changes in the consumption of a food group are accessible to

441 the people, depending on their characteristics. For example, the affordability, accessibility,
442 or individual preferences could be factors modifying the implementation of the changes that
443 we have identified here.

444 Moreover, this work only considered nutrient intake coming from food groups. In a
445 population with a high use of dietary supplement, it should be useful to identify whether
446 specific solutions could be found while considering nutrient intakes from supplements.
447 However, the percentage of individuals taking supplements in the study population was
448 relatively low (28.5%), and supplements were not taken regularly (48). Importantly, using of
449 supplements has been found to have no impact on the risk of insufficient intakes in the
450 population (49). The accessibility of supplements is not the same for all the population, and
451 supplements may be more beneficial in lower socioeconomic status individuals who eat
452 diets of lower quality and have higher risks of insufficient nutrient intakes (27). However, in
453 future studies, it could be interesting to evaluate what could be the effect of
454 supplementation on the trajectories and if better solutions could be found if an effective
455 supplementation strategy was set. Likewise, the effect of fortifying specific foods or food
456 staples may also be the subject of further studies.

457

458 One of the other advantages of this method is the possible use of multiple variables to
459 characterize the nodes (diets). These variables can be used as constraints to invalidate some
460 pathways. These variables can also be used to describe the trajectory with respect to other
461 dimensions of the diets.

462 **Conclusion**

463 Methods based on graph theory can be used to draw all possible nutrient-secure pathways of
464 changes to reach a reference diet and identify the detailed integral pathway that is optimal
465 overall for improving long-term health. Here, we could notably show that an increase in the
466 portion size of whole/semi-refined bread is a critical first step in this pathway before increases
467 in fruit and vegetables. It can show how small and realistic changes should be prioritized
468 while keeping with the final objective. Such analysis models may be useful for refining food-
469 based dietary guidelines on practical grounds.

470

471 **Acknowledgements**

The authors' responsibilities were as follows – EP, JW and FM designed research; EP was primarily responsible for preprocessing the data, conducting trajectory optimization and interpreting the results, with advisory inputs from JW, AD, HF and FM; AD and HF provided some data and tools; EP drafted the manuscript; EP and FM wrote the manuscript with critical comments from JW, AD, and HF. All authors approved the final manuscript. The authors report no conflicts of interest.

472

473 **References**

- 474 1. Murray CJL, Aravkin AY, Zheng P, Abbafati C, Abbas KM, Abbasi-Kangevari
475 M, Abd-Allah F, Abdelalim A, Abdollahi M, Abdollahpour I, et al. Global burden
476 of 87 risk factors in 204 countries and territories, 1990–2019: a systematic
477 analysis for the Global Burden of Disease Study 2019. *The Lancet*
478 2020;396:1223–49.
- 479 2. Springmann M, Clark M, Mason-D’Croz D, Wiebe K, Bodirsky BL, Lassaletta L,
480 de Vries W, Vermeulen SJ, Herrero M, Carlson KM, et al. Options for keeping
481 the food system within environmental limits. *Nature* 2018;562:519–25.
- 482 3. Sustainable healthy diets: guiding principles [Internet]. World Health
483 Organization (WHO), Food and Agriculture Organization of the United Nations
484 (FAO); 2019 Oct p. 37. Available from: [https://www.who.int/publications-detail-
485 redirect/9789241516648](https://www.who.int/publications-detail-redirect/9789241516648)
- 486 4. The US Burden of Disease Collaborators. The State of US Health, 1990-2016:
487 Burden of Diseases, Injuries, and Risk Factors Among US States. *JAMA*
488 2018;319:1444–72.
- 489 5. The French Agency for Food Environmental and Occupational Health & Safety
490 (ANSES). Avis de l’Agence nationale de sécurité sanitaire de l’alimentation, de
491 l’environnement et du travail relatif à « Actualisation des références
492 nutritionnelles françaises en vitamines et minéraux » (saisine n°2018-SA-0238,
493 saisine liée n°2012-SA-0103, 2021). 2021; Available from:
494 <http://www.anses.fr/fr/system/files/NUT2018SA0238Ra.pdf>
- 495 6. Dussiot A, Fouillet H, Wang J, Salomé M, Huneau J-F, Kesse-Guyot E, Mariotti
496 F. Modeled healthy eating patterns are largely constrained by currently estimated
497 requirements for bioavailable iron and zinc—a diet optimization study in French
498 adults. *Am J Clin Nutr* 2022;115:958–69.
- 499 7. Gazan R, Brouzes CMC, Vieux F, Maillot M, Lluch A, Darmon N. Mathematical
500 Optimization to Explore Tomorrow’s Sustainable Diets: A Narrative Review.
501 *Adv Nutr Oxford Academic*; 2018;9:602–16.
- 502 8. Stewart C, Piernas C, Cook B, Jebb SA. Trends in UK meat consumption:
503 analysis of data from years 1–11 (2008–09 to 2018–19) of the National Diet and
504 Nutrition Survey rolling programme. *Lancet Planet Health* 2021;5:e699–708.
- 505 9. Deliens T, Mullie P, Clarys P. Plant-based dietary patterns in Flemish adults: a
506 10-year trend analysis. *Eur J Nutr* 2022;61:561–5.
- 507 10. World Health Organization. Regional Office for Europe, Policies EO on HS and,
508 Candari CJ, Cylus J, Nolte E. Assessing the economic costs of unhealthy diets
509 and low physical activity: an evidence review and proposed framework

- 510 [Internet]. World Health Organization. Regional Office for Europe; 2017 [cited
511 2022 Jul 27]. 90 p. Available from: <https://apps.who.int/iris/handle/10665/326302>
- 512 11. Lea EJ, Crawford D, Worsley A. Public views of the benefits and barriers to the
513 consumption of a plant-based diet. *Eur J Clin Nutr Nature Publishing Group*;
514 2006;60:828–37.
- 515 12. Pohjolainen P, Vinnari M, Jokinen P. Consumers’ perceived barriers to following
516 a plant-based diet. *Br Food J Emerald Group Publishing Limited*;
517 2015;117:1150–67.
- 518 13. Escalon H, Beck F, Bossard C. Associations entre la connaissance des
519 recommandations du Programme national nutrition santé et les comportements en
520 matière d’alimentation et d’activité physique. *Rev DÉpidémiologie Santé
521 Publique* 2013;61:37–47.
- 522 14. ANSES. the Third Individual and National Survey on Food Consumption (INCA3
523 Survey) [Internet]. 2017 [cited 2022 Jan 24]. Available from:
524 <https://www.anses.fr/en/system/files/NUT2014SA0234EN.pdf>
- 525 15. Agence Française de Sécurité Sanitaire des Aliments. Synthèse de l’étude
526 individuelle nationale des consommations alimentaires 2:(INCA 2) 2006-2007
527 [Internet]. 2009 [cited 2022 Jul 27]. Available from:
528 <https://www.anses.fr/fr/system/files/PASER-Ra-INCA2.pdf>
- 529 16. Dussiot A, Fouillet H, Perraud E, Salomé M, Huneau J-F, Kesse-Guyot E,
530 Mariotti F. Nutritional issues and dietary levers during gradual meat reduction –
531 A sequential diet optimization study to achieve progressively healthier diets. *Clin
532 Nutr* 2022;41:2597–606.
- 533 17. de Gavelle E, Leroy P, Perrimon M, Huneau J-F, Sirot V, Orset C, Fouillet H,
534 Soler L-G, Mariotti F. Modeled gradual changes in protein intake to increase
535 nutrient adequacy lead to greater sustainability when systematically targeting an
536 increase in the share of plant protein. *Clim Change* 2020;161:129–49.
- 537 18. Verger EO, Holmes BA, Huneau JF, Mariotti F. Simple Changes within Dietary
538 Subgroups Can Rapidly Improve the Nutrient Adequacy of the Diet of French
539 Adults. *J Nutr* 2014;144:929–36.
- 540 19. Dubuisson C, Dufour A, Carrillo S, Drouillet-Pinard P, Havard S, Volatier J-L.
541 The Third French Individual and National Food Consumption (INCA3) Survey
542 2014–2015: method, design and participation rate in the framework of a European
543 harmonization process. *Public Health Nutr* 2019;22:584–600.
- 544 20. French Agency for Food, Environmental and Occupational Health Safety
545 (ANSES). Opinion of the French Agency for Food, Environmental and
546 Occupational Health & Safety on “the Third Individual and National Survey on

- 547 Food Consumption (INCA3 survey)” [Internet]. 2017 [cited 2021 Jul 8].
548 Available from: <https://www.anses.fr/en/system/files/NUT2014SA0234EN.pdf>
- 549 21. Henry C. Basal metabolic rate studies in humans: measurement and development
550 of new equations. *Public Health Nutr* 2005;8:1133–52.
- 551 22. Black AE. Critical evaluation of energy intake using the Goldberg cut-off for
552 energy intake:basal metabolic rate. A practical guide to its calculation, use and
553 limitations. *Int J Obes* 2000;24:1119–30.
- 554 23. ANSES. Actualisation des repères du PNNS : élaboration des références
555 nutritionnelles. Maisons-Alfort, France: French Agency for Food, Environmental
556 and Occupational Health Safety (Anses). [Internet]. 2016 [cited 2022 May 13].
557 Available from: <https://www.anses.fr/fr/system/files/NUT2012SA0103Ra-2.pdf>
- 558 24. ANSES. Table de composition nutritionnelle des aliments CIQUAL [Internet].
559 2016 [cited 2021 Jun 22]. Available from: <https://ciqual.anses.fr/#>
- 560 25. Fouillet H, Dussiot A, Perraud E, Wang J, Huneau J-F, Kesse-Guyot E, Mariotti
561 F. Plant to animal protein ratio in the diet: nutrient adequacy, long-term health
562 and environmental pressure [Internet]. medRxiv; 2022 [cited 2022 Sep 14]. p.
563 2022.05.20.22275349. Available from:
564 <https://www.medrxiv.org/content/10.1101/2022.05.20.22275349v2>
- 565 26. Diestel R. Graph Theory [Internet]. 5th ed. Heidelberg: Springer-Verlag; 2017
566 [cited 2022 Jul 11]. Available from: <https://diestel-graph-theory.com/eBook.html>
- 567 27. Salomé M, Kesse-Guyot E, Fouillet H, Touvier M, Hercberg S, Huneau J-F,
568 Mariotti F. Development and evaluation of a new dietary index assessing nutrient
569 security by aggregating probabilistic estimates of the risk of nutrient deficiency in
570 two French adult populations. *Br J Nutr Cambridge University Press*;
571 2021;126:1225–36.
- 572 28. Dijkstra EW. A Short Introduction to the Art of Programming [Internet].
573 Technische Hogeschool Eindhoven; 1971. Available from:
574 <https://books.google.fr/books?id=JE8ZAQAIAAJ>
- 575 29. Fouillet H, Dussiot A, Perraud E, Wang J, Huneau J-F, Kesse-Guyot E, Mariotti
576 F. Plant to animal protein ratio in the diet: nutrient adequacy, long-term health
577 and environmental pressure. *Front Nutr* [Internet] 2023 [cited 2023 Jun 20];10.
578 Available from: <https://www.frontiersin.org/articles/10.3389/fnut.2023.1178121>
- 579 30. Rossum G van, Drake FL. The Python language reference. Release 3.0.1 [Repr.].
580 Hampton, NH: Python Software Foundation; 2010. 109 p.
- 581 31. Hagberg A, Swart P, S Chult D. Exploring network structure, dynamics, and
582 function using networkx [Internet]. Los Alamos National Lab. (LANL), Los
583 Alamos, NM (United States); 2008 Jan. Report No.: LA-UR-08-05495; LA-UR-

- 584 08-5495. Available from: [https://www.osti.gov/biblio/960616-exploring-network-](https://www.osti.gov/biblio/960616-exploring-network-structure-dynamics-function-using-networkx)
585 [structure-dynamics-function-using-networkx](https://www.osti.gov/biblio/960616-exploring-network-structure-dynamics-function-using-networkx)
- 586 32. Aynaoud T. python-louvain x.y: Louvain algorithm for community detection
587 [Internet]. 2020. Available from: <https://github.com/taynaud/python-louvain>
- 588 33. GBD 2013 Mortality and Causes of Death Collaborators. Global, regional, and
589 national age–sex specific all-cause and cause-specific mortality for 240 causes of
590 death, 1990–2013: a systematic analysis for the Global Burden of Disease Study
591 2013. *The Lancet* 2015;385:117–71.
- 592 34. THE LANCET. Global Burden of Disease: GBD cause and risk summaries
593 [Internet]. 2023 [cited 2023 Jun 16]. Available from:
594 <https://www.thelancet.com/gbd/summaries>
- 595 35. Wang J, Hasanalieva G, Wood L, Anagnostopoulos C, Ampadogiannis G,
596 Bempelou E, Kiouisi M, Markellou E, Iversen PO, Seal C, et al. Effect of wheat
597 species (*Triticum aestivum* vs *T. spelta*), farming system (organic vs
598 conventional) and flour type (wholegrain vs white) on composition of wheat flour
599 – Results of a retail survey in the UK and Germany – 3. Pesticide residue content.
600 *Food Chem X* 2020;7:100089.
- 601 36. eurostat. Organic farming statistics [Internet]. 2022 [cited 2022 Sep 15].
602 Available from: [https://ec.europa.eu/eurostat/statistics-](https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Organic_farming_statistics)
603 [explained/index.php?title=Organic_farming_statistics](https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Organic_farming_statistics)
- 604 37. Pereira PM de CC, Vicente AF dos RB. Meat nutritional composition and
605 nutritive role in the human diet. *Meat Sci* 2013;93:586–92.
- 606 38. Richter CK, Skulas-Ray AC, Champagne CM, Kris-Etherton PM. Plant Protein
607 and Animal Proteins: Do They Differentially Affect Cardiovascular Disease
608 Risk? *Adv Nutr* 2015;6:712–28.
- 609 39. Mariotti F. Animal and Plant Protein Sources and Cardiometabolic Health. *Adv*
610 *Nutr* 2019;10:S351–66.
- 611 40. Ankar A, Kumar A. Vitamin B12 Deficiency. StatPearls. [Internet] Treasure
612 Island (FL): StatPearls Publishing; 2022 [cited 2022 Nov 29]. Available from:
613 <http://www.ncbi.nlm.nih.gov/books/NBK441923/>
- 614 41. Perraud E, Wang J, Salomé M, Huneau J-F, Lapidus N, Mariotti F. Plant and
615 Animal Protein Intakes Largely Explain the Nutritional Quality and Health Value
616 of Diets Higher in Plants: A Path Analysis in French Adults. *Front Nutr* [Internet]
617 2022 [cited 2022 Aug 23];9. Available from:
618 <https://www.frontiersin.org/articles/10.3389/fnut.2022.924526>
- 619 42. Kesse-Guyot E, Touvier M, Henegar A, Czernichow S, Galan P, Hercberg S,
620 Castetbon K. Higher adherence to French dietary guidelines and chronic diseases

- 621 in the prospective SU.VI.MAX cohort. *Eur J Clin Nutr* Nature Publishing Group;
622 2011;65:887–94.
- 623 43. Magrini M-B, Fernandez-Inigo H, Doré A, Pauly O. How institutional food
624 services can contribute to sustainable agrifood systems? Investigating legume-
625 serving, legume-cooking and legume-sourcing through France in 2019. *Rev Agric*
626 *Food Environ Stud* 2021;102:297–318.
- 627 44. Michie S, Ashford S, Sniehotta FF, Dombrowski SU, Bishop A, French DP. A
628 refined taxonomy of behaviour change techniques to help people change their
629 physical activity and healthy eating behaviours: The CALO-RE taxonomy.
630 *Psychol Health* Routledge; 2011;26:1479–98.
- 631 45. Samdal GB, Eide GE, Barth T, Williams G, Meland E. Effective behaviour
632 change techniques for physical activity and healthy eating in overweight and
633 obese adults; systematic review and meta-regression analyses. *Int J Behav Nutr*
634 *Phys Act* 2017;14:42.
- 635 46. Mariotti F, Havard S, Morise A, Nadaud P, Sirot V, Wetzler S, Margaritis I.
636 Perspective: Modeling Healthy Eating Patterns for Food-Based Dietary
637 Guidelines—Scientific Concepts, Methodological Processes, Limitations, and
638 Lessons. *Adv Nutr* 2021;12:590–9.
- 639 47. Britten P, Cleveland LE, Koegel KL, Kuczynski KJ, Nickols-Richardson SM.
640 Impact of Typical Rather than Nutrient-Dense Food Choices in the US
641 Department of Agriculture Food Patterns. *J Acad Nutr Diet* 2012;112:1560–9.
- 642 48. ANSES. Étude individuelle nationale des consommations alimentaires 3 (INCA
643 3) [Internet]. 2017 Jun. Available from:
644 <https://www.anses.fr/fr/system/files/NUT2014SA0234Ra.pdf>
- 645 49. ANSES. AVIS de l'Agence nationale de sécurité sanitaire de l'alimentation, de
646 l'environnement et du travail relatif à l'évaluation des apports en vitamines et
647 minéraux issus de l'alimentation non enrichie, de l'alimentation enrichie et des
648 compléments alimentaires dans la population française : estimation des apports
649 usuels, des prévalences d'inadéquation et des risques de dépassement des limites
650 de sécurité [Internet]. 2015 [cited 2023 Jun 22]. Available from:
651 <https://www.anses.fr/fr/system/files/NUT2012sa0142.pdf>

652