



HAL
open science

Characterization of milk small extracellular vesicles to study adaptation to lactation in ruminants

Céline Boby, A Delavaud, J Pires, Laurent-Emmanuel Monfoulet, S Bes, S Emery, L Bernard, C Leroux, A Imbert, M Tourret, et al.

► To cite this version:

Céline Boby, A Delavaud, J Pires, Laurent-Emmanuel Monfoulet, S Bes, et al.. Characterization of milk small extracellular vesicles to study adaptation to lactation in ruminants. 74th Annual Meeting of the European Federation of Animal Science, Aug 2023, Lyon, France. 10.3920/978-90-8686-936-7 . hal-04195798

HAL Id: hal-04195798

<https://hal.inrae.fr/hal-04195798>

Submitted on 4 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Characterization of milk small extracellular vesicles to study adaptation to lactation in ruminants

C. Bobby¹, A. Delavaud¹, J. Pires¹, L.E. Monfoulet², S. Bes¹, S. Emery¹, L. Bernard¹, C. Leroux¹, A. Imbert¹, M. Tourret¹, F. Fournier³, D. Roux³, H. Sauerwein⁴ and M. Bonnet¹

¹Université Clermont Auvergne, INRAE, VetAgro Sup, UMR Herbivores, Centre INRAE CARA, 63122 Saint-Genès-Champanelle, France, ²Université Clermont Auvergne, INRAE, Human Nutrition Unit, Centre INRAE CARA, 63122 Saint-Genès-Champanelle, France, ³INRAE, Herbipôle, Centre INRAE CARA, 63122 Saint-Genès-Champanelle, France, ⁴University of Bonn, Institute of Animal Science, Physiology Unit, Katzenburgweg 7, 53115 Bonn, Germany; celine.bobby@inrae.fr

Small extracellular vesicles (EV) are secreted into the extracellular space by all cells. Due to the diversity of their cellular origin and the molecules they contain, small EVs have the ability to ensure extracellular communication and carry molecular signatures of their tissue of origin and its physiological state. In ruminants, early lactation is characterized by profound changes in energy balance and metabolic status. To explore physiological adaptations during early lactation, this study investigated the use of non-invasive milk EVs to identify specific indicators of inter-organ signalling. Milk samples were collected from 8 cows on weeks 2 and 7 postpartum, corresponding to negative and neutral energy balance, respectively. Small EVs were isolated by ultracentrifugation coupled with size exclusion chromatography and characterized by morphological, biophysical and biochemical criteria. Labelled-free shotgun quantitative proteomics was performed by nanoLC-MS/MS. Electron microscopy revealed cup-shaped vesicles with a diameter of about 100 nm, characteristic of small EVs. The diameter was confirmed by Tunable Resistive Pulse Sensing and the specificity of small EVs isolation by the presence of cytoplasmic (TSG101) and membrane (CD63) markers. A total of 508 proteins were identified in milk EVs at weeks 2 and 7 of lactation. Multilevel PCA analysis showed a clear separation between the 2 time points, indicating a strong effect of lactation stage on the protein composition of milk EVs. This proteomic dataset will be analysed to identify small EV molecular signatures of tissue interactions that coordinate nutrient partitioning and adaptation during early lactation.
