



HAL
open science

Monitoring liveweight in Sarda dairy sheep using a walk-over-weighing system

Mauro Decandia, Marco Acciaro, Valeria Giovanetti, Giovanni Molle,
Francesco Chessa, Irène Llach, Eliel González García

► To cite this version:

Mauro Decandia, Marco Acciaro, Valeria Giovanetti, Giovanni Molle, Francesco Chessa, et al.. Monitoring liveweight in Sarda dairy sheep using a walk-over-weighing system. 74th Annual Meeting of the European Federation of Animal Science, Aug 2023, Lyon, France. Wageningen Academic Publishers, 29, pp.504, Book of Abstracts of the 74th Annual Meeting of the European Federation of Animal Science. 10.3920/978-90-8686-936-7. hal-04195801

HAL Id: hal-04195801

<https://hal.inrae.fr/hal-04195801v1>

Submitted on 4 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Book of Abstracts of the 74th Annual Meeting of the European Federation of Animal Science



**Book of abstracts No. 29 (2023)
Lyon, France
26 August – 1 September, 2023**

Book of Abstracts of the 74th Annual Meeting of the European Federation of Animal Science

Lyon, France, 26 August – 1 September, 2023



EAAP Scientific Committee:

F. Miglior
L. Pinotti
L. Boyle
D. Kenny
M. Lee
M. De Marchi
V.A.P. Cadavez
S. Millet
R. Evans
L. Gasco
M. Pastell
G. Pollott (secretary)
H. Spoolder (chair)



Session 39

Monitoring liveweight in Sarda dairy sheep using a walk-over-weighing system

M. Decandia¹, M. Acciaro¹, V. Giovanetti¹, G. Molle¹, F. Chessa¹, I. Llach² and E. González-García²

¹AGRIS Sardegna, Loc. Bonassai S.S. 291 Sassari-Fertilia, 07100 Sassari, Italy,

²SELMET, INRAE, Montpellier SupAgro, CIRAD, Université Montpellier, 34060 Montpellier, France;

mdecandia@agrisricerca.it

Live weight (**LW**) monitoring is important in livestock management to check nutritional, reproduction and welfare status of animals. In dairy sheep system LW measurement is less frequent because it is time and labour consuming. The use of automated weighing system could facilitate this practice. Within the H2020 Techcare project, aiming to use innovative technologies to improve welfare management for small ruminant systems, a short-test was run to evaluate the precision and accuracy of LW measures carried out using an automated walk-over-weighing (**WoW**) scale in comparison with a static scale. Thirty-six dry Sarda dairy ewes (LW, mean \pm SE 55.49 \pm 0.92) were used in a 3-day session. Ewes were daily weighed 3 times following the same circuit, which included a first static LW measurement; once a static position was achieved and LW recorded, the animals continued and traversed the WoW scale for the WoW LW recording. The ewes were previously accustomed to the circuit during a pre-experimental week. Raw static and automatic LW data were first filtered for removing misbehaviours and outlier with Kalman filter algorithm. The concordance correlation coefficient (CCC) has been then calculated (Model Evaluation System) to simultaneously account for precision and accuracy. The two component of CCC (=0.98), correlation coefficient estimate (r , that measure precision) and the bias correction factor (C_b , that indicates the accuracy) were respectively 0.98 and 0.99 indicating high agreement between measures. The WoW system evaluated here is an alternative to the static scales conventionally used on dairy sheep farms. If sound filtration of raw data is applied, WoW could contribute to the close (daily) monitoring of individual LW without operator intervention (i.e. voluntary weighing), taking animal welfare into account (i.e. no stress related to the weighing session on static scales), and potentially detecting nutrition issues highlighted by LW changes.