



HAL
open science

Performance, meat quality and blood parameters in four strains of organic broilers differ according to range use

Claire Bonnefous, Anne Collin, Laurence A. Guilloteau, Karine Germain, Laure Ravon, Thierry Bordeau, Pascal Chartrin, Estelle Godet, Estelle Cailleau Audouin, Nathalie Couroussé, et al.

► To cite this version:

Claire Bonnefous, Anne Collin, Laurence A. Guilloteau, Karine Germain, Laure Ravon, et al.. Performance, meat quality and blood parameters in four strains of organic broilers differ according to range use. 2023. hal-04199787

HAL Id: hal-04199787

<https://hal.inrae.fr/hal-04199787v1>

Preprint submitted on 8 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Performance, meat quality and blood parameters in four strains of organic broilers differ according to range use

Claire Bonnefous (✉ claire.bonnefous@inrae.fr)

INRAE, Université de Tours, BOA

Anne Collin

INRAE, Université de Tours, BOA

Laurence A. Guilloteau

INRAE, Université de Tours, BOA

Karine Germain

INRAE, UE EASM

Laure Ravon

INRAE, UE EASM

Thierry Bordeau

INRAE, Université de Tours, BOA

Pascal Chartrin

INRAE, Université de Tours, BOA

Estelle Godet

INRAE, Université de Tours, BOA

Estelle Cailleau-Audouin

INRAE, Université de Tours, BOA

Nathalie Couroussé

INRAE, Université de Tours, BOA

Emilie Raynaud

INRAE, Université de Tours, BOA

Sandrine Mignon-Grasteau

INRAE, Université de Tours, BOA

Maxime Reverchon

SYSAF

Simona Mattioli

University of Perugia

Cesare Castellini

University of Perugia

Elisa Angelucci

University of Perugia

Vanessa Guesdon

France Junia

Ludovic Calandreau

INRAE, CNRS, IFCE, Université de Tours, PRC

Cécile Berri

INRAE, Université de Tours, BOA

Elisabeth Le Bihan-Duval


INRAE, Université de Tours, BOA

Article

Keywords: poultry, free-range, range use behaviour, growth performance, biomarkers

Posted Date: July 25th, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-3164049/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Chicken meat production in organic systems involves free-range access where animals can express foraging and locomotor behaviours. These behaviours may promote outdoor feed intake, but at the same time induce a loss of energy in exploring. More generally, the relationship of range use with metabolism, welfare, health, growth performance and meat quality needs to be better understood. We studied four strains of intermediate (JA757) to slow-growing (S757N, White Bresse and a dual-purpose strain) meat-type chickens with outdoor access. We selected 25 males high- (HR) and low-rangers (LR) per strain. Only in JA757, HR exhibited lower body weight before range access, which may have predisposed them to use the range more. Carcass weight and/or carcass yield were significantly lower in HR compared to LR, showing a negative trade-off between range use and growth performance in all strains. Breast meat yellowness was higher in HR compared to LR in JA757 and the dual-purpose strain, probably due to carotenoids intake from the grass. No relationship between range use and welfare indicators at slaughter was reported whatever the strain. Chicken metabolism differed by range use as HR and LR diverged for blood biomarkers of oxidative and metabolic status, innate and inflammatory system response.

Introduction

Meat consumption of free-ranging livestock has increased in recent decades¹, driven by the consumer's perception that free range and animal welfare are related². By providing an outdoor range, broilers have the possibility to express natural behaviours such as exploration and foraging to a higher degree than in indoor systems³. Nevertheless, a high variability of outdoor range use is often reported between flocks⁴⁻⁶. Several studies in chickens showed that range use differs between individuals and is consistent within an individual over time and across contexts or situations^{7,8}, therefore qualifying ranging behaviour as a potential personality trait⁹.

Free-range access can influence chicken meat quality and composition but also blood physiological parameters due to animals' foraging and locomotor activities^{10,11}, but results are sometimes inconsistent. The diversity of slow-growing genotypes studied, the varying level of free-range use and the lack of individual assessment of chickens' ranging behaviours may be the source of contradictory results. In fast-growing strains, the body weight (BW) of ranging birds was significantly lower than that of non-ranging birds^{12,13}. Accordingly, the body weight of slow-growing indoor birds was significantly higher (4–5%) compared to slow-growing birds with outdoor access¹⁰. However, higher BW was related to more time spent outdoors in another slow-growing strain studied by Marchewka et al.¹⁴. Similarly, several studies showed an effect of range use on slow-growing breast meat colour with darker, redder and yellower breast meat^{10,15} while no effect was reported in other studies^{16,17}. The inconsistency of these findings might be due to the genetic background.

To improve the sustainability of free-range systems considering economic, welfare and product quality impacts, a better understanding of the consequences and mechanism activated by range use is needed. Therefore, the objective of the current study was to investigate the relationship between range use (high vs low) and live performance (before and after outdoor access) as well as several indicators or blood parameters related to animal welfare and health, metabolism and the carcass and meat quality. For this study we considered four strains of broilers with intermediate to slow growth rates, to reflect the multiplicity of genetic strains used in free-range systems in Europe.

Results

Variation of BW according to range use

In the JA757 strain, HR exhibited a 14% lower live body weight than LR at 29 days of age, before range access (Fig. 2). After 21 days of range access, the body weight of HR was about 16% and 7% lower compared to LR in JA757 and White Bresse strains, respectively. The differences were broadly maintained at slaughter, with BW around 11% and 7% lower in HR compared to LR in JA757 and White Bresse strains, respectively. Live body weight was not significantly influenced by range use in the dual-purpose strain, and we only reported a tendency at 6.5% in the S757N strain with a reduction of about 6% in the HR compared to the LR group at 85 days old.

Range use relationship with carcass, tibial and meat quality traits

Carcass traits

Range use was negatively related to the carcass traits in all the four studied strains. We observed a reduction of 12%, 8% and 7% of carcass weight in the HR compared to the LR in JA757, S757N and the White Bresse, respectively (Table 1). We reported a relative reduction of 2% and 1% of carcass yield in the HR compared to the LR in S757N and the dual-purpose strains, respectively (Table 1). Similarly, we observed a reduction of thigh and breast weights in the HR compared to the LR in JA757 and S757N and of thigh weight in White Bresse. Moreover, a reduction of 3% and 2% of thigh yield was associated to greater range use in S757N and the dual-purpose strain, respectively.

Table 1
Effects of range use on growth and tibial quality traits (mean ± SD) in four different strains of organic broilers

Item	JA757			S757N			White Bresse			Dual-purpose		
	LR (n = 25)	HR (n = 25)	<i>P</i> -value	LR (n = 25)	HR (n = 25)	<i>P</i> -value	LR (n = 25)	HR (n = 25)	<i>P</i> -value	LR (n = 25)	HR (n = 25)	<i>P</i> -value
Carcass weight (g)	1973 ± 268	1748 ± 227	0.006	1740 ± 243	1606 ± 150	0.047	1802 ± 131	1672 ± 145	0.006	997 ± 120	1026 ± 125	0.605
Carcass yield (% of BW)	69.4 ± 1.3	68.6 ± 1.5	0.072	69.0 ± 1.6	67.6 ± 1.3	0.006	65.7 ± 1.4	65.1 ± 1.4	0.176	63.2 ± 1.3	62.3 ± 1.1	0.009
Breast weight (g)	233 ± 37	201 ± 31	0.006	183 ± 30	168 ± 18	0.047	176 ± 15	165 ± 19	0.068	83 ± 13	84 ± 12	0.702
Breast yield (% of BW)	16.4 ± 1.1	15.8 ± 1.0	0.072	14.5 ± 1.1	14.1 ± 1.0	0.236	12.8 ± 0.7	12.8 ± 0.8	0.994	10.5 ± 0.9	10.3 ± 0.6	0.605
Thigh weight (g)	351 ± 48	315 ± 37	0.012	322 ± 39	300 ± 33	0.047	358 ± 27	332 ± 27	0.006	195 ± 23	199 ± 24	0.653
Thigh yield (% of BW)	24.7 ± 0.9	24.8 ± 0.5	0.518	26.0 ± 1.3	25.2 ± 1.0	0.047	26.2 ± 0.9	25.9 ± 0.8	0.316	24.8 ± 0.5	24.2 ± 0.6	0.018
	LR (n = 11)	HR (n = 12)	<i>P</i> -value	LR (n = 10)	HR (n = 12)	<i>P</i> -value	LR (n = 11)	HR (n = 12)	<i>P</i> -value	LR (n = 13)	HR (n = 12)	<i>P</i> -value
Tibia length (mm)	134.0 ± 5.6	128.2 ± 4.5	0.006	137.7 ± 5.3	137.0 ± 6.5	0.765	153.0 ± 2.3	150.5 ± 4.3	0.288	126.8 ± 5.8	128.1 ± 5.9	0.591
Tibia diameter (mm)	8.2 ± 1.1	7.8 ± 0.6	0.053	8.1 ± 0.8	7.8 ± 0.9	0.758	8.9 ± 0.8	8.8 ± 0.7	0.786	7.6 ± 0.6	7.1 ± 0.4	0.042
Tibia bone-breaking strength (N)	238.4 ± 54.7	242.6 ± 29.8	0.021	248.2 ± 33.6	230.6 ± 50	0.758	311.6 ± 61.5	296.8 ± 46.5	0.786	189.3 ± 34.5	173.2 ± 26.2	0.302

Leg health and welfare

We measured several welfare indicators at slaughter (Table 2): hock burns, pododermatitis, struggling activity on the shackle line and total duration of wing flapping. None of these welfare indicators were significantly different depending on range use. We evaluated leg health thanks to tibial measurements (Table 3), we observed a 4% shorter, 5% thinner even though not significantly different (*p*-value = 0.053) and 2% stronger tibial bone in HR than in LR in the JA757 strain. In the dual-purpose strain, we also reported 7% thinner tibial bone in HR than LR.

Table 2 Effects of range use on welfare indicators in four different strains of organic broilers

	JA757			S757N			White Bresse			Dual-purpose																	
	LR	HR	<i>P</i> -value	LR	HR	<i>P</i> -value	LR	HR	<i>P</i> -value	LR	HR	<i>P</i> -value															
Total duration of wing flapping (mean ± SD)	4,11 ± 4,82			3,65 ± 4,85			1,64 ± 4,43			3,79 ± 4,97																	
Scores of leg damage	0	1	2	0	1	2	0	1	2	0	1	2															
Hock burn (% of flock)	60	32	8	72	28	0	8	0	92	8	0	1000	5	44	4	20	64	16	0.192	100	0	0	92	8	0		
Pododermatitis (% of flock)	4	48	48	0	60	40	0.880	2	56	16	28	60	12	1000	1	72	12	8	84	8	1.000	25	75	0	40	60	0
Activity on the slaughter line (Y/N)	Y	N	Y	Y	N	Y	Y	N	Y	Y	N	Y	Y	N	Y	Y	N	Y	Y	N	Y	Y	N				
Chickens (%)	36	64	28	72	28	72	0.880	28	72	20	80	1000	12	88	8	92	1000	2	71	28	72	1000	9				

Table 3

Effects of range use on *Pectoralis major* weight, pH and quality measurements (mean \pm SD) in four different strains of organic broilers

Item	JA757			S757N			White Bresse			Dual-Purpose		
	LR (n = 25)	HR (n = 25)	P-value	LR (n = 25)	HR (n = 25)	P-value	LR (n = 25)	HR (n = 25)	P-value	LR (n = 25)	HR (n = 25)	P-value
Muscle characteristics												
Pectoralis major muscle weight (g)	181 \pm 29	156 \pm 25	0.006	139 \pm 24	126 \pm 14	0.225	128 \pm 12	120 \pm 14	0.243	62 \pm 9	63 \pm 9	0.802
pH 15 min post-mortem (pH15)	6.56 \pm 0.13	6.49 \pm 0.12	0.053	6.40 \pm 0.14	6.38 \pm 0.15	0.837	6.46 \pm 0.13	6.48 \pm 0.13	0.903	6.15 \pm 0.15	6.17 \pm 0.14	0.776
Ultimate pH (pHu)	5.67 \pm 0.07	5.78 \pm 0.12	0.004	5.77 \pm 0.10	5.76 \pm 0.11	0.837	5.49 \pm 0.11	5.46 \pm 0.10	0.903	5.65 \pm 0.08	5.61 \pm 0.11	0.495
Meat quality traits												
Lightness (L*)	48.1 \pm 1.6	46.5 \pm 2.1	0.007	45.1 \pm 2.0	46.0 \pm 1.8	0.504	47.7 \pm 3.1	48.5 \pm 3.5	0.903	44.9 \pm 2.5	45.4 \pm 2.8	0.776
Redness (a*)	-0.4 \pm 0.5	0.3 \pm 0.5	< 0.001	0.7 \pm 1.0	0.6 \pm 0.9	0.921	0.7 \pm 0.8	0.7 \pm 1.1	0.947	1.8 \pm 1.5	2.1 \pm 1.2	0.776
Yellowness (b*)	10.2 \pm 1.2	11.3 \pm 1.4	0.009	11.1 \pm 1.4	11.1 \pm 1.7	0.973	11.9 \pm 1.3	11.8 \pm 1.0	0.903	10.2 \pm 1.4	12.0 \pm 1.7	0.002
Drip loss (DL, %)	1.1 \pm 0.4	1.1 \pm 0.3	0.487	0.9 \pm 0.3	0.9 \pm 0.3	0.837	1.1 \pm 0.3	1.2 \pm 0.3	0.903	1.5 \pm 0.5	1.5 \pm 0.7	0.776
Thawing-cooking loss (TCL, %)	14.3 \pm 2.6	15.6 \pm 2.8	0.121	13.5 \pm 4.7	14.7 \pm 7.4	0.837	11.0 \pm 1.7	11.1 \pm 1.8	0.903	17.0 \pm 3.4	16.4 \pm 3.3	0.776
Warner Bratzler shear force (WB, N/cm ²)	17.0 \pm 2.5	16.3 \pm 2.1	0.323	16.2 \pm 2.9	16.7 \pm 2.2	0.837	16.3 \pm 1.7	16.2 \pm 2.1	0.903	22.4 \pm 5.5	20.6 \pm 7.6	0.776

Breast and thigh meat quality and composition

In the JA757 strain, HR chickens had a higher pHu (+ 0.09 pH units) associated with darker, redder and more yellow meat than LR chickens. In the dual-purpose strain, breast meat was also yellower in HR compared to LR chickens (Table 3).

Regarding thigh meat, whatever the strain and the level of range use, the main fatty acids were palmitic (C16:0), stearic (C18:0), oleic (C18:1) and linoleic (C18:2) acids. Range use only increased α -linolenic acid (C18:3 n-6) content in the dual-purpose strain, with a 22% higher percentage observed in HR compared to LR chickens (Table 4).

Table 4
Effects of range use on thigh muscle pH and meat quality measurements (mean \pm SD) in four strains of organic broilers

Item	JA757			S757N			White Bresse			Dual-Purpose		
	LR (n = 25)	HR (n = 25)	<i>P</i> -value	LR (n = 25)	HR (n = 25)	<i>P</i> -value	LR (n = 25)	HR (n = 25)	<i>P</i> -value	LR (n = 25)	HR (n = 25)	<i>P</i> -value
Muscle characteristics												
pHu in the thigh muscle	6.02 \pm 0.10	6.02 \pm 0.09	0.808	5.87 \pm 0.12	5.86 \pm 0.07	0.977	5.75 \pm 0.11	5.73 \pm 0.09	0.562	5.90 \pm 0.09	5.90 \pm 0.09	0.922
Chemical (%) and fatty acids composition (% of total fatty acids)												
	LR (n = 11)	HR (n = 12)	<i>P</i> -value	LR (n = 10)	HR (n = 12)	<i>P</i> -value	LR (n = 11)	HR (n = 12)	<i>P</i> -value	LR (n = 13)	HR (n = 12)	<i>P</i> -value
Lipids	6.8 \pm 1.1	6.2 \pm 1.0	0.399	5.5 \pm 2.3	5.5 \pm 2.0	0.999	4.5 \pm 0.6	4.1 \pm 0.8	0.405	4.2 \pm 0.9	4.3 \pm 0.7	0.859
Myristic (C14:0)	0.5 \pm 0.0	0.4 \pm 0.1	0.223	0.4 \pm 0.0	0.4 \pm 0.0	0.999	0.5 \pm 0.1	0.4 \pm 0.1	0.662	0.5 \pm 0.0	0.5 \pm 0.1	0.393
Palmitic (C16:0)	20.7 \pm 1.2	19.5 \pm 0.8	0.095	19.6 \pm 0.8	19.8 \pm 0.9	0.999	19.7 \pm 1.0	19.3 \pm 1.6	0.664	19.6 \pm 1.2	18.6 \pm 1.3	0.300
Stearic (C18:0)	6.0 \pm 1.0	6.9 \pm 1.3	0.223	8.1 \pm 1.6	8.2 \pm 1.0	0.999	8.4 \pm 0.8	9.1 \pm 1.2	0.348	8.5 \pm 0.8	9.0 \pm 0.6	0.393
Palmitoleic (C16:1)	3.5 \pm 0.8	2.6 \pm 0.6	0.095	1.9 \pm 0.3	1.7 \pm 0.3	0.525	2.0 \pm 0.3	1.6 \pm 0.3	0.150	1.4 \pm 0.3	1.3 \pm 0.3	0.393
Oleic (C18:1)	37.3 \pm 1.4	37.1 \pm 1.6	0.828	36.5 \pm 2.3	36.7 \pm 1.6	0.999	36.3 \pm 1.4	35.4 \pm 1.1	0.348	35.4 \pm 1.5	34.9 \pm 1.5	0.526
Paullinic (C20:1)	0.34 \pm 0.02	0.35 \pm 0.04	0.593	0.37 \pm 0.06	0.39 \pm 0.04	0.525	0.35 \pm 0.06	0.38 \pm 0.08	0.365	0.36 \pm 0.02	0.37 \pm 0.02	0.300
Linoleic (C18:2n-6)	27.3 \pm 1.9	28.6 \pm 1.3	0.223	27.8 \pm 0.9	28.1 \pm 1.3	0.999	28.1 \pm 1.6	28.1 \pm 3.1	0.948	29 \pm 1.9	29.9 \pm 1.6	0.393
Alpha-linolenic (C18:3n-6)	2.1 \pm 0.2	2.3 \pm 0.1	0.095	1.9 \pm 0.1	2.0 \pm 0.2	0.904	1.7 \pm 0.2	1.7 \pm 0.3	0.674	1.8 \pm 0.2	2.2 \pm 0.2	0.007
Saturated fatty acids	28.2 \pm 2.1	28.5 \pm 2.5	0.828	27.7 \pm 0.9	29.5 \pm 1.7	0.270	28.7 \pm 1.4	28.6 \pm 1.8	0.948	28.8 \pm 1.3	28.2 \pm 1.7	0.526
Monounsaturated fatty acids	40.3 \pm 2.6	39.8 \pm 2.8	0.828	39.8 \pm 2.2	38.5 \pm 2.7	0.999	38.9 \pm 1.6	37.4 \pm 1.3	0.170	36.9 \pm 1.5	36.7 \pm 1.7	0.827
Polyunsaturated fatty acids	31.8 \pm 2.9	33.1 \pm 3.2	0.728	33.2 \pm 2.1	33 \pm 2.8	0.999	32.1 \pm 2.0	32.9 \pm 3.4	0.638	34.4 \pm 1.1	35.3 \pm 2	0.393
n-6/ n-3	6.8 \pm 1.1	7.0 \pm 1.3	0.828	6.8 \pm 1.1	6.6 \pm 1.4	0.999	7.2 \pm 1.7	6.7 \pm 1.1	0.663	6.7 \pm 0.7	6.3 \pm 0.5	0.393

Range use and blood physiological parameters at slaughter

As observed in thigh meat, the main fatty acids of plasma in the four studied strains were palmitic (C16:0), stearic (C18:0), oleic (C18:1) and linoleic (C18:2). We reported a reduction by 26% of the n-6/n-3 ratio with greater range use in the S757N strain, while we observed an increase by 62% of this ratio with greater range use in the White Bresse strain (Table 5). In the White Bresse strain, HR chickens also had a 24% higher triglyceride concentration than LR chickens. Additionally, their Total Antioxidant Status was increased by 10%. In the S757N strain, HR chickens had lower blood concentration or activity for uric acid (-19%), creatine kinase (-17%), haptoglobin-like (-30%) and lysozyme (-25%) than LR chickens. Similar effects were observed in the JA757 strain for creatine kinase (-31%) and lysozyme (-29%). In this genetic strain, the HR birds were also characterized by lower Total Antioxidant Status (-8%) and tocopherols concentration (-35%), but a higher value of hydrogen peroxide (+22%). Regarding the dual-purpose strain, the only significant difference was in creatine kinase activity, decreased by 22% with range use.

Table 5

Effects of range use on broiler's blood physiological parameters (mean \pm SD) at slaughter time in four different strains of organic broilers

	JA757			S757N			White Bresse			Dual-Purpose		
	LR (n = 25)	HR (n = 25)	<i>P</i> -value	LR (n = 25)	HR (n = 25)	<i>P</i> -value	LR (n = 25)	HR (n = 25)	<i>P</i> -value	LR (n = 25)	HR (n = 25)	<i>P</i> -value
Uric acid ² (mg/L)	41 \pm 15	33 \pm 12	0.108	43 \pm 11	35 \pm 7	0.020	28 \pm 9	32 \pm 8	0.269	28 \pm 6	32 \pm 10	0.586
Glucose ² (mg/L)	2282 \pm 110	2296 \pm 115	0.747	2454 \pm 218	2384 \pm 173	0.325	2514 \pm 193	2584 \pm 124	0.274	2433 \pm 138	2392 \pm 107	0.516
Carbonyl ¹ (nmol /mg of protein)	0.9 \pm 0.4	0.9 \pm 0.5	0.765	0.8 \pm 0.2	0.7 \pm 0.2	0.325	0.7 \pm 0.2	0.8 \pm 0.2	0.269	1.2 \pm 0.5	1.0 \pm 0.5	0.468
Triglycerides ² (mg/L)	231 \pm 51	199 \pm 54	0.068	183 \pm 49	165 \pm 38	0.285	209 \pm 45	259 \pm 64	0.024	163 \pm 36	187 \pm 29	0.060
Creatine Kinase activity ¹ (UI/L)	5801 \pm 1984	4013 \pm 1045	0.003	3871 \pm 951	3228 \pm 912	0.054	3195 \pm 839	3765 \pm 922	0.120	4848 \pm 1621	3775 \pm 1274	0.007
Infectious bronchitis antibody titres ²	6363 \pm 1580	5988 \pm 1699	0.679	3668 \pm 1978	3232 \pm 1870	0.482	5174 \pm 2026	5769 \pm 1660	0.393	4101 \pm 2450	3410 \pm 2671	0.516
Haptoglobin-like activity ² (mg/ml)	2.2 \pm 0.4	2.0 \pm 0.6	0.549	2.3 \pm 0.6	1.6 \pm 0.6	0.003	2.3 \pm 0.7	2.2 \pm 0.5	0.628	1.4 \pm 0.3	1.3 \pm 0.3	0.707
Lysozyme activity ² (mg/L)	5.5 \pm 1.9	3.9 \pm 1.2	0.002	4.0 \pm 1.1	3.0 \pm 0.7	0.003	4.0 \pm 1.3	3.9 \pm 1.4	0.690	2.8 \pm 1	2.9 \pm 0.9	0.707
H ₂ O ₂ ¹ (μ M)	10.2 \pm 2.8	12.4 \pm 3.7	0.046	9.2 \pm 4	11.4 \pm 4.5	0.144	11.0 \pm 2.4	10.5 \pm 2	0.626	18.5 \pm 2.6	18.4 \pm 3.3	0.946
Total Antioxidant Status ² (mmol/L)	1.2 \pm 0.2	1.1 \pm 0.1	0.003	1.1 \pm 0.2	1.1 \pm 0.1	0.373	1.0 \pm 0.1	1.1 \pm 0.1	0.024	1.0 \pm 0.1	1.0 \pm 0.1	0.801
Σ Tocols ² (μ mol/L)	2.6 \pm 0.8	1.7 \pm 0.7	< 0.001	2.8 \pm 1.4	2.3 \pm 1.4	0.146	2.2 \pm 1.5	2.3 \pm 1	0.404	4.9 \pm 4.0	3.7 \pm 2.4	0.516
Glutathione Peroxidase activity ² (U/L)	15433 \pm 2258	15205 \pm 1640	0.747	17206 \pm 2747	16716 \pm 3050	0.561	18393 \pm 2417	17577 \pm 2020	0.346	17135 \pm 2053	16666 \pm 1979	0.754
Fatty acids¹(% of total fatty acids)												
Myristic (C14:0)	0.15 \pm 0.11	0.20 \pm 0.11	0.204	0.12 \pm 0.06	0.14 \pm 0.04	0.369	0.17 \pm 0.11	0.17 \pm 0.10	1	0.14 \pm 0.05	0.12 \pm 0.05	0.468
Palmitic (C16:0)	16.8 \pm 1.8	16.4 \pm 1.5	0.677	16.8 \pm 2.1	17.3 \pm 1.4	0.476	15.3 \pm 2.2	15.5 \pm 1.7	1	15.6 \pm 1.4	15.5 \pm 1.3	0.861
Stearic (C18:0)	18.9 \pm 3.3	19.2 \pm 3.1	0.854	18.9 \pm 2.9	20.2 \pm 2.6	0.221	12.8 \pm 1.9	12.9 \pm 2.1	1	19.2 \pm 3.4	19.3 \pm 3.9	0.861
Palmitoleic (C16:1)	0.54 \pm 0.24	0.48 \pm 0.25	0.654	0.34 \pm 0.13	0.26 \pm 0.09	0.138	0.36 \pm 0.36	0.27 \pm 0.21	0.810	0.19 \pm 0.08	0.21 \pm 0.08	0.512
Oleic (C18:1)	12.2 \pm 3.7	12.7 \pm 3.6	0.960	11.3 \pm 1.8	10.5 \pm 1.2	0.221	9.9 \pm 3.1	10.5 \pm 2	0.891	9.4 \pm 1.6	9.7 \pm 1.1	0.468
Paullinic (C20:1)	0.13 \pm 0.16	0.16 \pm 0.19	0.960	0.12 \pm 0.15	0.09 \pm 0.13	0.554	-	-	NA	0.10 \pm 0.10	0.11 \pm 0.14	0.468
Linoleic (C18:2n-6)	16.4 \pm 2.9	17.6 \pm 4	0.654	18.2 \pm 5.1	15.4 \pm 3.4	0.221	21.6 \pm 5.4	23.5 \pm 3.3	0.891	18.5 \pm 4.6	19.1 \pm 4.2	0.790
Alpha-linolenic (C18:3n-6)	0.57 \pm 0.33	0.64 \pm 0.3	0.654	0.39 \pm 0.16	0.37 \pm 0.17	0.320	0.51 \pm 0.13	0.53 \pm 0.10	0.891	0.57 \pm 0.31	0.60 \pm 0.35	0.861

¹ Blood analysis on serum samples.² Blood analysis on plasma samples.

	JA757			S757N			White Bresse			Dual-Purpose		
Saturated fatty acids	36 ± 4	36 ± 4	0.960	36 ± 5	38 ± 3	0.221	28 ± 3	28 ± 4	0.891	35 ± 4	35 ± 5	0.861
Monounsaturated fatty acids	13 ± 4	13 ± 4	0.960	12 ± 2	11 ± 1	0.221	10 ± 3	11 ± 2	0.891	10 ± 2	10 ± 1	0.468
Polyunsaturated fatty acids	36 ± 4	40 ± 8	0.204	36 ± 5	35 ± 3	0.554	40 ± 7	39 ± 5	0.891	40 ± 3	41 ± 4	0.861
n-6/ n-3	4.3 ± 2.5	3.6 ± 2.2	0.654	7.5 ± 1.7	5.6 ± 2.1	0.002	4.4 ± 2.1	7.1 ± 1.2	< 0.001	4.3 ± 1.3	4.0 ± 1.7	0.468
¹ Blood analysis on serum samples.												
² Blood analysis on plasma samples.												

Discussion

Given the high societal expectations for animal welfare and the development of production systems offering animals access to the outdoors, it is important to acquire knowledge about their range use capabilities and the consequences on physiological and productive functions. Because outdoor access can relate to different types of production, our study considered 4 strains of chickens with different growth rates (from 16g/d to 36g/d) and different rearing times (from 71 to 106 days).

A first point to mention is that among the strains studied, the intra-population variability in range use (assessed by the FDI calculation) and in turn the gap between high- and low-rangers differed greatly, which may partly explain the variable relationships observed between range use and other studied parameters according to the strain.

Relationship between range use and growth performance

High-rangers of the JA757 strain had lower BW before and after range access than low-rangers, suggesting a possible impact of BW on range use in these medium-growing birds. Previous studies on fast-growing strains already reported weight difference prior to range access¹³. Moreover, when studying behavioural budget before range access, we reported a positive correlation between active behaviours (locomotion and foraging) and the later range use in the JA757 strain⁸. Altogether, these results suggest that in fast- or intermediate-growing strains, early-life locomotor activity might predict subsequent range use ability and that part of the effect is mediated by lower body weight. This does not seem to be the case in the slower-growing strains we studied, for which we did not report any difference in BW before range access between high and low rangers, nor any correlation between active behaviours indoors before range access and later range use⁸. The determinant of outdoor range use is most likely multifactorial and strain dependent. When considering the final body weight or the carcass weight at slaughter, a negative trade-off with range use was obvious for all the genotypes except the slowest-growing dual-purpose strain. Considering that locomotor activity on the range and range use are positively correlated¹⁸, our results seem consistent but highlight the issue of balancing the expression of natural behaviours within free-range and economic traits.

Range use and welfare indicators at slaughter

We did not observe any difference for leg health between HR and LR in the four studied strains. This is consistent with previous studies showing that outdoor range use did not impact foot pad dermatitis and hock burns scores in fast-growing strains¹³ or in slow-growing strains^{14,19}. The results obtained in strain JA757 were in line with previous studies that reported decreased tibia length and increased bone strength in birds with outdoor compared to birds reared indoors²⁰. As a decreased tibial length and midpoint cross-sectional area in females of fast-growing strains was associated with forced exercise²¹, better tibia bone health in HR compared to the LR may be due to their higher locomotor activity on the range. Therefore, our results indicated that higher range use interlinked with a supposed higher locomotor activity may positively affect bone health in the intermediate-growing line that was slaughtered at the younger age (around ten weeks) compared to other strains. Finally, it has been previously shown that struggling on the shackle line was more intense in slow-growing than in fast-growing lines²². The duration of wing flapping was even shown to be heritable in a Label Rouge type line²³ whose growing rate corresponded to the strain S757N studied here. The present study did not highlight any relationship between range use and struggling behaviour at slaughter, which seems to have different determinisms.

Range use relationship with meat quality

Regarding breast meat quality, we observed a higher pH_u and a darker meat in HR than in LR in the JA757 strain only. Darker meat has already been reported in slow-growing birds with outdoor access compared to those reared indoors¹⁰. The darker breast meat colour (lower L* value) of the high-ranger chickens is consistent with their higher ultimate pH, which reveals lower glycogen store in their breast muscle²⁴, maybe related to a higher locomotor activity on the range. We also reported for this strain a slightly lower pH₁₅, which is a proxy of the muscle glycolytic activity early postmortem. It has been well described that in chicken (especially slow-growing ones) low breast muscle pH₁₅ is correlated to high struggling activity on the shackle line^{25,26}. Therefore, even though we did not observe significant difference in struggling activity on the shackle line, decreased pH₁₅ reported in high-ranger chickens may suggest higher muscle glycolytic activity before slaughter, which may have contributed to their higher pH_u as already shown²⁶. The differences in pH_u and pH₁₅ observed between HR and LR in the strain JA757 were neither related to differences in drip and cooking loss, nor in tenderness. No effect of range use on these traits was observed in the other three strains, which is consistent with previous results¹⁵. Additionally, breast meat of HR was yellower than that of LR in both JA757 and dual-purpose strains. Increased yellowness has been reported in several other studies^{10,27,28} and could be caused by the intake of fresh plants (e.g., grass or clover) containing carotenoids^{11,29}. We can hypothesize that JA757 and dual-purpose high-rangers dedicated a higher percentage of their time to foraging activities, and/or their metabolism is able to assimilate more pigments than other strains from the ingested plants, leading to a higher enrichment of their muscles in carotenoids. Overall, range use and breast meat quality of slow-growing strains were barely associated while, even though non-detrimental, more pronounced colour differences were observed in strain JA757, which may be related to differences in pigment content or energy metabolism in the muscle.

Whatever the strain, no difference in thigh ultimate pH (proxy of glycogen reserve) and proximal composition were reported between high and low-ranger, which is in line with previous results^{10,17}. However, when looking at the fatty acid composition, we showed a higher percentage of α -linolenic acid (C18:3 n-6) in the thigh muscle of HR compared to LR of the dual-purpose strain, without impact on the n-6/n-3 ratio. As α -linolenic acid is an omega 3 precursor, there could be a nutritional interest to its increase and, although the pasture fatty acids profile was not investigated, we may hypothesize that α -linolenic acid increase could be due to higher grass ingestion¹¹.

Broilers' blood parameters divergence with range use

Several biomarkers of the metabolism, inflammation, immunity and redox status were measured to assess the relationship between range use and chicken's physiological and health status at slaughter age. Most of the differences between HR and LR were observed in the most productive strains of this study, JA757 and S757N. Indeed, we reported a reduced lysozyme activity in the two strains, coupled with a reduced haptoglobin-like activity in the S757N strain. Lysozyme is an innate immunity protein indicative of acute or chronic inflammation³⁰ and the haptoglobin-like activity is indicative of an inflammatory status³¹. Our results are in accordance with previous results showing lower levels of lysozyme activity in strains spending more time outdoors³² and in rearing systems with outdoor access compared to indoor rearing³³. Whether range use has an immunosuppressing effect or a protective effect on anti-inflammatory status depends on the level of expression and therefore concentration in the serum or plasma but also on the level of expression before access. Our results showed similar levels of haptoglobin-like concentration before range access and a reduced increase of haptoglobin-like concentration in the HR compared to the LR (Supplementary Figure S1). Therefore, we suggest a possible positive impact of range use with lower acute and chronic inflammation in the S757N strains but we cannot draw a clear conclusion regarding the lysozyme concentration as we do not have measurements before range access.

Creatine kinase is a muscle enzyme that plays a role in the production of energy and is closely associated with muscle growth rate and activity³⁴. We observed a reduction in creatine kinase activity with range use in all the strains except the White Bresse. This result seems consistent with the reduction in breast muscle mass observed in high rangers from the JA757 and S757N strains but suggests another physiological determinant for the dual-purpose strain for which no difference on breast muscle mass or yield with range use was found.

Additionally, high range use also reduced blood concentrations of antioxidants. In the strain JA757, total antioxidant status (TAS) and the sum of tocopherols (including tocopherols) decreased with range use, while pro-oxidative substances (H₂O₂) increased. Following similar trends, uric acid concentration, which acts as an antioxidant^{35,36}, also decreased with range use in the strain S757N. Our observations are quite consistent with previous results that showed that locomotor activity (which is likely higher in HR chickens than in LR chickens) of Ross 308 broilers and other slow-growing strains are correlated to higher oxidative indicators and lower antioxidant concentration in the blood^{11,18,37}. Therefore, range use seems to induce an imbalanced redox status in the JA757 strain, characterized by the highest growth rate among the studied strains.

Regarding blood lipids, it is worth noting that although no changes in FA concentration and profile was observed between high- and low-rangers, n-6/n-3 ratio decreased with greater range use in S757N. It has been shown that pasture composition exhibits lower n-6/n-3 ratio and higher tocopherol concentration than diets classically provided to chickens^{11,38}, which can explain changes in n-6/n-3 observed in the

blood of S757N. In the White Bresse strain, an increase in total antioxidant status, triglyceride and n-6/n-3 ratio was observed in high-ranger chickens, which is difficult to explain. Because no difference in triglycerides were observed earlier during rearing (64 days of age) between HR and LR of the White Bresse strain (Supplementary Figure S2), it is difficult to conclude if lipid changes observed at slaughter in this strain are related to differences in range use or as a response to fasting or other stressful conditions that may affect bird metabolism before slaughter.

To summarize our results, we reported in Fig. 3 some of the main relationships evidenced between range use and indicators including performance, health, welfare and meat quality traits. The physiologic response to range use may vary in intensity depending on the strain and therefore not all of the relationships highlighted in Fig. 3 are reported in all of the strains. Especially in the White Bresse strain that do not seem adapt to range use with similar physiologic response than JA757, S757N and the dual-purpose strain. It can be hypothesized that a higher range use, by favouring foraging and nutrients uptake from the range, can improve the ratio between saturated and unsaturated lipids in blood and carotenoids content in meat. Nevertheless, due to the high energy spent by high rangers for locomotor activity, the consequences appear less favourable for body growth and oxidative stress, while leg health is improved. Finally, range use seems to limit the stimulation of the immune and inflammatory systems which could be related to the lower density of birds outdoors. More markers would be needed to refine the relationships between range use and each of the functions studied and complete this simplistic representation.

Conclusions

This study reported that range use could be predicted by early body weight in intermediate-growing strains but not in slow-growing strains. Whereas only the consequences of a greater range use are visible on the body weight of slow-growing strains. Greater range use by chickens leads to a decrease in growth performances for all strains except for the slowest-growing dual-purpose strain. Animal welfare indicators did not differ according to range use. Interestingly, our study reveals that the variations related to range use are more pronounced for the fastest growing strains, with some slight differences observed on tibial bone, muscle metabolism, meat quality (thigh and breast), and on several blood parameters related to redox, inflammatory and immune status, likely related to foraging activity and vegetable consumption on range. Further research could assess the relationship between the physical activity of the free-range chickens, their level of consumption on the range, and the associated physiological and metabolic responses.

Materials and methods

Animals, housing and experimental design

The four slow to medium growing breeds studied were Hubbard JA757 (used for certified chicken production), naked-neck Hubbard S757N (used for Label Rouge production), White Bresse (a local breed used for the French Bresse production) and an experimental dual-purpose breed (developed from a crossing between two laying strains). Their respective growth rates (GR) were 36, 26, 23 and 16 g/d and rearing duration was 71, 85, 106 and 99 days. Two batches of one-day-old chicks (male/female ratio of 1), issued from different hatcheries, were received 15 days apart. They were all vaccinated against coccidiosis, Marek's and Gumboro diseases and infectious bronchitis. Chickens housing, lighting programme and identification is fully detailed in Bonnefous et al. ⁸. Only 100 males per strain were followed by scan sampling. We chose to study males due to a potential favourable effect of this sex for range use ^{39,40}. Range access was provided to all chickens from 36 days of age. Ranges measured 2,500 m² (50 m x 50 m) and were provided with grass and unevenly distributed trees (density of 60 to 76 trees per hectare). We provided water and 100% organic feed (including 3 diets based on soya meal, maize, wheat and sunflower meal with in addition barley, alfalfa and triticale from day 29 to day 57 or with faba beans and triticale in addition from day 57 to the end of the rearing period) and characterized by 15.8 MJ/kg and 20.8% Crude Protein (CP) from day 1 to day 28, 15.6 MJ/kg and 18.7% CP from day 29 to day 57, and 15.3 MJ/kg and 17.4% CP from day 57 to the end of the rearing period) *ad libitum*. All strains had access to the same feed formulated according to the S757N strain nutritional needs, which is commonly used in organic rearing in France.

Measurements

Except for range use indicator, definition of the measurements according to the ontologies is available in Supplementary Table S1.

Individual range use

Individual range use was assessed according to the method validated by Ferreira et al. ³⁹ and fully described in Bonnefous et al. ⁸. Briefly, we reported the distance between the broiler and the poultry house to calculate a distance index per day of observation. We calculated the final distance index (FDI) of each individual by adding the distance index from the first to the last day of scan sampling observation. Based on the FDI of each of the 100 individuals studied per strain, we ranked them and divided them into two extreme groups of 25 birds per

strain. We named the group with the highest FDI 'high-ranger' (HR) and the group with the lowest FDI 'low-ranger' (LR) (Fig. 1). The average difference between the two groups was equivalent to 1.6–1.9 standard deviation of the trait depending on the strain.

Live body weight

Chicken live body weights were measured before range access at 29 days of life, after range access at 57 days of life and at slaughter at 71, 85, 106 and 99 days for the JA757, S757N, White Bresse and dual-purpose breeds, respectively, following a fasting period of 8 hours.

Struggling activity, foot pad dermatitis and hock burn evaluation on the slaughter line

Following the measurement of live body weight, birds were hung on the slaughter line and electrically stunned in a water bath before bleeding. Struggling activity between hanging and stunning was evaluated by two measurements: straightening up (SU) of the body (head over the legs) observed as a binary score equal to 0 when the bird did not try to stand up and 1 otherwise, and the total duration of wing flapping (TDWF)²⁶. After stunning, bleeding and scalding, foot pad dermatitis and hock burns were evaluated by visual inspection with a 3-point system⁴¹.

Carcass traits

Carcass weight was measured 1-day post-slaughter and carcass yield was calculated in relation to live body weight measured at the slaughterhouse before stunning. Left drumsticks of 23, 22, 23 and 25 birds of the JA757, S757N, White Bresse and dual-purpose strain respectively, were deboned. Tibia diameter and length were measured with an e-calliper and bone-breaking strength of the tibia was determined with an Instron Universal Testing Instrument (model 5543, INSTRON, Norwood, USA)⁴².

Weight and quality indicators of breast meat

At 15 min post-mortem, the pH (pH15) was measured in the left *Pectoralis (P.) major* muscle as described in Berri et al.³⁴. Two grams of fresh muscle were immediately homogenized in 18 ml of 5 mM iodoacetate/ 15 M KCl solution. The pH of the homogenate was measured with a portable pH meter (Model ph330i, WTW GmbH, Vienna, Austria) equipped with a combined glass electrode. After carcass dissection 1-day post-slaughter, the right breast muscle (*P. major* and *minor* muscles) weight was measured. The corresponding total breast yield was calculated in relation to live body weight measured at the slaughterhouse before stunning. At 24 h post-mortem, the pH (pHu) was measured by direct insertion of the electrode in the *P. major* muscle (Model ph330i, WTW GmbH, Vienna, Austria). Colour was measured at 24 h post-mortem on the upper ventral side of this muscle by using a spectrophotometer (CM 700 D, Konica Minolta Inc, Tokyo, Japan), applying the CIELAB trichromatic system as lightness (L*), redness (a*) and yellowness (b*) values. Drip loss (DL) of the same muscle between 24 and 72 h post-mortem was measured as described in Le Bihan-Duval et al.⁴³. After weighing, breast muscles were placed in a plastic bag, hung from a hook, and stored at 2°C for 48 h. After hanging, the sample was wiped with absorbent paper and weighed again. The difference in weight corresponded to the DL and was expressed as the percentage of the initial muscle weight. After DL measurement, *P. major* was vacuum-packed and stored at -20°C until further analyses. Following Le Bihan-Duval et al.²⁴, breast muscle was thawed overnight at 4°C, cooked in a water-bath at 85°C for 7 to 10 min depending on the muscle weight to obtain an internal endpoint temperature of 72°C, and cooled in crushed ice for 10 min. The thawing-cooking loss was expressed as a percentage of the fresh muscle weight. The toughness of cooked meat was evaluated by the Warner Bratzler (WB) shear test using an Instron Universal Testing Instrument (model 5543, INSTRON, Norwood, USA).

Weight and quality indicators of thigh meat

Thigh muscle weight was measured after carcass dissection, 1-day post-slaughter. Corresponding ratios were calculated in relation to live body weight measured at the slaughterhouse before stunning. At 24 h post-mortem, the pH (pHu) was measured by direct insertion of the electrode in the muscle. Proximate and fatty acids composition of a mix of all thigh muscles were calculated from chemical measurements on 23, 22, 23 and 25 birds of the JA757, S757N, White Bresse and dual-purpose strains, respectively. Dry matter, protein and lipid contents were analysed as follows: oven method⁴⁴ for moisture and minerals, Kjeldahl copper catalyst method⁴⁴ for protein, and chloroform/methanol procedure of Folch et al.⁴⁵ for total lipid content. Fatty acid composition was determined after transmethylation of lipids⁴⁶ by gas chromatography (Perkin Elmer Autosystem, St Quentin en Yvelines, France). Injector and detector (FID) temperatures were 250°C, and the carrier gas was nitrogen with a head column pressure of 16.5 psi using a capillary column (25 m x 0.22 mm, BPX, SGE, Villeneuve St Georges, France). Finally, methyl esters were identified and quantified by comparison with standards (Sigma, St Quentin Fallavier, France).

Serum and plasma analyses

We collected blood just after stunning from the carotid bloodletting, half in a heparinized tube and the other half in a dry tube. Blood samples were then centrifuged, after one hour in ice or at room temperature respectively for plasma and serum, then kept at -80°C until further analysis. Details on coefficient of variation intra and inter-assay are present in Supplementary Table S2.

Commercial kits (Thermo Fisher Diagnostics SAS, France) were used to determine plasma glucose (mg/L), uric acid (mg/L) and triglyceride (mg/L) concentrations. Total plasma antioxidant activity was determined through total antioxidant status (TAS) measurement (mmol/L) (Randox Laboratories, UK). The activities of serum creatine kinase (CK) and plasma glutathione peroxidase (GPx) were measured with commercial kits (Biolabo, Malzy, France and Randox Laboratories, London, United Kingdom respectively). The plasma haptoglobin-like activity (mg/mL), that increases in response to acute infection or inflammation ⁴⁷, was measured using a commercial kit (Tridelta Development Limited, Maynooth, Ireland). Protocols listed above were used in accordance with the supplier instructions and adapted to the Thermo Scientific Arena 20XT photometric analyzer (Thermo Fisher Scientific, Courtaboeuf, France). The commercial kit (OxiSelect™, Cell Biolabs, San Diego, USA) was used to quantify serum hydrogen peroxide (µM).

The commercial ID Screen® IBV Indirect kit (IDVET, Grabels, France) was used to qualify the vaccine response to infectious bronchitis by ELISA in plasma. The concentration of antibodies specifically targeting infectious bronchitis antigen was evaluated spectrophotometrically at 450 nm and thus calculated and expressed in antibody titres in order to assess the immune system activity.

The detection of protein carbonyl groups in serum was made following the method of Dalle-Donne et al., (2003), using 2,4-dinitrofenilhidrazina (DNPH) as reactive. Briefly, 500 µL of serum was diluted to 1:40 with phosphate-buffered saline (PBS) before two centrifugations at 1,317 g for 5 minutes. The solution was extracted after each centrifugation. One mL of trichloroacetic acid (TCA) was added to the extracted solution before centrifugation at 1,317 g for 3 minutes. The pellets from trichloroacetic acid (TCA) extracts were mixed with 1 mL of 10 mM DNPH in 2 M hydrochloric acid (HCl). Samples were incubated for 1h at room temperature and then centrifuged at 1,317 g for 5 min. To remove unreacted DNPH, supernatants were discarded and the pellets were washed 3 times with 1 mL of ethanol – ethylacetate (1:1, v/v). The pellets were then dissolved in 1.5 mL of 6 M guanidine-HCl and centrifuged as above to pellet insoluble particles. The carbonyl content of the resulting supernatants was evaluated spectrophotometrically at 366 nm using a molar extinction coefficient of 22,000 1/M*cm. Protein concentrations were measured using the Bradford method with Coomassie Brilliant Blue G-250 ⁴⁹, with bovine serum albumin as the standard. Finally, carbonyl proteins values were expressed as nmol of carbonyl/ mg of protein in the guanidine chloride solution.

The tocopherols (i.e., α-tocopherol and its isoforms γ and δ, and α and γ-tocotrienol) levels were measured in plasma according to Schüep and Rettenmaier ⁵⁰. Briefly, 0.5 mL of plasma was mixed with 0.5 mL of ethanol solution, 1 mL of saline solution and 4 mL of hexane / ethanol solution of 0.06% butylated hydroxytoluene (BHT). The mixture was sonicated in a sonication bath at intensity 30 for 10 minutes. Following centrifugation, 3.5 mL of the supernatant was transferred into a glass tube, dried under nitrogen flow, and resuspended in 200 µL of acetonitrile. The solution was reextracted twice and 50 µL of final acetonitrile solution was then injected into the High-Performance Liquid Chromatography with Fluorescence Detection (HPLC-FD) (pump model Perkin Elmer series 200, equipped with an autosampler system, model AS 950 – 10, Jasco, Tokyo, Japan) on a Sinergy Hydro-RP column (4 mm, 4.6 x 100 mm; Phenomenex, Bologna, Italy). The flow rate was 2 mL/min. All tocopherols and tocotrienols were identified using a Fluorescence Detector (model Jasco, FP-1525 - excitation and emission wavelengths of 295 and 328 nm, respectively) and quantified using external calibration curves prepared with increasing amounts of pure standard solutions (Sigma-Aldrich, Bornem, Belgium) in ethanol. The tocopherols were expressed as µmol/L and their sum was used for statistical analysis.

The lipid fraction for fatty acid evaluation was extracted from serum following the method reported by Folch et al. ⁴⁵. To obtain the fatty acid methyl esters, the lipid extract was dried with a rotavapor and 300 µL of n-hexane was added. Finally, the transmethylation procedure was performed with 150 µL of 2 M potassium hydroxide (KOH) methanol solution at 60°C for 10 min, following the method reported by Glass and Christopherson ⁵¹. For fatty acid quantity calculation, heneicosanoic acid was used as the internal standard (C21:0, Sigma-Aldrich analytical standard). To calculate the sum of the total saturated (SFA), total monounsaturated (MUFA), and total polyunsaturated (PUFA) acids from the n-3 and n-6 series, we used the average amount of each fatty acid. We then calculated the n-6/n-3 fatty acid ratio. Fatty acids results were expressed as % of total fatty acids.

Lysozymes within plasma were evaluated following the method of Osserman and Lawlor ⁵². Briefly, 40 µL of serum was distributed in a well within a petri dish containing a micro-organism sensitive to lysozymes' lytic activity (*Micrococcus lysodeikticus*). After an incubation of 18h at 37°C, the diameter of the ring was measured with an e-calliper. The concentration of lysozymes in the plasma was calculated and expressed in µg/ml (or mg/L), using a petri dish with known concentration of lysozymes as reference.

Statistical analysis

In order to evaluate the relationship between range use groups and body performances, meat quality or blood quantitative parameters within each line, we compared the means of the two groups using Student's test or Wilcoxon-Mann-Whitney's test depending on the normality that were evaluated by Shapiro's test. For qualitative parameters, we used a Fisher exact test. We applied a false discovery rate correction to our p-values per category of parameters as presented in the result tables. Significance was held when $P < 0.05$ and only significant differences have been discussed. We carried out our analyses using R version 4.1.2.

Declarations

Ethics declaration

This study was conducted at the experimental unit UE 1206 EASM of INRAE, France (<https://doi.org/10.15454/1.5572418326133655E12>), from February to June 2021 and slaughter was conducted at the experimental unit UE PEAT of INRAE, France (<https://doi.org/10.15454/1.5572326250887292E12>). It received Ethics committee approval by the Ethics Committee COMETHEA Poitou-Charentes (APAFIS#28675-2020120215483186 v3) in agreement with French and European legislation, and was carried out in accordance with relevant guidelines and regulations. This ethics committee is registered by the National Committee under the number C2EA 84. Reporting in the manuscript follows recommendations in ARRIVE guidelines.

Data availability

The data were deposited in an official repository entrepot.recherche.data.gouv.fr/dataverse/inrae. The data-sets generated and analysed during the current study are available on this website: <https://entrepot.recherche.data.gouv.fr/dataverse/inrae>. The data-sets digital object identifier is <https://doi.org/10.57745/JS5HLR>.

Acknowledgements

We are grateful to all members of the INRAE EASM unit, especially the animal caretakers, for their help with animal care. We thank all members of the INRAE PEAT unit and the INRAE BOA unit for their help at the slaughterhouse. We thank Alexandre Barbin and Nina Kruger for their help with data collection, Catherine Hurtaud for her help with ATOL ontology, and Pierre Blavy for his advice on statistical analysis and we are grateful to Julie Collet for her statistical support and reviewing. We also thank the breeding companies for providing 1-day-old chicks.

This study was funded by the PPILOW project and the French National Research Institute for Agriculture, Food, and Environment (INRAE). The PPILOW project has received funding from the European Union's Horizon 2020 research and innovation programme under Grant Agreement No 816172.

Author contributions

C. Bonnefous, A.C., L.R., K.G., S.M-G., V.G., L.C., M.R., S.M., L.A.G., C.C., C. Berri. and E. B-D designed the experiment, C. Bonnefous, K.G., L.R., T.B., P.C., E.G., E.C-A., N.C., E.R., and E.A. participated to data collection, C. Bonnefous analysed the data and wrote the first draft of the manuscript. All authors revised and approved the final manuscript.

Additional information

Competing interests: The authors declare no competing interests.

Correspondence and requests for materials should be addressed to C. Bonnefous or A.C

Author ORCIDs

Bonnefous Claire: <https://orcid.org/0000-0002-3522-5686>

Collin Anne : <https://orcid.org/0000-0002-3410-6108>

Guilloteau Laurence A. : <https://orcid.org/0000-0001-7089-2196>

Germain Karine : <https://orcid.org/0009-0005-6638-9404>

Ravon Laure : <https://orcid.org/0009-0009-7339-0875>

Bordeau Thierry : <https://orcid.org/0009-0005-0986-9647>

Chartrin Pascal : <https://orcid.org/0009-0003-0437-6195>

Godet Estelle: <https://orcid.org/0009-0001-4884-462X>

Cailleau-Audouin Estelle: <https://orcid.org/0009-0004-2403-4907>

Couroussé Nathalie: <https://orcid.org/0009-0004-3501-5330>

Raynaud Emilie :

Sandrine Mignon-Grasteau: <https://orcid.org/0000-0002-0142-0109>

Guesdon Vanessa: <https://orcid.org/0000-0003-3806-6163>

Calandreau Ludovic: <https://orcid.org/0000-0002-7535-5733>

Berri Cécile: <https://orcid.org/0000-0002-3613-1825>

Reverchon Maxime: <https://orcid.org/0000-0001-8329-7708>

Mattioli Simona: <https://orcid.org/0000-0001-5063-6785>

Castellini Cesare: <https://orcid.org/0000-0002-6134-0901>

Angelucci Elisa : <https://orcid.org/0000-0002-0040-5257>

Le Bihan-Duval Elisabeth: <https://orcid.org/0000-0002-0712-5549>


References

1. FiBL & IFOAM. The World of Organic Agriculture Statistics and Emerging Trends 2021. 120–123 <https://www.organic-world.net/yearbook/yearbook-2021/yearbook-2021-contents.html> (2021).
2. Pinto da Rosa, P. *et al.* Impact of different chicken meat production systems on consumers' purchase perception. *Br. Poult. Sci.* **62**, 387–395 (2021).
3. El-Deek, A. & El-Sabrou, K. Behaviour and meat quality of chicken under different housing systems. *Worlds. Poult. Sci. J.* **75**, 105–114 (2019).
4. Campbell, D. L. M., Horton, B. J. & Hinch, G. N. Using radio-frequency identification technology to measure synchronised ranging of free-range laying hens. *Animals* **8**, 1–16 (2018).
5. Dawkins, M. S., Cook, P. A., Whittingham, M. J., Mansell, K. A. & Harper, A. E. What makes free-range broiler chickens range? In situ measurement of habitat preference. *Anim. Behav.* **66**, 151–160 (2003).
6. Rodriguez-Aurrekoetxea, A., Leone, E. H. & Estevez, I. Environmental complexity and use of space in slow growing free range chickens. *Appl. Anim. Behav. Sci.* **161**, 86–94 (2014).
7. Ferreira, V. H. B. *et al.* Foraging Behavior Shows Individual-Consistency Over Time, and Predicts Range Use in Slow-Growing Free-Range Male Broiler Chickens. *Front. Vet. Sci.* **9**, 1–12 (2022).
8. Bonnefous, C. *et al.* Behavioural indicators of range use in four broiler strains. *Appl. Anim. Behav. Sci.* **260**, (2023).
9. Carere, C. & Maestriperi, D. Animal personalities: who cares and why? in *Animal Personalities. Behavior, physiology, and evolution* 1–9 (University of Chicago Press, 2013). doi:10.7208/chicago/9780226922065.003.0001.
10. Stadig, L. M. *et al.* Effects of free-range access on production parameters and meat quality, composition and taste in slow-growing broiler chickens. *Poult. Sci.* **95**, 2971–2978 (2016).
11. Mattioli, S. *et al.* Intake of nutrients (polyunsaturated fatty acids, tocopherols, and carotenoids) and storage efficiency in different slow-growing chickens genotypes reared in extensive systems. *PLoS One* **17(11)**, (2022).
12. Durali, T., Groves, P., Cowieson, A. & Singh, M. Evaluating range usage of commercial free range broilers and its effect on birds performance using radio frequency identification (RFID) technology. in *25th Annual Australian Poultry Science Symposium Sydney, Australia*. 103–106 (2014).
13. Taylor, P. S., Hemsworth, P. H., Groves, P. J., Gebhardt-Henrich, S. G. & Rault, J. L. Ranging behavior relates to welfare indicators pre- and post-range access in commercial free-range broilers. *Poult. Sci.* **97**, 1861–1871 (2018).

14. Marchewka, J. *et al.* Associations between welfare and ranging profile in free-range commercial and heritage meat-purpose chickens (*Gallus gallus domesticus*). *Poult. Sci.* **99**, 4141–4152 (2020).
15. Fanatico, A. C., Cavitt, L. C., Pillai, P. B., Emmert, J. L. & Owens, C. M. Evaluation of slower-growing broiler genotypes grown with and without outdoor access: Meat quality. *Poult. Sci.* **84**, 1785–1790 (2005).
16. Almasi, A., Andrassyne, B. G., Milisits, G., Kustosne, P. O. & Suto, Z. Effects of different rearing systems on muscle and meat quality traits of slow- and medium-growing male chickens. *Br. Poult. Sci.* **56**, 320–324 (2015).
17. Batkowska, J., Brodacki, A., Zięba, G., Horbańczuk, J. O. & Łukaszewicz, M. Growth performance, Carcass traits and physical properties of chicken meat as affected by genotype and production system. *Arch. Tierzucht* **58**, 325–333 (2015).
18. Mattioli, S. *et al.* How the kinetic behavior of organic chickens affects productive performance and blood and meat oxidative status: a study of six poultry genotypes. *Poult. Sci.* **100**, (2021).
19. Stadig, L. M., Rodenburg, T. B., Ampe, B., Reubens, B. & Tuytens, F. A. M. Effect of free-range access, shelter type and weather conditions on free-range use and welfare of slow-growing broiler chickens. *Appl. Anim. Behav. Sci.* **192**, 15–23 (2017).
20. Fanatico, A. C., Pillai, P. B., Cavitt, L. C., Owens, C. M. & Emmert, J. L. Evaluation of slower-growing broiler genotypes grown with and without outdoor access: Growth performance and carcass yield. *Poult. Sci.* **84**, 1321–1327 (2005).
21. Foutz, T. L., Griffin, A. K., Halper, J. T. & Rowland, G. N. Effects of increased physical activity on juvenile avian bone. *Trans. ASABE* **50**, 213–220 (2007).
22. Debut, M. *et al.* Behavioural and physiological responses of three chicken breeds to pre-slaughter shackling and acute heat stress. *Br. Poult. Sci.* **46**, 527–535 (2005).
23. Chabault, M. *et al.* Analysis of a slow-growing line reveals wide genetic variability of carcass and meat quality-related traits. *BMC Genet.* **13**, 1–8 (2012).
24. Le Bihan-Duval, E. *et al.* Chicken meat quality: Genetic variability and relationship with growth and muscle characteristics. *BMC Genet.* **9**, 1–6 (2008).
25. Debut, M. *et al.* Variation of chicken technological meat quality in relation to genotype and preslaughter stress conditions. *Poult. Sci.* **82**, 1829–1838 (2003).
26. Berri, C. *et al.* Variations in chicken breast meat quality: Implications of struggle and muscle glycogen content at death. *Br. Poult. Sci.* **46**, 572–579 (2005).
27. Castellini, C., Mugnai, C. & Dal Bosco, A. Effect of organic system on broiler carcass and meat quality. *Meat Sci.* **60**, 219–225 (2002).
28. Aksoy, T., Nariç, D., Önenç, A. & İlaslan Çürek, D. Effects of season, genotype and rearing system on some meat quality traits for broilers raised in semi-intensive systems. *Trop. Anim. Health Prod.* **53**, 1–8 (2021).
29. Dal Bosco, A. *et al.* Effect of range enrichment on performance, behavior, and forage intake of free-range chickens. *J. Appl. Poult. Res.* **23**, 137–145 (2014).
30. Carroll, S. F. & Martinez, R. J. Role of rabbit lysozyme in in vitro serum and plasma serum bactericidal reactions against *Bacillus subtilis*. *Infect. Immun.* **25**, 810–819 (1979).
31. O'Reilly, E. L. & Eckersall, P. D. Acute phase proteins: A review of their function, behaviour and measurement in chickens. *Worlds. Poult. Sci. J.* **70**, 27–43 (2014).
32. Castellini, C. *et al.* Adaptation to organic rearing system of eight different chicken genotypes: Behaviour, welfare and performance. *Ital. J. Anim. Sci.* **15**, 37–46 (2016).
33. Franciosini, M. P. *et al.* Influence of different rearing systems on natural immune parameters in broiler turkeys. *Poult. Sci.* **90**, 1462–1466 (2011).
34. Berri, C. *et al.* Consequence of muscle hypertrophy on characteristics of Pectoralis major muscle and breast meat quality of broiler chickens. *J. Anim. Sci.* **85**, 2005–2011 (2007).
35. Yu, B. P. Cellular defenses against damage from reactive oxygen species. *Physiol. Rev.* **74**, 139–162 (1994).
36. Davies, K. J. A., Sevanian, A., Muakkassah-Kelly, S. F. & Hochstein, P. Uric acid-iron ion complexes. A new aspect of the antioxidant functions of uric acid. *Biochem. J.* **235**, 747–754 (1986).
37. Mattioli, S. *et al.* Adaptive response to exercise of fast-growing and slow-growing chicken strains: Blood oxidative status and non-enzymatic antioxidant defense. *Poult. Sci.* **96**, 4096–4102 (2017).
38. Dal Bosco, A., Mugnai, C., Ruggeri, S., Mattioli, S. & Castellini, C. Fatty acid composition of meat and estimated indices of lipid metabolism in different poultry genotypes reared under organic system. *Poult. Sci.* **91**, 2039–2045 (2012).

39. Ferreira, V. H. B. *et al.* Relationship between ranging behavior and spatial memory of free-range chickens. *Behav. Processes* **166**, 1–8 (2019).
40. Chapuis, H. *et al.* Characterization of organic free range broiler exploratory behaviour obtained through RFID technology. *9ème Journées la Rech. Avic. Tours, Fr. 29-30 Mars, 2011* (2011).
41. Bilgili, S. F., Alley, M. A., Hess, J. B. & Nagaraj, M. Influence of age and sex on footpad quality and yield in broiler chickens reared on low and high density diets. *J. Appl. Poult. Res.* **15**, 433–441 (2006).
42. Rousseau, X. *et al.* Metabolism and nutrition: Adaptive response of broilers to dietary phosphorus and calcium restrictions. *Poult. Sci.* **95**, 2849–2860 (2016).
43. Le Bihan-Duval, E., Millet, N. & Remignon, H. Broiler meat quality: Effect of selection for increased carcass quality and estimates of genetic parameters. *Poult. Sci.* **78**, 822–826 (1999).
44. AOAC. *Official methods of analyses*. (1990).
45. Folch, J., Lees, M. & Sloane Stanley, G. . A simple method for the isolation and purification of total lipids from animal tissues. *J. Biol. Chem.* **226**, 497–509 (1957).
46. Morrison, W. R. & Smith, L. M. Preparation of fatty acid methyl esters and dimethylacetals from lipids with boron fluoride–methanol. *J. Lipid Res.* **5**, 600–608 (1964).
47. Oh, S. K., Pavlotsky, N. & Tauber, A. I. Specific binding of haptoglobin to human neutrophils and its functional consequences. *J. Leukoc. Biol.* **47**, 142–148 (1990).
48. Dalle-Donne, I., Rossi, R., Giustarini, D., Milzani, A. & Colombo, R. Protein carbonyl groups as biomarkers of oxidative stress. *Clin. Chim. Acta* **329**, 23–38 (2003).
49. Bradford, M. M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**, 248–254 (1976).
50. Schüep, W. & Rettenmaier, R. Analysis of Vitamin E homologs in plasma and tissue: High-Performance Liquid Chromatography. *Methods Enzymol.* **234**, 294–302 (1994).
51. Glass, R. L. & Christopherson, S. W. A method for the differential analysis of mixtures of esterified and free fatty acids. *Chem. Phys. Lipids* **3**, 405–408 (1969).
52. Osserman, E. F. & Lawlor, D. P. Serum and urinary lysozyme (muramidase) in monocytic and monomyelocytic leukemia. *J. Exp. Med.* **124**, 921–952 (1966).


Figures



Strain	JA757		S757N		White Bresse		Dual-purpose	
	LR	HR	LR	HR	LR	HR	LR	HR
Max	40.0	1207.5	160.0	765.0	507.5	2449.0	92.5	907.5
Min	0.0	295.0	5.0	427.5	137.3	1105.3	0.0	247.5
Mean	16.4	557.2	70.8	580.7	304.9	1412.6	28.7	457.4
Median	15.0	442.5	70.0	577.5	307.5	1375.3	20.0	420.0

Figure 1

Boxplots showing the FDI of JA757 strain (n = 25 per range use group), S757N strain (n = 25 per range use group), White Bresse strain (n = 25 per range use group) and dual-purpose strain (n = 25 per range use group) indicating minimum, 25th percentile values of FDI, median, 75% percentile values of FDI, maximum of FDI of broilers.



Strain	JA757				S757N				White Bresse				Dual-purpose											
	D29		D57		D71		D29		D57		D85		D29		D57		D99							
Range use	LR	HR	LR	HR	LR	HR	LR	HR	LR	HR	LR	HR	LR	HR	LR	HR	LR	HR						
	Max	811	761	2631	2254	3440	3085	639	617	1923	1881	3230	2765	574	581	1657	1633	3073	2934	408	364	1036	914	1835
Min	382	343	1748	1412	2205	1980	350	357	1058	1061	1720	2030	358	322	1205	917	2369	2115	202	234	511	464	1212	1202
Mean	602	521	2221	1851	2840	2543	507	504	1590	1485	2517	2375	472	457	1465	1367	2736	2566	277	298	713	719	1593	1653
Media	.2	.0	.5	.2	.4	.4	.8	.6	.5	.0	.4	.8	.8	.5	.9	.4	.6	.2	.7	.3	.6	.4	2	1
n	.0	.0	.0	.0	.0	.0	.0	.0	.5	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	5	0

Figure 2

Boxplots showing the BW of (a) JA757 strain (n = 25 per range use group), (b) S757N strain (n = 25 per range use group), (c) White Bresse strain (n = 25 per range use group) and (d) dual-purpose strain (n = 25 per range use group) at three weighing times (29 days old, 57 days old and slaughter age), indicating minimum, 25th percentile values of BW, median, 75% percentile values of BW, maximum of BW of broilers depending on their range use indicated by no pattern for low-rangers and patterns with stripes for high-rangers and the p-value after t-test to compare range use categories by age for each strain.




Figure 3

Graphical conclusion summarizing the potential relationships that were outlined in our study. In black are the observed results in our study and in blue are the suppositions.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryFigureS1.pdf](#)
- [SupplementaryFigureS2.pdf](#)
- [SupplementaryTableS1.docx](#)
- [SupplementaryTableS2.docx](#)