



HAL
open science

The Givetian euphyllphyte assemblage from Oum el Jerane, eastern Anti-Atlas, Morocco

Brigitte Meyer-Berthaud, Candice Bert, Anne-Laure Decombeix, Marion Lacan, Merlin Ramel

► To cite this version:

Brigitte Meyer-Berthaud, Candice Bert, Anne-Laure Decombeix, Marion Lacan, Merlin Ramel. The Givetian euphyllphyte assemblage from Oum el Jerane, eastern Anti-Atlas, Morocco. 8th International Agora Paleobotanica Meeting, Josep Sanjuan (Universitat de Barcelona, IRBio); Aixà Tosal (Universitat de Barcelona, IRBio); Carles Martın-Closas (Universitat de Barcelona, IRBio), Sep 2023, Bellver de Cerdanya, Spain. hal-04209948

HAL Id: hal-04209948

<https://hal.inrae.fr/hal-04209948v1>

Submitted on 18 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinee au depot et a la diffusion de documents scientifiques de niveau recherche, publies ou non, emanant des etablissements d'enseignement et de recherche francais ou etrangers, des laboratoires publics ou prives.



Distributed under a Creative Commons Attribution 4.0 International License

The Givetian euphyllophyte assemblage from Oum el Jerane, eastern Anti-Atlas, Morocco

Brigitte MEYER-BERTHAUD¹, Candys BERT¹, Anne-Laure DECOMBEIX¹, Marion LACAND², Merlin RAMEL¹

1. UMR AMAP, Univ. Montpellier, CNRS, CIRAD, INRA, IRD, Montpellier, France
2. Research laboratory on cold forests, UQAT, Canada

The Middle Devonian is a pivotal period, marking a major transition from the early land plants that are now extinct to the structurally complex plants that gave rise to the major vascular plant groups of today. Our current understanding of the patterns and dynamics of this transition are mainly based on data from the paleocontinent Laurussia (Capel et al. 2022). When, at what rate, and how did this transition occur in Gondwana, the largest but least sampled paleocontinent for this time period? Our team is involved in a long-term project to document the taxonomic composition of Devonian plant assemblages from the northern edge of Gondwana and help answer this type of questions.

This talk focusses on the anatomically preserved euphyllophytes discovered at Oum El Jerane, a new Givetian (Middle Devonian) plant locality of Anti-Atlas in southeastern Morocco. The fossiliferous beds are marine and dated from the upper Givetian by the invertebrate fauna. The specimens consist of limonitized fragments of axes measuring 4-14 mm wide and up to 30 mm in length

One of the best preserved specimens is a last-order branch of *Arachnoxylon minor* (Iridopteridales). The Cladoxylopsida are represented by two specimens corresponding to relatively small plants, a branch referable to cf. *Pseudosporochnus hueberi* and the proximal portion of a stem bearing roots assignable to a new genus. The aneurophytalean progymnosperms are represented by woody axes of *Triloboxylon* cf. *arnoldii* and a last order branch bearing paired appendages suspected to belong to a new species of *Triloboxylon*.

The generic composition of the Oum El Jerane assemblage is similar to that of the well-known assemblages from New York (Beck & Wight, 1988) indicating that floral exchanges occurred between northern Gondwana and Laurussia. Differences occur at the specific level. They suggest a certain level of provinciality of the Moroccan flora.

References

- Beck, C.B. & Wight, D.C, 1988. Progymnosperms. In *Origin and Evolution of gymnosperms*. C.B. Beck (Ed). Columbia University Press, New York, 1-84.

Capel, E. Cleal, C. J. Servais, T. & Cascales-Miñana, B, 2023. New insights into Silurian–Devonian palaeophytogeography. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 613, 111393.