



**HAL**  
open science

## Designing sustainable agrosystems by copying the biogeochemical organization of natural ecosystems

Sébastien Fontaine, Luc Abbadie, Gaël Alvarez, Michaël Aubert, Sébastien Barot, Juliette Bloor, Delphine Derrien, Olivier Duchene, Nicolas Gross, Ludovic Henneron, et al.

### ► To cite this version:

Sébastien Fontaine, Luc Abbadie, Gaël Alvarez, Michaël Aubert, Sébastien Barot, et al.. Designing sustainable agrosystems by copying the biogeochemical organization of natural ecosystems. EJP SOIL Annual Science Days 2023, EJP SOIL, Jun 2023, Riga, Latvia. hal-04217071

**HAL Id: hal-04217071**

**<https://hal.inrae.fr/hal-04217071>**

Submitted on 2 Oct 2023



**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Designing sustainable agrosystems by copying the biogeochemical organization of natural ecosystems


S. Fontaine, L. Abbadie, M. Aubert, S. Barot, J.M.G. Bloor, D. Derrien, O. Duchene, N. Gross, L. Henneron, X. Le Roux, N. Loeuille, J. Michel, S. Recous, D. Wipf, G. Alvarez

# Cropping systems: can get out of the issue of tradeoffs ?



# Old forest & prairie multifunctionality


Yield (T ha<sup>-1</sup>)




Jenkinson et al 1994

LTE Rothmasted

No fertilization  
Continuous exportation




Harvested N (kg N ha<sup>-1</sup>)




Glover et al 2010

Study of pair sites (Kansas)



# Old forest & prairie multifunctionality

Soil C stock (g C kg<sup>-1</sup> soil)




Balesdent et al., 1988

## Soil inventories under different land uses



Long-term C storage (kg C m<sup>-2</sup>)




Schlesinger 1990

## Chronosequence study



# Old forest & prairie multifunctionality

Nitrate leaching ( $\text{kg N ha}^{-1}$ )




Mesocosm study with lysimeters



Cros 2019

Ecosystem multifunctionality  
results from a better  
organisation of species and  
functions




Ecosystem performances depends on the synchrony between plant N demand and soil N supply

**A high level of synchrony**


- **↑** biomass production by **↓** N limitation
- **↓** excess of soluble N, N losses (<5%)
- **↑** nutrient and C retention in SOM



# Need of an organization...provided by 4 systems




Soil organic nutrient reserves



Nutrient retrieved from Atmosphere and soil minerals

Common nutrient market


Time flies...come to visit my POSTER !



Designing productive,  
sustainable ecosystems requires  
to improve the efficiency of the  
system as a whole

(not specific functions)

# Step 1 : characterizing the local pedoclimatic content





Soil texture, pH  
Soil mineral reactivity



Evapotranspiration  
Rainfall  
Extremes temperatures

Identify the more adapted organization systems



# Step 2 : installing the organization systems in agrosystems

On the POSTER, a magic table suggests how to install the targeted systems in agrosystem

Let's continue around a Riga Black Basalm !

Synchrony	Conditions of synchrony	Combination of practices to set up for promoting the targeted synchrony
Sync-MAOM	<ul style="list-style-type: none"> <li>-Acquisitive plant species</li> <li>-Continuous activity of microbes M &amp; I</li> <li>-Reserve of MAOM in soil</li> </ul>	<ul style="list-style-type: none"> <li>- Use or genetically select acquisitive species with high C rhizodeposition rate</li> <li>- The C/N of plant species or organic residues must be high enough. Ideally, the different plant species have contrasting C/N ratios (a, c, j, k)</li> <li>- Maintain a continuous cover of active plants fueling microbes in energy (all pictures but e)</li> <li>- Recycle organic nutrients at local scale (farm-watershed) to preserve soil organic reserve on the long-term (d, e, f)</li> </ul>
Sync-FreeOM	<ul style="list-style-type: none"> <li>-Conservative plant species</li> <li>-Mycorrhizal fungi</li> <li>-Reserve of FreeOM in soil</li> </ul>	<ul style="list-style-type: none"> <li>- Use or genetically select conservative species producing recalcitrant litter with reactive compounds (e, f)*</li> <li>- Amendment of recalcitrant organic residues harboring reactive compounds partly charged in organic nutrients (e)</li> <li>- Recycle organic nutrients at local scale (farm-watershed) to preserve soil organic reserve (d, e, f)</li> </ul>
Sync-Inorganic	<ul style="list-style-type: none"> <li>-Plant symbiosis with mycorrhizal fungi &amp; N<sub>2</sub> fixing bacteria</li> <li>-Nutrients stored in bedrock, soil minerals and/or precipitates</li> </ul>	<ul style="list-style-type: none"> <li>- Use or genetically select of species with strong capacity of mobilizing nutrients from rock and soil minerals</li> <li>- Use of plant with deep roots colonizing bedrock (g, h, i)</li> <li>- Use of legumes (a, c, k)</li> <li>- Inoculation with mixed mycorrhizal fungi &amp; N<sub>2</sub> fixing bacteria in highly degraded soils</li> </ul>
Sync-Market	<ul style="list-style-type: none"> <li>-Plant species with complementary nutritional needs</li> <li>-Common mycorrhizal networks</li> </ul>	<ul style="list-style-type: none"> <li>- Mix plant species with different nutrient acquisition strategies and carbon:nutrient ratios (a, c, l, j, k)</li> <li>- Promote perennial plants (f, g, h, l, k) and/or permanent plant cover (all pictures but e) to fuel mycorrhizae in energy-rich carbon</li> <li>- No or limited use of soil tillage (b) and pesticides to preserve mycorrhizae networks</li> <li>- Inoculation with mixed mycorrhizal fungi in highly degraded soils</li> </ul>
Increasing overall synchrony	<ul style="list-style-type: none"> <li>-Synchrony systems adapted to pedoclimatic context</li> <li>-Complementary synchrony systems</li> <li>-Plant plasticity &amp; reserve</li> </ul>	<ul style="list-style-type: none"> <li>- Analyzing the soil profile and climate, adapt the proportion of the different synchrony systems accordingly</li> <li>- Mixing plant species with different nutrient acquisition strategies (a, c, l, j, k)</li> <li>- Select crops species for their suitability to association</li> <li>- Promote perennial plants with high reserve and organ plasticity (f, g, h, l, j, k)</li> </ul>

