



HAL
open science

Genome-Scale CRISPR-Cas9 Knockout Screening of genes involved in pluripotency of Rabbit induced Pluripotent Stem Cells.

Hong Thu Pham, Zhang Zhirong, Nathalie Beaujean, Sylvie Rival-Gervier, Romeo Ricci, Marielle Afanassieff, Pierre Savatier

► **To cite this version:**

Hong Thu Pham, Zhang Zhirong, Nathalie Beaujean, Sylvie Rival-Gervier, Romeo Ricci, et al.. Genome-Scale CRISPR-Cas9 Knockout Screening of genes involved in pluripotency of Rabbit induced Pluripotent Stem Cells.. 2nd meeting StemPhase, Jun 2022, Rennes (Campus de Beaulieu), France. hal-04217476

HAL Id: hal-04217476

<https://hal.inrae.fr/hal-04217476>

Submitted on 25 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Poster + Flash-talk

Hong Thu Pham

Genome-Scale CRISPR-Cas9 Knockout Screening of genes involved in pluripotency of Rabbit induced Pluripotent Stem Cells.

Hong Thu PHAM¹, Zhirong ZHANG², Nathalie BEAUJEAN¹, Sylvie Gervier¹, Roméo RICCP², Marielle AFANASSIEFF¹, and Pierre SAVATIER¹.

¹ Institut Cellule souche et Cerveau, INSERM U1208, INRAE USC1361, Université Claude Bernard Lyon1, 69675 BRON Cedex

² IGBMC, CNRS UMR 7104, INSERM U1258, 67404 Illkirch CEDEX

Pluripotency refers to the ability of a stem cell to give rise to all cell types in mature organisms. Two types of pluripotent stem cells (PSCs) could be generated in vitro: embryonic stem cells (ESCs) derived from the inner cell mass of early embryos and induced pluripotent stem cells (iPSCs) resulted from reprogramming differentiated cells by overexpression of pluripotency factors. In mice, iPSCs could be stabilized in two pluripotency states depending on culture conditions: the naïve state sustained by the LIF/gp130/Stat3 pathway and the primed state supported by FGF2 and Activin A/TGFb/Smad pathways. Only mouse iPSCs in the naïve state are able to colonize recipient embryos and give rise to chimeric animals. In rabbits, only primed iPSCs have been generated and until now, attempts to reprogram them to the naïve state have failed. We decide to use of Genome-Scale CRISPR-Cas9 Knockout Screening (GeCKO), a powerful tool for discovering gene functions, to find the genes that hinder or promote the production of naïve rabbit iPSCs. With this method, we will target more than 14,000 genes on the rabbit induced pluripotent stem cell line, named B19, using a GeCKO library including 84,400 single guide RNAs. For preliminary steps, a DOX-inducible Cas9 protein expression system was transfected into B19 cells. Then, we used a lentiviral delivery of gRNAs to knockout the *Kdm4a* as a testing gene to improve the system and set-up the GeCKO screening. The B19-cas9#12 cells will now be transduced with the Gecko library. After Cas9 induction and culture under naïve state conditions, cells will be sorted into two populations based on specific expression of two naïve markers (EOS-GFP and FOLR1): the naïve one selected on high expressions of EOS-GFP and FOLR1 and the primed population selected on low expressions of EOS-GFP and FOLR1. Amplification and sequencing of Cas9-targeted genes in these two cell populations will give us signaling pathways to induce or repress to reprogram primed rbiPSCs towards the naïve state.