



**HAL**  
open science

## Farming Systems and egg production

Joël Gautron

► **To cite this version:**

Joël Gautron. Farming Systems and egg production. Master. Master 2 Européen Global Quality, Clermmont-Ferrand, France. 2020. hal-04217782

**HAL Id: hal-04217782**

**<https://hal.inrae.fr/hal-04217782>**

Submitted on 26 Sep 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# Farming systems and egg production

The chicken or the egg ?


Joël GAUTRON  
[Joel.gautron@inrae.fr](mailto:Joel.gautron@inrae.fr)

Directeur de recherche  
Research Director  
UMR BOA  
(33) 2 47 42 75 40



# The chicken egg

The egg, a basic ingredient for food  
Isolated chamber for embryo development




**Must contains the entire components necessary for embryo**

- Well-balanced nutritious ingredients
- Lot of compound (> 1000) with a broad range of biological activities
- Protective systems (natural defenses)

Physical defense (Mainly shell)

Chemical defense (Proteins with antimicrobial activities)

# The egg composition is well adapted for embryo development




## The well adapted composition of the yellow

- *Feminal gamete*  
(Clear disk of 3.5 mm)
- *Nutritional reserves (lipids, proteins) and defenses (antibodies)*
- *Surrounded by a thin and translucent membrane*

# The egg composition is well adapted for embryo development

The egg white, water rich (88%), proteins and glucides

- *Various textures of white*
- *Antimicrobial molecules*



The well adapted composition of the yellow

- *Feminal gamete*  
(Clear disk of 3.5 mm)
- *Nutritional reserves (lipids, proteins) and defenses (antibodies)*
- *Surrounded by a thin and translucent membrane*


# The egg composition is well adapted for embryo development

The egg white, water rich (88%), proteins and glucides

- *Various textures of white*
- *Antimicrobial molecules*

Chalazae to maintain the egg in suspension

*Protection of the yolk from chocks*



Mineral eggshell

- *Ensure the physical protection*
- *Assure la protection thermique*
- *Assure les échanges gazeux*
- *Source de calcium pour l'embryon*

The well adapted composition of the yellow


- *Feminal gamete*
- *(Clear disk of 3.5 mm)*
- *Nutritional reserves (lipids, proteins) and defenses (antibodies)*
- *Surrounded by a thin and translucent membrane*




# The egg as food product



# The egg's journey





Catégorie A (œufs coquille)  
 Catégorie B (ovoproduits)  
 Catégorie C (Pet food)



Réhault et al., 2019; Nys et al., 2017

# Distribution of total French consumption




ITAVI d'après SSP, Kantar et Douanes



# Shell and yolk colours

Shell colour is only dependent of genetic



Brown



Leghorn



Marans

Yolk colour depends of diet carotenoids



alfalfa



Corn



marigold



Paprika

No impact on flavours and taste (but important for consumers)

# Egg as nutritional ingredient for humans

## Nutritional characteristics for 2 eggs (100 g)

- **Calories : 155**
- **Total proteins: 12,3 g**
  - High quality biological value (reference WHO 100/ Cow milk 86)
- **Total lipids: 11,9 g**
  - phospholipids rich: 31 % (soit 3,4 g)
  - majority of unsaturated fatty acids
  - cholesterol : 0,42 g (1,2 g / 100 g de jaune)
  - High digestibility value : 98% Triglycerids, 90% Phospholipids
- **Vitamins rich:**
  - A,D,E, B1, B6, B12, biotine (jaune), B2, folic acid, niacine (white)
- **Minerals:** phosphorus, iron and suffur


# Egg as nutritional ingredient for humans

## Egg and Cholesterol


Can we eat eggs every day?

**Yes**, studies have shown that if cholesterol levels are normal, you can eat many eggs a day without affecting cholesterol levels.

If your cholesterol level is high, you should reduce your intake to 4 per week by cutting down on other sources of animal protein and fats.



26 mars 1984



19 juillet 1999



# Egg as nutritional ingredient for humans

## → Yellow to emulsify

- An emulsion is an intimate mixture of two immiscible liquid substances
- The lecithin in the egg yolk is used as an emulsifier in the preparation of sauces in the kitchen.



## → White is swelling

- Beating the whites to snow means introducing air bubbles into a liquid mixture of water and protein to create a foam. This operation is called foaming.
- These properties vary during storage (gas exchanges between the inside and outside of the egg).



# Egg as nutritional ingredient for humans



**Biscuiterie/ pâtisserie/  
viennoiserie**

Colorant, liant,  
coagulant, moussant

Moussant, foisonnant,  
anti-cristallisant

Emulsifiant, colorant,  
Brillance (dorure)

**Confiserie**

Anti-cristallisant  
Foisonnant

Pouvoir aromatique

**Glaces**

Liant

Emulsifiant

**Charcuterie (quenelle)**

Liant, émulsifiant

**Pâtes alimentaire**

Colorant, liant,  
pouvoir aromatique

**Mayonnaise/sauces  
chaudes**

Agent de texture

Emulsifiant, agent de  
texture






# Egg production

# Chicken eggs, products for human nutrition

**Chickens cross selected for egg production**



**More than 300 eggs produced in one year of production, i.e. ten times the weight of the hen.**

**Transformation of plant matter into animal products = an enormous metabolic challenge!**

# Hystory of egg production

- Before the war: Domesticated chickens => mostly self-consumption
- After the war: need to meet the demand (in quantity) and to control the sanitary conditions (in quality: zero risk) => confinement and breeding in cages.
- 80 90s => "productive egg".

# Eggs and chicken strains



## Table eggs

A basic ingredient for human consumption



## Hatching eggs

Close and self sufficient chamber to allow the development of the chicken  
(Breeders for layers and meat production)

**Layer strains**  
(more than 300 eggs per year)  
Non marketable for meat


**Meat strains**  
(150 eggs per years)  
Non marketable for eggs

# Eggs and chicken strains



## Table eggs

A basic ingredient for human consumption



Itavi d'après FAO, Commission et FranceAgriMer

# Hystory of egg production

- Before the war: Domesticated chickens => mostly self-consumption
- After the war: need to meet the demand (in quantity) and to control the sanitary conditions (in quality: zero risk) => confinement and breeding in cages.
- 80 90s => "productive egg".
- Since the end of the 90s: new consumer demands: strong awareness of citizens on agricultural production systems in general and animal production in particular, including poultry and eggs diversification of farming methods
- The current European production models are the result of this social demand Welfare Directive for laying hens (1999/74/EC).
- This regulation is also the result of scientific research to satisfy the 5 freedoms of animal welfare: no hunger, no thirst, free of movement, no fear/distress, while allowing the expression of natural behaviour.

# Eggs and layers

## Health, economic and ethical issues



### Table eggs

14,7 billions eggs per year in France

A basic ingredient for human consumption

- ✓ Risks of toxi-infections for the consumer (Salmonellosis)
- ✓ Economic losses (about 8%): downgraded eggs linked to degraded egg qualities (dirty, cracked or broken shells, poor internal qualities leading to problems of white/yolk separation)



### Hatchery eggs

1,1 billion eggs per year in France

Close and self sufficient chamber to allow the development of the chicken

- ✓ 10% clear eggs (absence of fertilization, embryonic mortality)
- ✓ Elimination of male chicks from the laying strain (early in ovo sexing)

# Egg production system in Europe

## Welfare Directive for laying hens (1999/74/EC)

### Rules for rearing of hens

Enriched cages

Alternative systems

- The animals must be inspected at least once a day.
- The presence of perches is mandatory. If no minimum height is specified, hens must be able to put their fingers underneath.
- The presence of a nest is required. The nest is a separate space whose floor is not made of wire mesh. This nest can be provided for one or more hens.
- The nest is not considered a usable surface.
- The light program must follow a 24-hour rhythm. An uninterrupted period of darkness of an indicative duration of approximately 8 hours must be practiced in order to allow the animals to rest and to avoid eye problems. Light intensity must be sufficient to allow the animals to see and be seen, especially by the breeder during daily inspection.



# Egg production system in Europe

**Welfare Directive for laying hens (1999/74/EC)**

**Rules for rearing of hens**

**Enriched cages**

**Alternative systems**

**Barn or aviary systems  
Indoor or Outdoor**

**Code 3: Cage fitted with new standards**

**Code 2: Raised on the ground or in an aviary without outside access**

**Code 1: Aviary or ground + outdoor access**

**Code 0: Aviary or ground + outdoor access + Organic production**




# Code 3: Enriched cages


All European production of code 3 eggs has been in cages since 2012: laying hen welfare directive (1999/74/EC).

- Cages with an area of at least 2000cm<sup>2</sup>.
- Height of the cage increased
- Installation of perches (15cm /chicken)
- Installation of separate nests
- Scraping and pecking area
- 12 cm feeder/hen
- 100,000 hens in two buildings

before 2012



since 2012



# Code 3: Enriched cages

**Diet: 100% vegetable, minerals and vitamins**

**Cereals (such as wheat, corn...), protein crops (soybeans, field peas, lupin), vegetable oils, vitamins, food supplements (amino acids or synthetic coloring), source of calcium (3.5% instead of 1% in broilers)**





# Code 2: Barn or aviary without outdoor access

- 30,000 hens in a building
- Maximum density in the building 9 hens per m<sup>2</sup>
- Food 100% vegetable, minerals and vitamins (idem code 1)





# Code 1: Aviary or ground + outdoor access

30,000 hens max. per farm.

Maximum density in the building 9 hens per m<sup>2</sup>

With access to an outdoor course (4m<sup>2</sup> per hen, 12 ha of course for 30 000 hens)

Food (100% vegetable, minerals and vitamins)





# Code 1: Label Rouge specificities

## Label Rouge specifications:

- Two buildings of 6000 hens maximum
- Access to an outdoor course (5m<sup>2</sup> per hen, 6 ha for 12 000 hens)
- Food (100% vegetable, minerals and vitamins, 50% minimum of cereals, no additives)





# Code 0: Organic eggs

Maximum 12 000 hens, buildings of 3000 hens maximum

Maximum density in the building 6 hens per m<sup>2</sup>

Access to an outside run (4 m<sup>2</sup> per hen, max 4,8 ha)


Food: 100% vegetable, minerals and vitamins, without synthetic additives 95% minimum of raw materials from AB

Mainly prevention, stimulation of natural defenses. Lists of authorized veterinary drugs if needed



# Evolution of egg production system in France


## Number of hens in France (Thousands)






# Characteristics of various systems

## Durées de ponte (j)




## Nbre œufs pondus/poule




# Characteristics of various systems

## IC



## Taux mortalité %



# Characteristics of various systems



|   | Cage  | Volière ou sol sans parcours | Volière ou sol + parcours | Label Rouge                   | Bio                            |
|---|---|------------------------------|---------------------------|-------------------------------|--------------------------------|
| Accès au parcours   | NON   | NON                          | OUI                       | OUI<br>5m <sup>2</sup> /poule | OUI<br>4 m <sup>2</sup> /poule |
| Densité en bâtiment<br>(nb poules / m2 accessible aux poules)             | 13,3<br>(6 étages)                          | 9,0<br>(2 étages)            | 9,0<br>(2 étages)         | 9,0                           | 6,0                            |
| Mortalité (%)   | 3-4%  | 6-8%                         | 6-8%                      | 6-8%                          | 8-10%                          |
| Taux de poussière dans le bâtiment<br>(santé de l'éleveur et de la poule) | Faible                                      | Fort                         | Fort                      | Moyen                         | Moyen                          |
| Utilisation vaccins et produits véto                                      | Similaire (que lorsque l'animal est malade) |                              |                           |                               |                                |
| Impact environnemental<br>(Bilan carbone)                                 | Faible<br>(IC=2.2)                          | Moyen<br>(IC=2.4)            | Moyen<br>(IC=2.4)         | Moyen<br>(IC=2.6)             | Moyen<br>(IC=2.6)              |
| Impact environnemental<br>Utilisation de terres                           | Faible                                      | Moyen                        | Moyen                     | Moyen                         | Fort                           |

# Characteristics of various systems

|  | Cage              | Volière ou sol sans parcours | Volière ou sol + parcours | Label Rouge | Bio   |
|--|-------------------|------------------------------|---------------------------|-------------|-------|
| Qualité organoleptique (goût) et nutritionnelle            | Pas de différence |                              |                           |             |       |
| Qualité sanitaire (salmonelle)                             | Pas de différence |                              |                           |             |       |
| Coût de production sortie élevage<br>(€ les 100 oeufs)     | 6,41              | 7.35                         | 7,82                      | 8,65        | 13,64 |
| Cout de production base 100                                | 100               | 115                          | 122                       | 135         | 213   |
| Prix de vente consommateur<br>(€ les 6) <i>Kantar 2014</i> | 0,89              | 0,94                         | 1,36                      | 1,87        | 1,96  |
| Prix de vente consommateur base 100                        | 100               | 106                          | 153                       | 210         | 220   |

# Evolution of egg production systems in UE

## Evolution des systèmes de production dans l'UE



Source Commission européenne



**20**  
**HEURES**

**2**

# Discussion

**What is your favorite eggs and why ?**

**And Now ?**  
**Egg in the next decade ?**



# The specialized chicken lines



## Layer hens

(340 eggs per year)  
Meat is not marketable



ZW female chicken



~~ZZ male chicken~~

## Broiler Production

(<150 eggs per year)  
Non marketable low  
quality eggs



6 billions of males are killed every year in the world



Ethical and societal concern

## Alternatives ?




INRAE



# Reduce the number of layers

## ✓ Use of molt cycles




75 % of marketable eggs is the beneficial limit

Second and third laying cycles are possible after molting of the layer

→ Need to induce artificial molt with water and feeding privation not allowed in EU

Research is needed to induce moulting while respecting animal welfare

# Increasing persistency of laying hens



Second and third laying cycles are possible after molting of the layer


→ Need to induce artificial molt with water and feeding privation not allowed in EU

# Increasing persistency of laying hens

Breeding companies claim that they will have developed the « long life » layer, which will be capable of producing 500 eggs in a production cycle lasting 100 weeks by 2020 (Van Sambeek, 2010)

Bain et al., 2016 estimated « than even 25 more eggs per hen could potentially reduce the UK flock, including breeding hens by 2,5 millions birds per annum. »

limited by the health charter in France...



# Dual purpose chickens



## Layers strain

(340 eggs per year)  
Meat is not marketable



## Broiler strains

(<150 eggs per year)  
Non marketable low  
quality eggs



## Crossbreed

Females are reared  
for egg production

Low number of eggs  
Quality ?



Males are reared for  
meat production

Low meat yield  
Different meat texture (consumer  
education)



Unfavorable  
environmental impact

**Need to evaluate the productivity, the quality, the behaviour of animals in various housing systems and various environmental conditions, health and costs**



# IN OVO SEXING

Gender determination before the birth

- ✓ Must be fast (20 000 to 30 000 eggs per hour)
- ✓ Must be cheap
- ✓ Must be precise (98.5 %)
- ✓ Without detrimental consequences on the hatchability and the viability of the chicken
- ✓ Must be done before 9 days of embryonic development to avoid any nociception

**→ A frantic race between states, scientists and industry to offer alternative solutions and hit the jackpot.**

# How to determine sex in ovo


- Destructives and non destructive methods
- Biological approaches
  - Hormonal detection
  - Metabolite marker detection
- Physicochemistry approaches
  - Dimorphic volatile odors between male and females
- Physical and optical approaches
  - FTIR spectroscopy
  - Raman spectroscopy
  - Magnetic resonance imaging
  - Hyperspectral analysis
- Genetic engineering
  - Genome editing

# In ovo sex determination

## Hormonal testing

SELEGGT – Hormonal testing (<http://www.seleggt.com/>) (Allemagne)

- ✓ Test performed at 9 days of incubation
- ✓ Small hole 12 mm in the shell
- ✓ Samples used a patented test to measure the level of Estrone sulfate only present in females



- ✓ Accuracy 97-98%
- ✓ Prototype développé
- ✓ Almost 100 000 eggs already hatched
- ✓ Price 1-3 cents per egg, 7 Euros per pullet
- ✓ Low throughput of approximately 5,000 eggs

**The only available commercial solution, but not enough fast for the entire egg market**

# In ovo sex determination

## Hormonal testing

### EMBREX – Hormonal testing (USA)

- ✓ Test performed at 17 days of incubation
- ✓ Samples used a patented test to measure the level of Estradiol 17B
- ✓ Accuracy 100% ?
- ✓ Prototype not developed
- ✓ Price unknown

## Metabolite markers

### In ovo – Biomarker detection (<https://inovo.nl/solutions/in-ovo-egg-sexing/>) (Netherlands)

- ✓ Use of H NRM spectroscopy
- ✓ Fast 2 sec/egg
- ✓ marketing planned in 2020

# In ovo sex determination

## Physicochemical methods

### Vital farms & Novotrans – odor detection (USA)

- ✓ Test performed at ?? days of incubation
- ✓ Vacuum system to trap the odors
- ✓ Patented in USA 2019, June
- ✓ Accuracy ????
- ✓ Prototype ???
- ✓ Price ???
- ✓ throughput ???

## Genetic engineering

### EggXYt – Genetic alteration (<https://www.eggxyt.com/>) (Israël)

- ✓ Genome editing
- ✓ Feasible the day of lay
- ✓ Fluorescent detection through the shell
- ✓ Transgenic chickens
- ✓ Consumer acceptance ?
- ✓ Accuracy 100 %
- ✓ Price ???

# In ovo sex determination

## Physical methods

### Spectroscopy – Analysis of fluorescence signals (tu-dresden.de) (Allemagne)

- ✓ Test performed at 3 days of incubation
- ✓ DNA quantity is different from male to female (2%)
- ✓ Blood vessels are illuminated and fluorescence is measured at 910 nm
- ✓ Accuracy 90 %
- ✓ No impact on hatchability
- ✓ Prototype ?

### Hypereye – Hyperspectral imaging (Canada)

- ✓ Test performed at the day of lay
- ✓ Hyperspectral method
- ✓ Specific signature using mathematical algorithms
- ✓ Prototype was announced for 2018
- ✓ Throughput of 50 000 eggs per hours
- ✓ 1 to 5 canadian dollars cents per egg



# In ovo sex determination

## Physical methods

### Spectroscopy – hyperspectral imaging- Evonta technology Dresden (Allemagne)

- ✓ Test performed at 14 days of incubation
- ✓ Difference in feather color
- ✓ Use of strain with different feathers depending on sex
- ✓ Accuracy 97 %
- ✓ Prototype ?
- ✓ Late determination 14 days
- ✓ Needs strains with feather differences

### SOO –non invasive physical methods (France)

- ✓ Two complementary technics to predict the sex embryo. Raman hyperspectral and biocaptors
- ✓ Accuracy ??
- ✓ Prototype ?

# How to determine sex in ovo

| Project     | Country     | Method                      | Day of egg incubation | Accuracy % | Price Cents/egg | Number of egg / H | Marketing       | Remarks   |
|-------------|-------------|-----------------------------|-----------------------|------------|-----------------|-------------------|-----------------|---|
| SELEGGT     | Germany     | Hormonal                    | 9                     | 97-98      | 1-3             | 5000              | Yes             | The only available commercial solution, but not enough fast for the entire egg market |
| EMBREX      | USA         | Hormonal                    | 17                    | 100 ?      | ???             | ???               | No              | No prototype, 17 days is too late   |
| In ovo      | Netherlands | Metabolite                  | 9                     | ??         | ???             | ???               | Planned in 2020 | 2 Sec/egg   |
| Vital farms | USA         | Odor detection              | ???                   | ???        | ???             | ???               | ???             |   |
| EggXYt      | ISRAEL      | Genome editing              | 0                     | 100        | ???             | A lot             | Yes             | Transgenic chickens, consumer acceptance  |
| Dresden     | Germany     | Raman                       | 3,5                   | 90         | ???             | ???               | Prototype       | Accuracy to improve   |
| Hypereye    | Canada      | Hyperspectral               | 0                     | 99         | 1-5             | 30 000<br>50 000  | ????            | Large number of eggs, No news since 2018  |
| Evonta      | Germany     | Hyperspectral Feather color | 14                    | 97         | ???             | ???               | ???             | Late determination, Needs strains with feather differences                            |
| SOO         | France      | Raman Biocaptors            | ???                   | ???        | ???             | ???               | ???             |   |
| PPILLOW     | France      | Electromagnetic signature   | ???                   | ???        | ???             | ???               | No              |   |

**Many of them are still in development**

**Research and development must be continued (New methods, Big data, deep learning, artificial intelligence)**

# Alternatives? Take-home messages

**Few alternatives to the culling of male day-old chicks of layer lines:**

- **Increasing of laying period and use of molt cycles to reduce the number of births chickens for renewal**
- **Developing dual purpose chickens, but not for a mass market**
- **Strengthening the development of *in ovo* sexing processes**

**BUT**



- **Today whatever method used, no fast and robust method is operational**
- **A race against time: There is an increased need for research and development before considering the industrial scale**
- **This change will be really challenging for producers with many technical and economical adaptations**

**To conclude**

# The Chicken or the egg ?

This is the question that is poorly formulated

Birds are descendants of dinosaurs



The chicken or dinosaurs ?