



HAL
open science

Rhizoactinobacteria enhance growth and antioxidant activity in Thai jasmine rice (*Oryza sativa*) KDML105 seedlings under salt stress

Kawiporn Chinachanta, Arawan Shutsrirung, Choochad Santasup, Wasu Pathom-Aree, Doan Trung Luu, Laetitia Herrmann, Didier Lesueur, Chanakan Prom-U-Thai

► To cite this version:

Kawiporn Chinachanta, Arawan Shutsrirung, Choochad Santasup, Wasu Pathom-Aree, Doan Trung Luu, et al.. Rhizoactinobacteria enhance growth and antioxidant activity in Thai jasmine rice (*Oryza sativa*) KDML105 seedlings under salt stress. *Plants*, 2023, 12, pp.3441. 10.3390/plants12193441 . hal-04220870v1

HAL Id: hal-04220870

<https://hal.inrae.fr/hal-04220870v1>

Submitted on 28 Sep 2023 (v1), last revised 29 Sep 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Rhizoactinobacteria enhance growth and antioxidant activity in Thai jasmine rice (*Oryza sativa*) KDML105 seedlings under salt stress

Kawiporn Chinachanta^{1,2}, Arawan Shutsrirung¹, Choochad Santasup¹, Wasu Pathom-aree^{2,3}, Doan Trung Luu⁴, Laetitia Herrmann^{5,8}, Didier Lesueur^{5,6,7,8,9} and Chanakan Prom-u-thai^{1,10*}

¹ Department of Plant and Soil Sciences, Faculty of Agriculture, Chiang Mai University, Chiang Mai 50200, Thailand; kawiporn.ch@cmu.ac.th (K.C.); arawan.s@cmu.ac.th (A.S.); choochad.s@cmu.ac.th (C.S.)

² Center of Excellent in Microbial Diversity and Sustainable Utilization Chiang Mai University, Chiang Mai 50200, Thailand

³ Research Center of Microbial Diversity and Sustainable Utilization, Department of Biology, Faculty of Science, Chiang Mai University, Chiang Mai 50200, Thailand; wasu.p@cmu.ac.th

⁴ IPSiM, Univ Montpellier, CNRS, INRAE, Institut Agro, Montpellier, France; doan.luu@cnrs.fr

⁵ Alliance of Bioversity International and Centre International of Tropical Agriculture (CIAT), Asia Hub, Common Microbial Biotechnology Platform (CMBP), Hanoi 10000, Vietnam; L.Herrmann@cgiar.org (L.H.); didier.lesueur@cirad.fr (D.L.)

⁶ Centre de Coopération Internationale en Recherche Agronomique pour le Développement (CIRAD), UMR Eco&Sols, Hanoi 10000, Vietnam

⁷ Eco&Sols, Université de Montpellier (UMR), CIRAD, Institut National de la Recherche Agricole, Alimentaire et Environnementale (INRAE), Institut de Recherche pour le Développement (IRD), Montpellier SupAgro, 34060 Montpellier, France

⁸ School of Life and Environmental Sciences, Faculty of Science, Engineering and Built Environment, Deakin University, Melbourne, VIC 3125, Australia

⁹ Chinese Academy of Tropical Agricultural Sciences, Rubber Research Institute, Haikou, China

¹⁰ Lanna Rice Research Center, Chiang Mai University, Chiang Mai 50200, Thailand; chanakan.p@cmu.ac.th

* Correspondence: chanakan.p@cmu.ac.th; Tel.: +66 053 0944913

Abstract: Salinity is one of the most devastating abiotic stresses which hamper the growth and production of rice. Nine indole-3-acetic acid (IAA) producing salt-tolerant-plant growth-promoting rhizobacteria (ST-PGPR) were inoculated into Thai jasmine rice (*Oryza sativa* L.) variety Khao Dawk Mali 105 (KDML105) seedlings grown under different concentrations of NaCl (0, 50, 100, and 150 mM). ST-PGPR strains significantly promote growth parameters, chlorophyll content, nutrient uptake (N, P, K, Ca, and Mg), antioxidant activity, and proline accumulation in the seedlings under both normal and saline conditions compared to the respective controls. The K⁺/Na⁺ ratio of the inoculated seedlings was much higher than that of the controls, indicating greater salt tolerance. The highest salt-tolerant and IAA producing strain *Sinomonas* sp. ORF 15-23, yielded the highest values of all the parameters, particularly at 50 mM NaCl. The percentage increases in these parameters relative to the controls were ranged from > 90% to 306%. Therefore, *Sinomonas* sp. ORF15-23 was considered a promising ST-PGPR to be developed as bioinoculants for enhancing the growth, salt tolerance, and aroma of the KDML105 rice in salt-affected areas. The environmentally friendly technologies such as ST-PGPR bioinoculants could also support the sustainability of KDML105 geographical indication (GI) products. However, the efficiency of *Sinomonas* sp. ORF15-23 should be evaluated under field conditions for their effect on rice nutrient uptake and growth, including the 2AP level.

Keywords: aromatic rice; climate resilient agriculture; plant growth-promoting actinomycetes; salinity stress mitigation; salt stress alleviation; salt tolerant rhizobacteria; microbial bioinoculants.

1. Introduction

Citation: To be added by editorial staff during production.

Academic Editor: Firstname Last-name

Received: date

Revised: date

Accepted: date

Published: date



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Salt-affected inland soil [mainly halite (NaCl)] is one of the major constraints affecting rice production in Northeastern Thailand. Affected areas included Thung Kula Rong Hai (TKR), a famous area for the production of unique aroma and high grain quality of Khao Dawk Mali 105 (KDML105) rice variety known as Thai Jasmine rice (*Oryza sativa* L.). While saline soils affect approximately 50% of the country's rice-cultivated area [1], the sustainable KDML105 rice production in the TKR areas faces significant challenges due to the increase in soil salinity resulting from lengthened dry seasons from global climate change and the use of chemical fertilizers. The negative impact of excessive salinity includes an imbalance in cellular ionic flux and excessive concentrations of Na⁺, Cl⁻, Mg²⁺, K⁺, and Ca²⁺ ions inside the plant cells, thereby creating oxidative stress through the production of reactive oxygen species (ROS) that impair photosynthesis and cellular metabolism and leading to reductions in plant growth and yield [2,3]. In rice, excess soluble salts in the soil directly affect plant growth-promoting rhizobacteria (PGPR) and reduce yield components, including stand establishment, the numbers of panicles, tillers, and spikelets per plant, floret sterility, individual grain size, and delayed heading [4,5]. However, some groups of rhizosphere microbes, particularly salt-tolerant PGPR (ST-PGPR), can survive in high salt environments due to their ability to cope with osmotic stress; such microbes can improve plant growth as well as plant tolerance to salinity. The protective activities of ST-PGPR are related to their ability to acquire nutrients from the soil, to produce phytohormones and osmoprotectants, and to induce systemic resistance (ISR) [6,7]. Thus, these defense mechanisms could be very helpful for plants in severely saline conditions and thus promote plant growth under normal and stressful environmental conditions. The literature has confirmed of several identified strains in various genera, e.g., *Planococcus*, *Pseudomonas*, *Bacillus*, *Enterobacter*, and *Azotobacter*, that play significant roles in improving crop yield in wheat, rice, maize, and groundnut under salinity stress [8-11]. The use of PGPR has been reported to improve the growth of non-aromatic and aromatic rice; the latter is most preferred by consumers, and the inoculation of PGPR has markedly increased the chlorophyll content, photosynthetic capacity, and growth of rice [12]. The ST-PGPR strain TY0307 exhibited promising ability regarding salt tolerance, proline accumulation, and yield of rice under salt-stress conditions [6,11]. Additionally, rice rhizobacterial strains of *Pseudomonas*, *Enterobacter*, and *Acinetobacter* were reported to produce 2-acetyl-1-pyrroline (2AP), the primary aromatic compound in the rice variety Basmati-370 [13]. The application of locally isolated ST-PGPR strains could be an effective long-term and sustainable solution for rice cultivation in salt-affected soils in the current agricultural systems that must cope with the effect of climate change [14,15]. Using alternative strategies for mitigation of salinity may not be feasible, as they may have negative impacts on the agroecosystems.

The use of ST-PGPR in plant growth and maintenance of plant homeostasis under saline conditions is gaining increased attention as a strategy for solving the problem of salt stress. In the present investigation, we hypothesized that the inoculation of ST-PGPR obtained from the KDML105 rice rhizosphere grown in the TKR region would enhance the growth, salt tolerance, and aroma intensity of the KDML105 variety. Therefore, IAA-producing ST-PGPR strains previously screened by our group [16-17] were used in the present study. Firstly, the 2AP production potential in the culture broth of the ST-PGPR strains was selected [16], and the selected strains were then used to inoculate KDML105 rice seedlings. The objective of the present study was to evaluate the effects of ST-PGPR strains on growth parameters, chlorophyll content, nutrient concentration, antioxidant capacity, and proline concentrations of the rice seedlings germinated under different levels of salinity.

2. Results

2.1. Effect of ST-PGPR inoculation on KDML105 rice seedlings

The inoculation effects of nine ST-PGPR strains on KDML105 rice seedlings grown under normal (0 mM NaCl) and saline conditions (50, 100, and 150 mM NaCl) were evaluated. Uninoculated seedlings at 0, 50, 100, and 150 mM NaCl were considered as control-0 (pure control), control-50, control-100, and control-150, respectively. The term ‘controls’ was used for the uninoculated seedling treatments of each of the salinity levels. These abbreviations for each of the respective control(s) are used throughout the paper. The results showed that the inoculation of ST-PGPR had significant positive effects on growth parameters, chlorophyll content (SPAD units), antioxidant activity (DPPH), and proline concentration of the seedlings both under normal and saline conditions.

2.2. Growth parameters

ST-PGPR strains and salinity had significant interactions that affected both of shoot and root length and dry biomass ($p < 0.0001$) (Table 1). Under normal growing condition (0 mM NaCl), the inoculation of ST-PGPR strains clearly enhanced KDML105 rice seedling growth compared to the control-0, except for strain CRF5-8. Strains *Micrococcus* sp. ORF15-20 and *Sinomonas* sp. ORF15-23 yielded the highest length for shoot and root, while CRF-5-8 showed the shortest shoot and root, similar to those of the control-0 (Figure 1a). Figure 1b shows KDML105 rice seedling growth as affected by the inoculation of ST-PGPR strain *Sinomonas* sp. ORF15-23 under various NaCl concentrations. The nine selected ST-PGPR strains differed significantly in promoting the seedling growth under salinity stress. Among all the tested strains, *Sinomonas* sp. ORF15-23, the most salt-tolerant strain, promoted the highest shoot length and biomass at all levels of NaCl, followed by *Micrococcus* sp. ORF15-20, *Enterobacter* sp. ORF10-12, and *Micrococcus* sp. ORF15-19. In general, these growth parameters were promoted by the inoculation of most of the tested ST-PGPR strains when the NaCl concentration was increased from 0 to 50 mM, but growth declined progressively at concentrations beyond 50 mM. However, on average, the inoculated treatments provided higher seedling biomass than those of their respective controls at all levels of NaCl concentration. The non-IAA-producing strain CRF5-8 gave the lowest shoot and root length, with values slightly lower than its respective controls at each NaCl concentration. In contrast, the least salt-tolerant strain *Burkholderia* sp. CRF16-3 provided the lowest seedling biomass, with values similar to its respective controls at each NaCl concentration.



Figure 1. Growth-promoting effects of ST-PGPR strains inoculation on KDML105 rice seedlings under normal condition (a), and examples of seedlings inoculated with *Sinomonas* sp. ORF15-23 under various NaCl concentrations (b).

Table 1 Growth-promoting effects of ST-PGPR inoculation on shoot length, root length, and dry weight of KDML105 rice seedlings under normal and saline conditions.

ST-PGPR strains	Shoot length (cm)				Root length (cm)				Dry weight (g)			
	NaCl concentrations (mM)				NaCl concentrations (mM)				NaCl concentrations (mM)			
	0	50	100	150	0	50	100	150	0	50	100	150
Controls¹	21.72 d	20.23 d	10.60 d	5.43 d	9.59 cd	10.18 c	5.33 b	3.12 c	0.30 b	0.27 b	0.24 ab	0.16 c
Organic rice farming												
ORF4-13	24.41 bc	23.74 cd	18.25 ab	12.84 a	19.30 a	12.66 bc	5.67 b	3.43 c	0.32 b	0.38 b	0.29 ab	0.25 bc
ORF10-12	25.17 bc	29.51 abc	17.44 abc	14.96 a	10.01 cd	15.26 b	11.11 a	9.10 a	0.32 b	0.4 b	0.28 ab	0.28 ab
ORF15-19	23.48 cd	28.96 abc	9.69 d	7.59 b	9.83 cd	12.78 bc	4.89 b	3.77 c	0.31 b	0.34 b	0.34 a	0.29 ab
ORF15-20	27.42 ab	30.78 ab	15.80 bc	13.24 a	22.03 a	12.80 bc	6.36 b	5.14 bc	0.42 ab	0.39 ab	0.36 a	0.35 a
ORF15-23	29.42 a	32.50 a	18.81 a	15.40 a	19.34 a	20.77 a	11.48 a	6.88 ab	0.54 a	0.58 a	0.32 a	0.27 ab
Conventional rice farming												
CRF 5-8	20.84 d	13.43 e	9.18 d	8.23 b	7.78 d	6.83 c	4.89 b	3.87 c	0.35 b	0.26 b	0.27 ab	0.25 bc
CRF14-15	23.84 cd	24.50 cd	15.84 bc	6.84 b	11.36 bc	12.83 bc	10.55 a	4.09 c	0.32 b	0.27 b	0.30 ab	0.28 ab
CRF16-3	26.65 abc	22.37 d	19.14 a	7.35 b	10.69 cd	11.36 c	11.44 a	3.65	0.36 b	0.28 b	0.22 b	0.20 bc
CRF17-18	25.04 bc	25.00 bcd	15.00 c	7.35 b	14.38 b	13.24 bc	11.62 a	5.25 bc	0.36 b	0.31 b	0.27 ab	0.25 bc
Mean	25.10	25.102	14.98	9.92	13.43	12.87	8.33	4.82	0.36	0.37	0.29	0.26
F-test	*	*	*	*	*	*	*	*	*	*	*	*
% CV	5.76	10.00	8.31	14.34	10.39	11.10	15.48	8.04	7.64	6.90	8.25	5.11
Rhizobacterial isolates		**				**				**		
(A)												
NaCl concentration		**				**				**		
(B)												
A x B		**				**				**		
LSD _(0.01) (AxB)		2.61				2.08				0.08		
% CV		8.56				13.15				17.15		

¹Controls = control-0, control-50, control-100, and control-150, at 0, 50, 100, and 150 mM NaCl, respectively; Mean (n=3).

The average values followed by different letters within the same column were significantly different according to pairwise comparisons using an LSD test ($P \leq 0.01$).

** Significant at the 0.01 probability level.

2.3. Chlorophyll content of KDML105 rice seedlings

Leaf chlorophyll content of all the treatments decreased gradually with the increase of salt concentration, but there were different magnitudes of decrease among the treatments. The chlorophyll contents of the control-50, control-100, and control-150 seedlings were decreased by 3.04, 7.05, and 21.04%, respectively, compared to the control-0. However, the application of ST-PGPR strains enhanced the leaf chlorophyll content of the seedlings by 1.34-18.48%, ~0-16.5%, ~0-15.2%, and ~0-27.2%, at 0, 50, 100, and 150 mM NaCl, respectively, compared to the respective controls (Table 3). The maximum chlorophyll content was obtained with *Sinomonas* sp. ORF15-23 inoculation at all levels of salinity, with the percentage of increase ranging from 15.26-27.50 % compared to the respective controls. Under salinity stress, the inoculation of ST-PGPR strains significantly increased the total chlorophyll content compared to those of the controls (Table 2).

2.4. Antioxidant activity and proline in KDML105 rice seedlings

It appeared that salinity had a promoting effect on the antioxidant activity (DPPH radical scavenging activity) of the leaves of KDML105 rice seedlings, and the effect was significantly enhanced by the inoculation with the ST-PGPR strains. The values of DPPH radical scavenging activity ranged from 43.94 to 60.92 mg Trolox g mL⁻¹ for the uninoculated controls, and from 45.43 to 93.43 mg Trolox g mL⁻¹ for treatment inoculated with ST-PGPR strains. The highest concentration of tested NaCl (150 mM) provided the maximum antioxidant activity in the seedling leaves for each treatment, with values of 60.32 to 93.43 mg Trolox g mL⁻¹. On the average, the rank of the antioxidant activities was observed as *Sinomonas* sp. ORF15-23 > *Micrococcus* sp. ORF15-20 > *Micrococcus* sp. ORF15-19 > *Enterobacter* sp. ORF10-12 > *Sinomonas* sp. CRF14-15 > *Bacillus* sp. CRF17-18 > *Sinomonas* sp. ORF4-13 > *Burkholderia* sp. CRF 16-3 > controls > CRF 5-8 (Table 2).

The proline accumulation in leaves of KDML105 seedlings increased with an increasing NaCl concentration from 0 to 100 mM NaCl and decreased thereafter. Furthermore, the inoculation of ST-PGPR strains significantly increased the proline content of the leaves at all NaCl concentrations compared to the respective controls. The maximum increase in proline content was obtained by *Sinomonas* sp. ORF15-23 inoculation, with percentage increase of 107.8, 163.0, 80.2, and 121.7% at 0, 50, 100, and 150 mM NaCl compared to control-0, control-50, control-100, and control-150, respectively (Table 2).

Table 2. Effect of ST-PGPR inoculation on chlorophyll content, proline accumulation, and antioxidant activity in the KDML105 rice seedling leaves under different NaCl concentrations.

ST-PGPR strains	Chlorophyll (SPAD unit)				Proline ($\mu\text{mol g}^{-1}\text{FW min}^{-1}$)				DPPH radical scavenging activity (mg Trolox g mL ⁻¹)			
	NaCl concentrations (mM)				NaCl concentrations (mM)				NaCl concentrations (mM)			
	0	50	100	150	0	50	100	150	0	50	100	150
Controls	37.33 c	36.23 d	34.87 bc	30.84 c	14.84 d	17.84 f	30.43 d	15.94 d	43.94 f	57.47cd	61.92 e	60.92 d
Organic rice farming												
ORF4-13	39.84 bc	36.24 d	34.95 bc	30.36 c	25.04 b	29.93 cd	38.03 c	21.93 c	48.95df	56.04de	68.93cd	70.32 c
ORF10-12	40.32 bc	39.84 abc	38.74 a	37.72 a	29.05 a	35.29 b	42.93 b	29.84 b	62.04 b	71.52 b	76.34 b	80.43 b
ORF15-19	41.52 ab	40.95 ab	39.95 a	38.87 a	26.94 b	31.94 c	45.92 b	28.92 b	70.32 a	74.06ab	83.94 a	89.32 a
ORF15-20	39.42 bc	39.42abcd	35.39 b	33.28 b	24.95 bc	32.94 bc	45.23 b	30.94 b	69.94 a	74.95 a	82.95 a	89.42 a
ORF15-23	44.23 a	42.19 a	40.19 a	39.32 a	30.84 a	46.92 a	54.83 a	35.34 a	72.94 a	76.04 a	85.03 a	93.43 a
Conventional rice farming												
CRF5-8	37.93 c	36.92 cd	31.94 cd	30.48 c	16.94 c	24.92 e	23.02 e	20.94 c	45.43ef	49.54f	54.03 f	60.32 d
CRF14-15	40.32 bc	38.84 bcd	34.96 bc	32.05 bc	23.94 bc	27.94 d	34.29 cd	20.94 c	56.93 c	60.30 c	69.83 c	73.95bc
CRF16-3	37.83 c	36.93 cd	33.64 bcd	30.59 c	15.94 d	28.94 cd	27.43 de	16.92 d	52.43 d	56.03de	65.47de	69.34 c
CRF17-18	39.82 bc	38.85 bcd	30.50 d	31.93 bc	17.94 c	18.42 f	20.58 f	15.93 d	49.95d	53.05e	62.05 e	70.93 c
Mean	39.86	38.64	35.51	33.54	32.52	36.51	39.78	30.22	57.29	62.90	71.05	75.84
F-test	*	*	*	*	*	*	*	*	*	*	*	*
% CV	3.37	3.62	3.86	2.94	4.31	5.47	4.42	3.72	2.95	2.13	2.18	3.86
Rhizobacteria isolates (A)		**				**				**		
NaCl concentration (B)		**				**				**		
A x B		**				**				**		
LSD _(0.01) for (AxB)		1.99				3.16				1.80		
%CV		3.33				2.91				3.34		

¹Controls = control-0, control-50, control-100, and control-150, at 0, 50, 100, and 150 mM NaCl, respectively; Mean (n=3).

The average values followed by different letters within the same column were significantly different according to pairwise comparisons using an LSD test ($P \leq 0.01$).

** Significant at the 0.01 probability level.

2.4. Nutrient uptake

The relationships between NaCl concentrations, ST-PGPR inoculation, and nutrient uptake (N, P, K, Ca, Mg, and Na) of KDML105 rice seedlings are shown in Table 3. The analysis showed significant interaction ($P < 0.01$) between ST-PGPR strains and NaCl concentrations for nutrient uptake. The shoot N, P, and K uptake decreased in the controls under increasing NaCl concentration, particularly beyond 50 mM NaCl (Table 3). On average, the inoculated treatments provided higher shoot N, P, and K uptake than those of their respective controls. However, the shoot N, P, and K uptake in most of the inoculated treatments also showed a similar trend of negative salt stress effects as in their respective controls, but to a much lesser degree. Among all treatments, strain *Sinomonas* sp. ORF15-23 produced the highest levels of N, P, and K uptake at 50 mM NaCl, with percentage increase of 145.2, 186.2, and 272.6% compared to the control-50 (Table 3).

Compared to their respective controls, the inoculation of most ST-PGPR strains increased shoot N, P, and K uptake of the KDML105 rice seedlings, and the highest NaCl level (150 mM) provided the highest increasing percentage of 41.6-126.2, 69.6-157.8, and 11.2-301.7%, respectively. It was interesting to note that all the CRF strains from conventional rice farming resulted in a lower N uptake than the control-50 (Table 3). Among all the tested strains, only *Sinomonas* sp. ORF15-23 clearly enhanced N, P, and K uptake when the salinity increased from 0 to 50 mM NaCl; however, the uptake decreased progressively beyond 50 mM NaCl.

Table 3. Effect of ST-PGPR inoculation on nitrogen (N), phosphorus (P), and potassium (K) uptake by KDML105 rice seedlings under different NaCl concentrations, and significance level and LSD values for nutrient uptake by KDML105 rice seedlings.

ST-PGPR strains	N				P				K			
	(mg N plant ⁻¹)				(mg P plant ⁻¹)				(mg K plant ⁻¹)			
	NaCl concentrations (mM)				NaCl concentrations (mM)				NaCl concentrations (mM)			
	0	50	100	150	0	50	100	150	0	50	100	150
Controls 1	10.86 d (0.00)	10.26 de (0.00)	6.264 d (0.00)	3.74 d (0.00)	3.63 b (0.00)	3.21 bc (0.00)	2.69 bc (0.00)	1.34 c (0.00)	16.41 c (0.00)	10.26 f (0.00)	8.04 e (0.00)	3.92 c (0.00)
Organic rice farming												
ORF4-13	11.26 d (3.72)	10.03def (-2.22)	9.19 bc (46.76)	8.48 a (126.36)	3.90 ab (7.55)	4.52 bc (40.74)	3.36 abc (25.15)	2.73 ab (102.75)	17.09 c (4.13)	18.09 b (76.30)	11.11 cd (38.15)	9.58 b (144.26)
ORF10-12	11.58 d (6.67)	15.48 b (50.88)	9.58 bc (52.87)	6.80 abc (81.73)	3.87 ab (6.67)	4.76 b (48.15)	2.66 bc (-1.04)	2.44 abc (81.25)	17.34 c (5.67)	14.76bcd (43.86)	9.52 de (18.41)	10.84 b (176.43)
ORF15-19	10.63 d (-2.09)	11.22 cd (9.36)	11.66 a (86.17)	7.92 ab (111.46)	3.75 b (3.33)	3.91 bc (21.69)	3.67 ab (36.61)	2.96 ab (120.09)	16.83 c (2.58)	18.22 b (77.62)	16.63 a (106.79)	15.75 a (301.71)
ORF15-20	19.45 a (79.06)	13.77 bc (34.18)	9.32 bc (48.85)	6.27 abc (67.33)	5.17 ab (42.31)	4.68 bc (45.66)	4.07 a (51.34)	3.47 a (157.81)	17.93 c (9.29)	16.30 bc (58.89)	14.80 ab (84.03)	9.03 b (130.36)
ORF15-23	17.17 ab (58.12)	25.16 a (145.19)	10.02 ab (59.90)	6.94 abc (85.34)	6.59 a (81.49)	9.20 a (186.21)	3.65 ab (35.71)	2.97 ab (116.96)	28.24 a (72.10)	38.23 a (272.59)	12.90 bc (60.40)	9.48 b (127.30)
Conventional rice farming												
CRF5-8	11.83 d (8.93)	8.58def (-16.37)	8.15 c (30.17)	5.30 cd (41.56)	3.57 b (-1.65)	2.96 c (-7.75)	3.35 abc (24.55)	2.63 ab (95.31)	18.06 bc (10.05)	12.45def (21.38)	12.39 c (54.14)	8.91 b (141.71)
CRF14-15	10.72 d (-1.29)	8.15 ef (-20.53)	10.41 ab (66.19)	8.12 a (116.88)	3.94 ab (8.43)	3.24 bc (0.84)	3.36 abc (25.00)	3.19 ab (131.25)	18.53 bc (12.91)	13.85cde (35.00)	12.69 bc (57.84)	11.12 ab (183.67)
CRF16-3	15.08 bc (38.90)	8.85def (-13.76)	5.85 d (-6.58)	5.78 bcd (54.38)	4.36 ab (20.00)	3.36 bc (4.58)	2.38 c (-11.61)	2.28 bc (69.64)	21.10 b (28.56)	10.59 ef (3.16)	8.56 e (6.44)	4.36 c (11.22)

CRF17-18	12.96 cd (19.34)	7.29 f (-29.00)	6.13 d (-2.16)	5.80 bcd (54.91)	4.32 ab (19.01)	3.63 bc (12.89)	3.02 abc (12.50)	3.03 ab (125.07)	17.10 c (4.20)	14.42 cd (40.50)	9.26 de (15.19)	8.68 bc (121.30)
Mean	13.16	11.88	8.66	6.51	4.31	4.34	3.22	2.69	18.86	16.71	11.59	9.84
F-test	*	*	*	*	*	*	*	*	*	*	*	*
% CV	8.15	10.43	8.41	5.03	7.25	7.78	6.42	7.69	8.45	9.13	8.24	2.09
Rhizobacterial isolates (A)		**				**				**		
NaCl concentration (B)		**				**				**		
A x B		**				**				**		
LSD _(0.01) for (AxB)		1.5283				1.2928				2.1961		
%CV		9.35				21.84				30.21		

¹Controls = control-0, control-50, control-100, and control-150, at 0, 50, 100, and 150 mM NaCl, respectively; Mean (n=3).

Numbers in parentheses are percentage increases/ decreases of shoot N, P, and K uptake of the KDML105 rice as compared to their respective controls.

The average values followed by different letters within the same column were significantly different according to pairwise comparisons using an LSD test ($P \leq 0.01$).

Increasing salt concentration resulted in a decreased in Ca and Mg uptake by the KDML105 rice seedlings, both uninoculated and inoculated treatments, except for strain *Sinomonas* sp. ORF15-23 that showed markedly enhanced in Ca and Mg uptake when the concentration increased from 0 to 50 mM NaCl (Table 4). The highest amounts of Ca and Mg uptake were obtained with the strain *Sinomonas* sp. ORF15-23 at 50 mM NaCl, with percentage increases of 306.5 and 204.9% as compared to the control-50. However, beyond this salt level, the uptake decreased monotonically. At the same level of salt concentration, the inoculation of most ST-PGPR strains increased the Ca and Mg uptake of the KDML105 rice seedlings compared to their respective controls (Table 5). On average, the highest tested NaCl level (150 mM) resulted in the maximum percentage increases of Ca and Mg uptake of the inoculated seedlings, with values of 0-184.4 and 25.0-119.4%, respectively. In contrast to Ca and Mg uptake, Na uptake of the controls and the ORF-strains from organic rice farming practice was slightly increased at 50 mM NaCl compared to 0 mM NaCl. Nevertheless, the uptake decreased beyond the concentration of 50 mM NaCl. The inoculation of most ST-PGPR strains increased the Na uptake of the KDML105 rice seedlings compared to their respective controls at the same levels of salt concentration (Table 4).

Table 4. Effect of ST-PGPR inoculation on calcium (Ca), magnesium (Mg), and sodium (Na) uptake by KDML105 rice seedlings under different NaCl concentrations.

ST-PGPR strains	Ca				Mg				Na			
	(mg Ca plant ⁻¹)				(mg Mg plant ⁻¹)				(mg Na plant ⁻¹)			
	NaCl concentrations (mM)				NaCl concentrations (mM)				NaCl concentrations (mM)			
	0	50	100	150	0	50	100	150	0	50	100	150
Controls¹	0.66 b (0.00)	0.49bcd (0.00)	0.29 (0.00)	0.16 b (0.00)	1.68 b (0.00)	1.30 b (0.00)	0.79cde (0.00)	0.48 c (0.00)	9.96 d (0.00)	10.45 cd (0.00)	9.38de (0.00)	6.37 d (0.00)
Organic rice farming												
ORF4-13	0.67 b (1.82)	0.61bcd (25.10)	0.44 (51.04)	0.30 ab (87.50)	1.38 b (-18.10)	1.82 b (40.74)	1.07 b (35.48)	0.73abc (51.04)	10.21 cd (2.51)	13.45 b (28.72)	10.79bcd (15.03)	9.45 b (48.35)
ORF10-12	0.93 b	0.76 bc	0.34	0.020 b	1.57 b	1.72 b	0.76 de	0.70abc	10.08 d	14.24 b	10.052cd	10.11 b

	(40.61)	(56.38)	(16.67)	(22.50)	(-6.67)	(32.71)	(-4.55)	(45.83)	(1.20)	(36.28)	(7.16)	(58.71)
ORF15-19	0.71 b	0.54bcd	0.31	0.26 ab	1.64 b	1.19 b	1.05 b	1.01 ab	9.95 d	12.04 bc	12.27 ab	10.88 b
	(8.03)	(11.93)	(6.25)	(63.13)	(-2.20)	(-8.18)	(33.08)	(111.46)	(-0.10)	(15.23)	(30.81)	(70.80)
ORF15-20	0.84 b	0.82 b	0.58	0.46 a	2.02 ab	1.64 b	1.55 a	1.05 a	13.48 b	13.88 b	13.54 a	13.20 a
	(27.27)	(68.52)	(100.00)	(184.38)	(20.00)	(26.39)	(95.45)	(118.75)	(35.34)	(32.84)	(44.35)	(107.22)
ORF15-23	1.62 a	1.98 a	0.48	0.32 ab	3.13 a	3.95 a	1.09 b	1.05 a	17.01 a	1.630 a	1.130 bc	9.75 b
	(145.45)	(306.58)	(66.67)	(102.50)	(86.43)	(204.94)	(37.37)	(119.38)	(70.78)	(56.00)	(20.47)	(53.06)
Conventional rice farming												
CRF5-8	0.70 b	0.42 d	.030	0.26 b	2.03 ab	1.22 b	0.97 bc	0.75abc	11.24 cd	9.07 d	9.50cde	8.95 bc
	(6.06)	(-14.40)	(3.13)	(40.63)	(20.83)	(-5.71)	(22.73)	(56.25)	(12.85)	(-13.20)	(1.28)	(40.50)
CRF14-15	1.02 ab	0.46 cd	0.42	0.17 b	1.73 b	1.03 b	0.90bcd	1.01 ab	10.56 cd	9.50 cd	10.83bcd	10.42 b
	(55.15)	(-5.56)	(45.83)	(5.00)	(2.86)	(-20.83)	(13.64)	(110.00)	(6.02)	(-9.08)	(15.46)	(63.58)
CRF16-3	0.79 b	0.50bcd	0.22	0.22 b	1.69 b	1.15 b	0.70 e	0.60 bc	11.70 c	10.22 cd	8.14 e	7.36 cd
	(20.00)	(3.70)	(-23.61)	(37.50)	(0.71)	(-11.42)	(-11.11)	(25.00)	(17.47)	(-2.19)	(-13.22)	(15.54)
CRF17-18	0.83 b	0.43 cd	0.22	0.23 b	1.73 b	1.30 b	1.05 b	0.88abc	11.52 cd	10.76 cd	9.50cde	9.03 bc
	(25.45)	(-10.70)	(-25.00)	(40.63)	(2.86)	(0.46)	(32.95)	(82.29)	(15.66)	(2.98)	(1.28)	(41.76)
Mean	0.88	0.70	0.36	0.25	1.86	1.63	0.99	0.83	11.57	12.99	10.53	9.55
F-test	*	*	ns	*	*	*	*	*	*	*	*	*
% CV	3.34	2.94	5.23	3.84	8.53	3.85	8.33	2.18	5.88	8.48	7.77	9.08
Rhizobact												
erial			**			**				**		
isolates												
(A)												
NaCl												
concentrat		**				**				**		
ion (B)												
A x B		**				**				**		
LSD _(0.01)			0.2823			0.5456				1.3728		
for (AxB)												
%CV		31.73				25.28				7.57		

¹Controls = control-0, control-50, control-100, and control-150, at 0, 50, 100, and 150 mM NaCl, respectively; Mean (n=3).

Numbers in parentheses are percentage increases/ decreases of shoot N, P, and K uptake of the KDML105 rice as compared to their respective controls.

The average values followed by different letters within the same column were significantly different according to pairwise comparisons using the LSD test ($P \leq 0.01$).

** Significant at the 0.01 probability level.

One of the various strategies employed by rice to survive under salt stress is maintaining a high K^+/Na^+ ratio in the cells. We hypothesized that the ST-PGPR inoculation might help to promote this ratio, thereby increasing the chance of survival under stress conditions. Therefore, in this experiment, we calculated the K^+/Na^+ ratio in the KDML105 rice seedlings to evaluate the effect of ST-PGPR inoculation. The results indicated that under normal condition (0 mM NaCl), the K^+/Na^+ ratio of the control-0 seedlings (1.65) and inoculated seedlings (1.33-1.80) showed similar or slightly different values (Table 5). However, the K^+/Na^+ ratio of the uninoculated seedlings (controls) showed a marked reduction with increasing NaCl concentrations. The K^+/Na^+ ratio of the ST-PGPR inoculated seedlings was also reduced with increasing NaCl concentrations but to a lesser degree compared to those of the controls. Among all treatments, strain *Sinomonas* sp. ORF15-23 provided the highest K^+/Na^+ ratio at 50 and 100 mM NaCl (Table 5).

Table 5. Effects of ST-PGPR inoculation on K^+/Na^+ ratio in KDML105 rice seedling under different NaCl concentrations.

ST-PGPR strains	K^+/Na^+ ratio
-----------------	------------------

	NaCl concentrations (mM)			
	0	50	100	150
Controls¹	1.65 ab	0.99 c	0.86 b	0.62 ef
Organic rice farming				
ORF4-13	1.67 ab	1.34 ab	1.03 ab	1.01 cd
ORF10-12	1.72 ab	1.04 c	0.95 b	1.07 c
ORF15-19	1.69 ab	1.51 a	1.35 a	1.45 b
ORF15-20	1.33 b	1.17 bc	1.09 ab	0.69 def
ORF15-23	1.66 ab	2.34 a	1.49 ab	0.91 cde
Conventional rice farming				
CRF5-8	1.61ab	1.37 ab	1.30 a	1.06 a
CRF14-15	1.75 a	1.45 a	1.17 ab	1.06 c
CRF16-3	1.80 a	1.04 c	1.05 ab	0.59 f
CRF17-18	1.48 ab	1.34 ab	0.98 b	0.96 cd
Mean	1.64	1.27	1.10	0.94
F-test	*	*	*	*
% CV	10.60	10.32	13.366	13.75
Rhizobacterial isolates (A)			**	
NaCl concentration (B)			**	
A x B			**	
LSD _(0.01) for (AxB)			4.81	
%CV			11.84	

¹Controls = control-0, control-50, control-100, and control-150, at 0, 50, 100, and 150 mM NaCl, respectively. 253

Numbers in parentheses are the percentage increases/ decreases of shoot N, P, and K uptake of the KDML105 rice as compared to their respective controls. 254

The average values followed by different letters within the same column were significantly different according to pairwise comparisons using the LSD test ($P \leq 0.01$). 255

** Significant at the 0.01 probability level. 256

2.5. Relationship between the study variables by principal component analysis 257

The principal component analysis (PCA) explained 83.7 % of the study variables. The first principal component, PC1, explained 69.2 %, and the second PC2 explained 14.5 % of the variation (Figure 2). All of the study variables were positively influenced by the ORF-strains inoculation. A close positive relationship existed between the nutrient uptake and the seedling biomass. Na showed stronger positive correlations with 2AP, proline, and DPPH (antioxidant activity) than with other nutrients. The proline level had the highest positive correlations with both antioxidant activity and 2AP level. The ORF strains had stronger positive relationships with growth parameters, chlorophyll content, proline level, antioxidant activity, and nutrient uptake in fresh leaves of KDML105 rice seedlings than the CRF strains. 258
259
260
261
262
263
264
265
266
267
268
269
270




Figure 2. Principal component analysis showing the relationship between growth parameters (shoot and root lengths and fresh and dry biomass), chlorophyll content, nutrient (N, P, K, Ca, Mg, and Na) uptake, antioxidant activity, proline accumulation, and 2AP level of KDML105 rice seedlings as affected by ST-PGPR inoculation.

3. Discussion

The premium aromatic rice variety Khao Dawk Mali 105 (KDML105) comprises about 50% of the rainfed paddy rice production in a huge area of Thung Kula Rong Hai (TKR), Northeastern Thailand. In addition, the KDML105 rice produced in the TKR region possesses a stronger aroma than rice cultivated in other areas of the country as well as in other countries [18], and thus it is traded as a premium quality rice with high price tag in both local and global markets. It is well known that the yield and aroma quality of KDML105 in the TKR region has been negatively affected by naturally high salinity and drought conditions [19]. The problem is exacerbated by the increase in drought as a result of global climate change. The use of salt-tolerant plant growth-promoting rhizobacteria (ST-PGPR) is a promising, sustainable, and cost-effective alternative to chemical management that can be used to mitigate these problems [14,15]. In the present study, the shoot and root biomass, as well as the chlorophyll content (SPAD unit), of the KDML105 rice seedlings were significantly enhanced by most of the selected ST-PGPR compared to their respective controls (Table 1,2). Several other studies have confirmed that seed priming and the inoculation of ST-PGPR improved rice seed germination, chlorophyll content, and photosynthetic capacity as well as rice growth and yield [11-12,20]. In the present study, it was interesting to note that the highest IAA-producing and the highest salt-tolerant ST-PGPR, *Sinomonas* sp. ORF15-23 also yielded the highest chlorophyll content and other rice growth parameters at all levels of salinity (Table 2). In contrast, the inoculation of non-IAA-producing strain CRF5-8 [17] as well as the lowest salt-tolerant strain, *Burkholderia* sp. CRF16-3 resulted in the lowest seedling growth parameters that were similar to the respective controls (Table 1). This phenomenon highlighted the importance of the IAA-production and salt-tolerance property of the PGPR in promoting the rice growth under salt stress. IAA has been demonstrated to increase root growth and surface area, leading to higher nutrient uptake and thereby improving plant growth as well as stress tolerance [21-23]. Previous studies showed that the growth-promoting effects on rice under salt stress are attributable to strain variability of ST-PGPR, which could enhance salt tolerance by altering root morphology, modifying root- to-shoot communication, increasing nutrients uptake, maintaining ion homeostasis, decreasing oxidative damage, and elevating

photosynthetic capacity [24-27]. Therefore, key specific microbial species, not the microbial richness or diversity, determined the efficiency of growth promotion by each individual ST-PGPR. In the present study, ST-PGPR inoculation not only promoted rice seedling growth but also improved shoot N, P, K, Ca, and Mg uptake of the seedlings, particularly inoculation with *Sinomonas* sp. ORF15-23, compared to the uninoculated seedlings (Table 3,4). It is possible that *Sinomonas* sp. ORF15-23 was more compatible with the rice KDML105 than other ST-PGPR. However, the exact mechanisms behind this observation remains to be answered. Genomic analysis of whole genome sequence and transcriptomics would be able to gain insights into growth promoting and salt tolerance mechanisms of *Sinomonas* sp. ORF15-23 as exemplified in recent publication [28-29]. In other words, the use of promising ST-PGPR effectively mitigated the deleterious effect of excessive salinity levels. It would be interesting to continue examining the effect of ST-PGPR inoculation on grain productivity and quality under practical field conditions in the future studies.

Apart from stimulating plant growth, IAA produced by ST-PGPR also performs a key role in ameliorating stress in plants. Phytohormone-producing bacteria increase plant tolerance to salinity stress, thereby promoting plant growth under excessive salinity [8, 30-32]. Auxin produced by *Bacillus amyloliquefaciens* RWL-1 has been reported to increase salinity stress tolerance in rice (*Oryza sativa* L.) [31]. Rangseekeaw *et al.*, [33,34] investigated three plant growth promoting abilities (IAA and siderophore production and phosphate solubilization). IAA production by actinobacteria *D. abyssi* MT1.1^T at 150 mM NaCl was three-fold decreased as compared to those production at 0 mM NaCl. Similarly, the reduction in IAA production by *D. profundus* MT2.2^T (decreased from 12.20 to 7.73 $\mu\text{g mL}^{-1}$) and *D. nishinomiyaensis* DSM20448^T (decreased from 16.64 to 9.39 $\mu\text{g mL}^{-1}$), was recorded at 150 mM NaCl. There is some evidence that IAA production is increased with increasing NaCl concentration. The results of our previous study indicated that the highest IAA-producing strain *Sinomonas* sp. ORF15-23 could grow best under salt stress [17]. *Sinomonas* sp. ORF15-23 also exhibited the highest ability in promoting rice seedling growth in the present study (Table 1). This result implied that besides having mechanisms for stress tolerance (e.g., IAA production, antioxidant activity, and potassium intake) [17]. ST-PGPR also transmitted some level of tolerance to the rice seedling under green houses. Salt stress causes osmotic stress in the early phases, leading to the accumulation of reactive oxygen species (ROS) that are harmful to plant cells. For example, hydrogen peroxide (H_2O_2), an important nonradical ROS was found to increase in tomato under 150 mM NaCl stress compared to non-inoculated tomato without salt stress [33,34]. Antioxidant activity plays a vital role in detoxifying ROS induced by salinity stress [35]. In the present study, the antioxidant activity (DPPH radical scavenging activity) in the leaves of the rice seedlings increased with an increasing salt concentration, and the activity was significantly enhanced by the inoculation with ST-PGPR strains (Table 2).

To maintain osmotic balance and optimum ROS concentration under stress conditions, plants synthesize antioxidants and osmoprotectants (osmolytes) such as proline [33-34, 36-37], an amino acid that is one of the most important osmolytes in response to salinity stress. In the present study, the proline content significantly increased with the inoculation of ST-PGPR strains, particularly *Sinomonas* sp. ORF15-23 that provided the maximum proline increase (163%) at 50 mM NaCl (Table 2). In addition, the PCA indicated a close relationship between DPPH radical scavenging activity and proline level (Figure 2). Proline accumulation in plants is a primary defense response to environmental stresses, including excessive salinity. The role of proline during stress generally includes osmotic adjustment, detoxification of ROS, and protection of membrane integrity as well as storage of organic carbon and nitrogen [38-39]. Under stressful conditions, it has been observed that proline also functions as a radical scavenger, thus performing a dual function as an osmolyte compound and an antioxidant [40]. Several studies have shown that proline effectively enhanced the salt tolerance and growth of various crops such as olives, tobacco, and rice seedlings [41-43]. The inoculation of bacterial isolate RWL-1 yielded greater synthesis of

various amino acids, including proline, under salinity stress [31]. Under salt-stress conditions, proline accumulation was observed in rice inoculated with ST-PGPR strain TY0307, resulting in enhanced salt tolerance, growth, and yield of rice [44]. Soil salinity induces adverse effects on seedling establishment and plant biomass accumulation [45-46]. Although rice possess inherent salt-tolerant strategies (4 dS m^{-1}), excessive soil salinity could damage seedling establishment and further inhibit growth of rice and soil microbes are of pivotal importance for plant growth, especially in adverse ecosystems such as those with saline soil conditions [47]. If the intensity increases in growing conditions will affect the amount of microorganisms will decrease and reduce activities that are beneficial to plants. Our results confirmed the increase of proline accumulation in rice when exposed to salinity stress and the enhancement of proline production by ST-PGPR inoculation that enhanced salt tolerance in the rice seedlings and thereby improved the growth of seedlings during salt stress (Table 2). In addition to function as an osmoprotectant and an antioxidant, proline has been recognized as the key precursor for the biosynthesis of 2AP, a major volatile compound of aromatic rice, including the KDML105 variety [48-49]. Several investigations have concluded that the 2AP content of KDML105 rice seedlings was markedly enhanced when exposed to salt stress, and this can be attributed to an increased accumulation of its precursor proline [48, 50-52]. Our findings agreed with these previous studies in that the 2AP content of all the treatments increased along with the proline content in KDML105 rice seedlings under salt stress, particularly between 0 and 50 mM NaCl (Table 2). The PCA indicated that the proline level (Figure 2) had the highest positive correlations with both 2AP and antioxidant activity [17,53]. In addition, the results of this study indicated that the 2AP level was significantly higher in the inoculated seedlings than in the uninoculated seedlings. It is interesting to note that the high 2AP-producing ST-PGPR strains *Sinomonas* sp. ORF15-23, *Enterobacter* sp. ORF10-12, and *Burkholderia* sp. CRF16-3 yielded the maximum 2AP content in the seedlings at 50, 100, and 150 mM NaCl, respectively (Table 6). Previous study has shown that the inoculation of high 2AP-producing rhizobacterial strains could increase the 2AP levels in the grains of the aromatic rice variety Basmati-370 [13].

In addition to osmoregulation and ROS scavenging (antioxidant activity), ion homeostasis (acid-base balance) is also considered an important defense mechanism of rice against salinity stress. The main toxic salt ions damaging to crop plants are Na^+ and Cl^- [54]. Under salt stress, extracellular Na^+ inhibits root K^+ uptake therefore a high K^+/Na^+ ratio is important for salt tolerance. In the present study, the ST-PGPR-inoculated seedlings had a higher K^+/Na^+ ratio than the uninoculated seedlings, and this may have led to the higher salt tolerance (Table 5). The inoculation of *Azospirillum* to salt-stressed maize restricted Na^+ uptake and enhanced the uptake of K^+ and Ca^{2+} in cv. 323, thus maintaining a high K^+/Na^+ ratio. The K^+/Na^+ ratio was significantly higher in salt tolerance maize cv. 324 than the salt-sensitive cv. 323 [55]. Under stressful conditions, IAA was shown to increase both proline and K contents and improve the nutritional, physiological, and metabolic activities of the plant [56]. Our observations are in accordance with this previous report in that the inoculation with the highest IAA-producing strain, *Sinomonas* sp. ORF15-23 resulted in the highest proline, Ca, and K uptake under salt stress (Table 2,3,4). The increase in proline, Ca, and K uptake might have led to improvements in the growth and salt tolerance of the rice seedlings. Therefore, K is one of the vital nutrients playing a critical role in plant stress. It was observed that high-affinity Na^+ uptake was found in K^+ -starved seedlings of several cereal crops, including rice. Furthermore, the Na^+ uptake was very rapid, and the K_m value was low under low K^+ and Ca^{2+} concentrations. However, high-affinity Na^+ uptake was sensitive to external K^+ [57-58]. These previous findings emphasize the importance of K in enhancing rice growth and salt tolerance under high salinity; thus, K should be available in sufficient quantity, particularly in the rhizosphere soil, throughout the growing season. One possible explanation could be that the exudation of specific compounds from ST-PGPR, and the growth promotion of roots both contributed

to the stimulation of microbial activity and modified the nutritional status in the rhizosphere under salt stress conditions [59-62]. The enhanced activities of IAA production, antioxidant activity, and potassium intake by ST-PGPR, could benefit the transformation of soil nutrients (such as K⁺ and Na⁺) and further promotes the overall availability of soil nutrients.

The results of this study revealed the promising benefits of the ST-PGPR strains for rice growth and aromatic quality (2AP) under both normal and saline conditions. The PCA indicated that the ST-PGPR rhizobacteria from organic rice farming practice (ORF strains) had stronger positive relationships with each of the study variables than the those from conventional rice farming practice (CRF strains) (Figure 2). Several studies have shown that plant adaptation to local/stress environments is driven by the co-adaptation of plants and rhizosphere microbes via a complex hormonal signaling pathway [63-64], and IAA appears to play a major role in microbe-plant interactions [65]. Exposure to excessive salinity was found to decrease maize and wheat root attachment by *Azospirillum brasilense* [66]. Similar finding was observed in this study as seen from a decrease in rhizobacterial count with an increasing salinity. However, the highest IAA-producing strain, *Sinomonas* sp. ORF15-23 maintained the highest count of 10⁸ CFU mL⁻¹ under all NaCl levels (Table 6). The high number of *Sinomonas* sp. ORF15-23 may be the reason for its ability in promoting the rice seedling growth and salt tolerance. Therefore, the use of ST-PGPR(s) could be an alternative option in alleviating salinity problems and enhancing rice yield and quality in KDML105 rice grown in the inland salt-affected areas such as Thung Kula Rong Hai (TKR). However, the use of the ST-PGPR inoculants in actual field conditions requires further investigation.

4. Materials and Methods

4.1. Rice Rhizobacterial Isolates

Nine KDML105 rice rhizobacterial strains that exhibited various degrees of tolerance to high salt concentrations (0 to 3% NaCl) were selected from our previous study [16], to evaluate their effects on KDML105 rice seedlings' growth and salt tolerance. All of the strains were able to produce IAA and promote the production of 2AP in KDML105 rice seedlings under salt stress. These selected strains were considered as salt-tolerant plant growth-promoting rhizobacteria (ST-PGPR). Five and four isolates were obtained from organic rice farming (ORF) and conventional rice farming (CRF), respectively [67]. *Micrococcus* sp. ORF15-19 and *Sinomonas* sp. ORF15-23 displayed the highest levels of salt tolerance, while *Burkholderia* sp. CRF16-3 displayed the lowest salt tolerance (Table 6).

Table 6. Effect of the inoculation of rhizobacterial isolates from organic and conventional farming practices on the IAA production, 2AP level of KDML105 rice seedlings and rhizobacterial count under different salt stress conditions.

Strain	Genus	IAA production (µg IAA mL ⁻¹)	2AP level of KDML105 rice seedlings (µg·kg ⁻¹)				Rhizobacterial population (CFU mL ⁻¹)			
			NaCl (% w/v)				NaCl (% w/v)			
			0	50	100	150	0	1	2	3
<i>Organic farming</i> ²										
ORF4-13	<i>Sinomonas</i> sp.	155.1	11.01	13.23	7.55	4.87	8.7×10 ⁸	2.3×10 ⁸	8.3×10 ⁷	6.7×10 ⁷
ORF10-12	<i>Enterobacter</i> sp.	47.7	14.31	18.7	12.87	7.14	2.2×10 ⁹	2.7×10 ⁸	1.0×10 ⁸	1.7×10 ⁷
ORF15-19	<i>Micrococcus</i> sp.	147.2	14.64	18.71	8.54	6.53	2.3×10 ⁹	1.1×10 ⁹	8.3×10 ⁸	1.5×10 ⁸
ORF15-20	<i>Micrococcus</i> sp.	127.8	15.39	18.24	6.62	5.88	7.2×10 ⁸	1.5×10 ⁸	1.3×10 ⁷	1.2×10 ⁷
ORF15-23	<i>Sinomonas</i> sp.	155.6	15.64	19.61	10.13	6.22	2.1×10 ⁹	1.3×10 ⁹	8.3×10 ⁸	2.1×10 ⁸
<i>Conventional farming</i> ²										
CRF5-8	unidentified	ND	12.44	13.64	8.21	4.65	1.2×10 ⁹	2.5×10 ⁸	6.2×10 ⁷	3.5×10 ⁷

CRF14-15	<i>Sinomonas</i> sp.	84.5	10.65	11.58	6.92	4.41	9.7×10 ⁸	3.3×10 ⁸	2.7×10 ⁸	6.7×10 ⁷
CRF16-3	<i>Burkholderia</i> sp.	7.3	14.06	17.43	10.12	9.43	3.8×10 ⁶	6.7×10 ⁵	3.3×10 ⁵	1.7×10 ⁵
CRF17-18	<i>Bacillus</i> sp.	55.1	11.03	12.01	6.43	5.75	1.1×10 ⁹	2.5×10 ⁸	7.8×10 ⁷	3.5×10 ⁷

¹ND = Not Detectable; ² Farming Practice.

Adapted from Chinachanta and Shutsrirung [7].

4.2. Effect of ST-PGPR inoculation on rice seedling growth under salt stress

The ability of nine selected ST-PGPR strains in enhancing KDML105 rice seedling growth under various NaCl concentrations was determined. The responses of the seedling to ST-PGPR inoculation were evaluated by analysis of the following growth parameters, chlorophyll content, nutrient concentration, antioxidant activity, and proline accumulation.

4.2.1. Preparation of ST-PGPR pellets and rice seedlings

This experiment was conducted using a completely randomized design (CRD) in a factorial scheme (10×4), with three replications, consisting of nine selected ST-PGPR strains plus one uninoculated control (ten treatments) and four NaCl concentrations (0, 50, 100, and 150 mM NaCl).

The nine selected ST-PGPR strains were grown in 25 mL nutrient broth (NB) for three days at 37 °C with shaking at 120 rpm. The ST-PGPR cells were collected by centrifugation at 10,000 rpm for 15 min to separate the culture broth from the pellet cells. The cell pellets were diluted with 100 mL sterile distilled water to obtain a cell concentration of 10⁶ colony-forming units (CFU) per mL (OD₆₀₀ ~ 0.2). This cell suspension was used as inoculum for seed biopriming and seedling inoculation. Sterile distilled water was used as the negative control (without ST-PGPR inoculation).

Rice (*Oryza sativa* L.) seeds variety KDML105 were used to evaluate the ability of selected ST-PGPR to promote growth and salt tolerance in rice. The seeds were surface sterilized in a mixture of 0.2% Tween 80 and 2% sodium hypochlorite for 3 min. The seeds were then washed three times with 70% ethanol, followed by rinsing five times with sterile water. The sterile seeds were soaked (seed biopriming) in the pellet suspension of each ST-PGPR strain according to the treatment and were then incubated in the dark at 25°C for 24 h [68]. The bioprimed seeds were then placed at an equal distance on sterile wet tissue paper in a Petri dish (20 seeds per plate) using sterile forceps (five replicates per treatment) and kept in a plant growth chamber under the dark at 25 °C. Four days after germination, 10 uniform seedlings from each treatment were selected and transplanted into a growth pouch containing Hoagland's nutrient solution (pH 7). The rice seedlings were initially irrigated with 1/4 strength Hoagland solution for five days, and the solution was replaced twice during this period. Then, the seedlings were irrigated with 1/2 strength Hoagland solution for two days. After that, the irrigation medium was changed to a full-strength Hoagland solution [69] with four salinity levels (0, 50, 100, and 150 mM NaCl). The average EC of an irrigation medium at each NaCl concentration was 2.06, 7.69, 13.78, and 19.51 dS m⁻¹, respectively. The full-strength solution were refreshed twice per week. The uninoculated (controls) and inoculated seedlings were grown in a climate-controlled room (12:12 light: dark photoperiod, 25±3°C, with a light level of approximately 5.8 klux).

The rice seedlings from each pouch were harvested at 30 days after transplanting, and then four replications of the seedlings were determined for growth parameters (shoot and root length; shoot and root dry weight). The leaves and root samples were dried to a constant weight at 65 °C for 48 hr. After that, the dry matter was weighed, and the dried samples were milled into powder, stored in plastic bags, and then kept in a desiccator for analysis of nutrient content. The remaining fresh seedlings (six replications) were used to determine antioxidant activity and proline accumulation in the leaves.

4.3. Chemical analysis

Leaf chlorophyll content was monitored at the third leaf stage after applying the salt stress to the seedlings (at 30 days) using a SPAD meter (SPAD-502, Minolta Camera Co., Ltd., Japan). The dried plant leaves were ground, homogenized, and used to determine the concentration of macronutrients. The total nitrogen (N) content (%) was determined by a modified Kjeldajl digestion (colorimetric) method [70]. The digestion was maintained at a boiling point of 350 °C. Ammonia was distilled from an alkaline medium and absorbed in an unstandardized boric acid solution and titrated with standard HCl solution. For the determination of total phosphorus (P), potassium (K), calcium (Ca), magnesium (Mg), and sodium (Na), the method described by Fageria [71] was applied. The total P concentration (%) in the samples was quantified spectrophotometrically using the vanado-molybdate phosphoric acid yellow colour method [72], with a UV-visible spectrophotometer (Shimadzu UV-VIS 1201, Shimadzu Co. Kyoto, Japan). The concentrations of K, Ca, Mg, and Na in the sample extracts were analyzed by an atomic absorption spectrophotometer (AAS) (Spectra AA240 FS, Varian, New Jersey, USA). Each sample was measured in triplicate. The nutrient uptake was calculated from the nutrient concentration and the dry matter of each sample using the following formula.

$$\text{Nutrient uptake (mg plant}^{-1}\text{)} = \frac{\text{Nutrient content (\%)} \times \text{Dry matter (mg plant}^{-1}\text{)}}{100}$$

Nutrient uptake = g plant⁻¹ (macronutrients) or mg plant⁻¹ (micronutrients)

Nutrient content (%) = Element concentration = in g kg⁻¹ (for macronutrients) or mg kg⁻¹ (for micronutrients) (1)

Dry matter = shoot dry weight = in g plant⁻¹ (for macronutrients) or mg kg⁻¹ (for micronutrients)

For the antioxidant activity analysis, the oven-dried leaf samples were defatted twice with hexane (1:20 w/v) for 30 min. The defatted rice leaf fraction was extracted twice with 99.9% methanol (1:20 w/v) in an electrical shaker overnight at room temperature and then filtered through Whatman No.1 filter paper. The extracts were evaporated to dryness at 50 °C by a vacuum rotary evaporator. The extract in the evaporator flask was eluted with methanol to a volume of 100 mL, then kept in a volumetric flask. The extracts were stored in the freezer at -18°C until use in further analysis. All analyses were performed within two weeks of extraction.

The free radical scavenging capacity was estimated following a previously reported procedure using 2,2'-diphenyl-1-picrylhydrazyl radical (DPPH) [73]. A synthetic antioxidant, BHT (99.0% purity, Rankem, India), was used as a reference. DPPH free radical-scavenging ability was calculated using the following formula:

$$\text{Scavenging ability (\%)} = \frac{[\text{Absorbance at 517 nm of the control} - \text{Absorbance at 517 nm of the sample}]/\text{Absorbance at 517 nm of the control} \times 100}{100} \quad (2)$$

Proline content was determined by standard method as described by [33]. Dried leaf powder of each sample (0.1 g) was used to extract the proline and the absorbance of the leaf extract was measured at 520 nm by a spectrophotometer (Shimadzu UV-VIS 1201, Shimadzu Co. Kyoto, Japan).and was recorded against pure toluene as a reference blank. The proline concentration was calculated from a standard curve prepared from pure proline (Sigma).

4.4. Statistical analysis

Two-way ANOVA together with LSD values at a 1% probability level [74] was used for analyzing collected data using Statistix 9 (Analytical Software, Inc., Tallahassee, FL, USA). Principal component analysis (PCA) is a statistical technique that allow easier analysis of a large dataset with visual by reducing the complexity and noise of the data, and highlight the most important features and relationships between observed parameters. In this study, the relationships between growth parameters (shoot and root length and fresh and dry biomass), chlorophyll content, nutrient uptake (N, P, K, Ca, Mg, and Na), antioxidant activity, proline, and 2AP level accumulation of KDML105 rice seedlings as affected by ST-PGPR inoculation were evaluated using PCA. The measured parameters were introduced as variables in the PCA using R 1.2.1335 [75].

5. Conclusions

The present investigation revealed that the inoculation of most of the tested ST-PGPR strains, particularly *Sinomonas* sp. ORF15-23, significantly reduced the extent of growth suppression due to excessive salinity, leading to incremental increases in rice seedling growth and salt tolerance. In addition, the 2AP (a key volatile aroma compound) level in the rice seedlings was markedly enhanced by ST-PGPR inoculation, and this may have led to high 2AP levels in the rice grains. **These findings suggest that *Sinomonas* sp. ORF15-23 can be used to enhance KDML105 rice seedlings growth and improve soil nutrient uptake in saline soil. This information provides a basis background for development of a microbial technology to aid in restoration of saline-degraded areas.** Nevertheless, further investigations under field conditions are needed for the development of the promising ST-PGPR strain(s) as a bio-inoculant for rice production in salinity affected area such as **the effect of ST-PGPR inoculation on grain quality and yield in future studies.**

6. Patents

Author Contributions: K.C. carried out the study, methodology, discussed the results, and wrote the manuscript; writing—original draft preparation, K.C. and A.S.; conceptualization and discussion, K.C., A.S., C.S., D.T.L., L.H., D.L. and C.P.; supervision and writing—review and editing, W.P., D.T.L., D.L. and C.P.; funding acquisition, K.C. W.P., D.T.L., and C.P. -a. All authors have read and agreed to the published version of the manuscript.

Funding: This study was funded by Junior Research Fellowship Program 2022 from France Embassy in Thailand and Agropolis Fondation, Montpellier, France to K.C. W.P. and K.C. acknowledge the partial support from Chiang Mai University through the Research Center of Microbial Diversity and Sustainable Utilization. C.P. and K.C. acknowledge the partial support from Chiang Mai University.

Data Availability Statement: Data sharing does not apply to this article as no datasets were generated or analyzed during the current study.

Acknowledgments: The authors are indebted to the farmers of Community Enterprise of Thamo Organic Agricultural Group, Surin Province, and Na So Rice Farmers Group (Nature Conservation Club), Kut Chum, Yasothon Province, for their help during field sampling, as well as the Alliance of Biodiversity International and Center International of Tropical Agriculture (CIAT), Asia Hub, Common Microbial Biotechnology Platform (CMBP), Hanoi, Vietnam, and the Institute for Plant Sciences of Montpellier (IPSiM), Univ Montpellier, CNRS, INRAE, Institut Agro, Montpellier, France. W.P. and K.C.-a. acknowledges the partial support from Chiang Mai University through the Research Center of Microbial Diversity and Sustainable Utilization. D.T.L. and K.C.-a. acknowledges the partial support from Junior Research Fellowship Program from France Embassy in Thailand and Agropolis Fondation, Montpellier, France. The authors would like to thank Dr. Dale Taneyhill for English editing throughout the manuscript. This paper was partially funded by Chiang Mai University, Thailand.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Arunin, S.; Pongwichian, P. Salt-affected soils and management in Thailand. *Bull. Soc. Sea Water Sci. Jpn.* **2015**, *69*, 319–325, doi: 10.11457/swsj.69.319. 588
2. Chen, S.; Gollop, N.; Heuer, H. Proteomic analysis of salt-stressed tomato (*Solanum lycopersicum*) seedlings: effect of genotype and exogenous application of glycinebetaine. *J. Exp. Bot.* **2009**, *60*, 2005–2019, doi: 10.1093/jxb/erp075. 589
3. Suzuki, N.; Bassil, E.; Hamilton, J.S.; Inupakutika, M.A.; Zandalinas, S.I.; Tripathy, D.; Luo, Y.; Dion, E.; Fukui, G.; Kumazaki, A.; Nakano, R.; Rivero, R.M.; Verbeck, G.F.; Azad, R.K.; Blumwald, E.; Mittler, R. ABA is required for plant acclimation to a combination of salt and heat stress. *PLoS One* **2016**, *11*(1), e0147625, doi: 10.1371/journal.pone.0147625. 590
4. Grattan, S.R., L. Zeng, M.C. Shannon and Roberts, S.R. Rice is more sensitive to salinity than previously thought. *Calif. Agric.* **2002**, *56* (6), 189–195. 591
5. Rad, H.E.; Aref F.; Rezaei, M. Evaluation of salinity stress affects rice growth and yield components in northern Iran. *Am. J. Sci. Res.* **2012**, *54*, 40–51. 592
6. Egamberdieva, D.; Wirth, S.; Bellingrath-Kimura, S.D.; Mishra, J.; Arora, N.K. Salt-tolerant plant growth promoting rhizobacteria for enhancing crop productivity of saline soils. *Front. Microbiol.* **2019**, *10*: 2791, doi: 10.3389/fmicb.2019.02791. 593
7. Chinachanta, K.; Shutsrirung, A. Screening for P- and K- Solubilizing, and Siderophore Producing Capacity of Rhizobacteria from Khao Dawk Mali 105 Aromatic Rice. *IOP Conf. Ser. Earth and Environ. Sci.* **2021**, *858*(1), 012004, doi: 10.1088/1755-1315/858/1/012004. 594
8. Rajput, L.U.B.N.A.; Imran, A.; Mubeen, F.; Hafeez, F.Y. Salt-tolerant PGPR strain *Planococcus rifietoensis* promotes the growth and yield of wheat (*Triticum aestivum* L.) cultivated in saline soil. *Pak. J. of Bot.* **2013**, *45*, 1955–1962. 595
9. Sarkar, A.; Ghosh, P.K.; Pramanik, K.; Mitra, S.; Soren, T.; Pandey, S.; Mondale, M.H.; KantiMaiti, T. A halotolerant *Enterobacter* sp. displaying ACC deaminase activity promotes rice seedling growth under salt stress. *Res. Microbiol.* **2018**, *169*(1), 20–32, doi: 10.1016/j.resmic.2017.08.005. 596
10. El-Esawi, M.A.; Alaraidh, I.A.; Alsahli, A.A.; Alamri, S.A.; Ali, H.M.; Alayafi, A.A. *Bacillus firmus* (SW5) augments salt tolerance in soybean (*Glycine max* L.) by modulating root system architecture, antioxidant defense systems and stress-responsive genes expression. *Plant Physiol. Biochem.* **2018**, *132*, 375–384, doi: 10.1016/j.plaphy.2018.09.026. 597
11. Zhang, S.; Fan, C.; Wang, Y.; Xia, Y.; Xiao, W.; Cui, X. Salt-tolerant and plant-growth-promoting bacteria isolated from high-yield paddy soil. *Can. J. Microbiol.* **2018**, *64*, 968–978, doi: 10.1139/cjm-2017-0571. 598
12. Sharma, A.; Shankhdhar, D.; Sharma, A.; Shankhdhar, S.C. Growth promotion of the rice genotypes by pgprs isolated from rice rhizosphere. *J. Soil Sci. Plant Nutr.* **2014**, *14* (2), 505–517, doi.org/10.4067/S0718-95162014005000040. 599
13. Deshmukh, Y.; Khare, P.; Patra, D. Rhizobacteria elevate principal basmati aroma compound accumulation in rice variety. *Rhizosphere* **2016**, *1*, 53–57, doi: 10.1016/j.rhisph.2016.07.001. 600
14. Jha, B.; Gontia I.; Hartmann, A.; The roots of the halophyte *Salicornia brachiata* are a source of new halotolerant diazotrophic bacteria with plant growth-promoting potential. *Plant Soil* **2012**, *356*, 265–277, doi: 10.1007/s11104-011-0877-9. 601
15. Viscardi, S.; Ventrino, V.; Duran, P.; Maggio, A.; De Pascale, S.; Mora, M.L.; Pepe, O. Assessment of plant growth promoting activities and abiotic stress tolerance of *Azotobacter chroococcum* strains for a potential use in sustainable agriculture. *J. Soil Sci. Plant Nutr.* **2016**, *16*, 848–863, doi: 10.4067/S0718-95162016005000060. 602
16. Chinachanta, K.; Shutsrirung, A.; Herrmann, L.; Lesueur, D.; Pathom-aree, W. Enhancement of the aroma compound 2-acetyl-1-pyrroline in thai jasmine rice (*Oryza sativa*) by rhizobacteria under salt stress. *Biology*, **2021**, *10*, 1065, doi: 10.3390/biology10101065. 603
17. Chinachanta, K.; Shutsrirung, A.; Herrmann L.; Lesueur, D. Isolation and characterization of KDML105 aromatic rice rhizobacteria producing indole-3-acetic acid: impact of organic and conventional paddy rice practices. *Lett. Appl. Microbiol.* **2021**, doi: 10.1111/lam.13602. 604
18. Yoshihashi, T.; Nguyen, T.T.H.; Kabaki, N. Area dependency of 2-acetyl-1-pyrroline content in an aromatic rice variety, Khao Dawk Mali 105. *Jpn. Agric. Res. Q.* **2004**, *38*(2), 105–109, doi: 10.6090/jarq.38.105. 605
19. Gupta, A.; Singh, A.N.; Tiwari, R.K.; Sahu, P.K.; Yadav, J.; Srivastava, A.K.; Kumar, S. Salinity Alleviation and reduction in oxidative stress by *Endophytic* and *Rhizospheric Microbes* in two rice cultivars. *Plants* **2023**, *12*, 976, https://doi.org/10.3390/plants12050976. 606
20. Kim, S.Y.; Lim, J.H.; Park, M.R.; Kim, Y.J.; Park, T.I.; Se, Y.W.; Choi, K.G.; Yun S.J. Enhanced antioxidant enzymes are associated with reduced hydrogen peroxide in barley roots under saline stress. *J. Biochem. Mol. Biol.* **2005**, *38*(2), 218–24, doi: 10.5483/bmb-rep.2005.38.2.218. 607
21. Diby, P.; Sarma, Y.R.; Srinivasan, V.; Anandaraj, M. *Pseudomonas fluorescens* mediated vigour in black pepper (*Piper nigrum* L.) under greenhouse cultivation. *Ann Microbiol.* **2005**, *55*(3), 171–74, doi: 10.1016/j.pmpp.2022.101836. 608
22. Paul, D.; Sarma, Y.R. Plant growth promoting rhizobacteria (PGPR) mediated root proliferation in black pepper (*Piper nigrum* L.) as evidenced through GS root software. *Arch. Phytopathol. Plant Prot.* **2006**, *39*(4), 311–314, doi: 10.1007/s13593-014-0233-6. 609
23. Paul, D.; Lade, H. Plant-growth-promoting rhizobacteria to improve crop growth in saline soils: a review. *Agron. Sustain. Dev.* **2014**, *34*(4), 737–752, doi: 10.1007/s13593-014-0233-6. 610
24. Chandrasekaran, M.; Boughattas, S.; Hu, S.; Oh, S.-H.; Sa, T. A meta-analysis of arbuscular mycorrhizal effects on plants grown under salt stress. *Mycorrhiza* **2014**, *24*, 611–625. 611
25. Pan, J.; Peng, F.; Xue, X.; You, Q.; Zhang, W.; Wang, T.; Huang, C. The growth promotion of two salt-tolerant plant groups with PGPR inoculation: A meta-analysis. *Sustainability* **2019**, *11*, 378, doi:10.3390/su11020378. 612

26. Egamberdieva, D.; Wirth, S.; Jabborova, D.; Räisänen, L.A.; Liao, H. Coordination between *Bradyrhizobium* and *Pseudomonas* alleviates salt stress in soybean through altering root system architecture. *J. Plant Interact.* **2017**, *12*, 100–107. 648
649
27. Ghazanfar, B.; Cheng, Z.; Cuinan, W.U.; Liu, H.; Hezi, L.I.; Rehman, R.N.U.; Ahmad, I.; Khan, A.R. *Glomus etunicatum* root inoculation and foliar application of acetyl salicylic acid induced NaCl tolerance by regulation of NaCl & LeNHXL gene expression and improved photosynthetic performance in tomato seedlings. *Pak. J. Bot.* **2016**, *48*, 1209–12117. 650
651
652
28. Niu, S.; Gao, Y.; Zi, H.; Liu, Y.; Liu, X.; Xiong, X.; Yao, Q.; Qin, Z.; Chen, N.; Guo, L.; Yang, Y.; Qin, P.; Lin, J.; Zhu, Y. The osmolyte-producing endophyte *Streptomyces albidoflavus* OsILf-2 induces drought and salt tolerance in rice via a multi-level mechanism. *The Crop Journal* **2022**, *10*, 375–386, doi:10.1016/j.cj.2021.06.00834. 653
654
655
29. Kruasuwan, W.; Lohmaneeratana, K.; Munnoch, J.T.; Vongsangnak, W.; Jantrasuriyarat, C.; Hoskisson, P.A.; Thamchaipenet, A. Transcriptome landscapes of salt-susceptible rice cultivar IR29 associated with a plant growth promoting endophytic *Streptomyces*. *Rice* **2023**, *16*, (6), doi:10.1186/s12284-023-00622-7 656
657
658
30. Egamberdieva, D. Alleviation of salt stress by plant growth regulators and IAA producing bacteria in wheat. *Acta Physiol Plant.* **2009**, *31*, 861–864, doi: 10.1007/s11738-009-0297-0. 659
660
31. Shahzad, R.; Khan, A.L.; Bilal, S.; Waqas, M.; Kang S-M.; Lee, I-J. Inoculation of abscisic acid-producing endophytic bacteria enhances salinity stress tolerance in *Oryza sativa*. *Environ. Exp. Bot.* **2017**, *136*, 68–77, doi:10.1016/j.envexpbot.2017.01.010. 661
662
32. Kang, S.; Shahzad, R.; Bilal, S.; Khan, A.L.; Park, Y.G.; Lee, K.E.; Asaf, S.; Khan, M.A.; Lee, I-J. Indole-3-acetic-acid and ACC deaminase producing *Leclercia adecarboxylata* MO1 improves *Solanum lycopersicum* L. growth and salinity stress tolerance by endogenous secondary metabolites regulation. *BMC Microbiol.* **2019**, *19*(80), doi: 10.1186/s12866-019-1450-6. 663
664
665
33. Rangseekaew, P.; Barros-Rodriguez, A.; Pathom-aree, W.; Manzanera, M. Deep-sea actinobacteria mitigate salinity stress in tomato seedlings and their biosafety testing. *Plants* **2021**, *10*(8): 1687. doi: 10.3390/plants10081687. 666
667
34. Rangseekaew, P.; Barros-Rodriguez, A.; Pathom-aree, W.; Manzanera, M. Plant beneficial deep-sea actinobacterium, *Derma-coccus abyssi* MT1.1^T promote growth of tomato (*Solanum lycopersicum*) under salinity stress. *Biology* **2022**, *11*(2): 191. doi: 10.3390/biology11020191. 668
669
670
35. Gupta, B.; Huang, B. Mechanism of Salinity tolerance in plants: physiological, biochemical, and molecular characterization. *Int. J. Genom.* **2014**, Article ID 701596, doi: 10.1155/2014/701596. 671
672
36. El-Rahman, S.S.A.; Mazen, M.M.; Mohamed, H.I.; Mahmoud, N.M. Induction of defence related enzymes and phenolic compounds in lupin (*Lupinus albus* L.) and their effects on host resistance against Fusarium wilt. *Eur. J. Plant Pathol.* **2012**, *134*, 105–116, doi: 10.1007/s10658-012-0028-z. 673
674
675
37. Singh, R.; Singh, S.; Parihar, P.; Mishra, R.K.; Tripathi, D.K.; Singh, V.P.; Chauhan D.K.; Prasad, S.M. Reactive oxygen species (ROS): Beneficial companions of plants developmental processes. *Front. Plant Sci.* **2016**, *7*, 1299, doi: 10.3389/fpls.2016.01299. 676
677
38. Peng, Y.L.; Gao, Z.W.; Gao, Y.; Liu, G.F.; Sheng, L.X.; Wang, D.L. Eco-physiological characteristics of alfalfa seedlings in response to various mixed salt-alkaline stresses. *J. Integr. Plant Biol.* **2008**, *50*(1), 29–39, doi: 10.1111/j.1744-7909.2007.00607.x. 678
679
39. Heuer, B. Role of Proline in Plant Response to Drought and Salinity **2010**; pp. 213–238, doi: 10.1201/b10329-12. 680
40. Matysik, J.; Alia, A.; Bhalu, B.; Mohanty, P. Molecular mechanisms of quenching of reactive oxygen species by proline under stress in plants. *Curr. Sci.* **2002**, *82*(5), 525–532. 681
682
41. Hoque, M.A.; Banu, M.N.A.; Nakamura, Y.; Shimoishi, Y.; Murata, Y. Proline and glycinebetaine enhance antioxidant defense and methylglyoxal detoxification systems and reduce NaCl-induced damage in cultured tobacco cells. *J. Plant Physiol.* **2008**, *165*(8), 813–824, doi: 10.1016/j.jplph.2007.07.013. 683
684
685
42. Ben Ahmed, C.; Ben Rouina, B.; Sensoy, S.; Boukhriss, M.; Ben Abdullah, F. Exogenous proline effects on photosynthetic performance and antioxidant defense system of young olive tree. *J. Agric Food Chem.* **2010**, *58*(7), 4216–4222, doi:10.1021/jf9041479. 686
687
43. Deivanai, S.; Xavier, R.; Vinod, V.; Timalata, K.; Lim, O.F. Role of exogenous proline in ameliorating salt stress at early stage in two rice cultivars. *J. Stress Physiol. Biochem.* **2011**, *7*, 157–174. 688
689
44. Zhang, S.; Fan, C.; Wang, Y.; Xia, Y.; Xiao, W.; Cui, X. Salt-tolerant and plant- growth-promoting bacteria isolated from high-yield paddy soil. *Can. J. Microbiol.* **2018**, *64*, 968–978, doi: 10.1139/cjm-2017-0571. 690
691
45. Munns, R.; Tester, M. Mechanisms of salinity tolerance. *Annu. Rev. Plant Biol.* **2008**, *59*, 651–681. 692
46. Munns, R. Comparative physiology of salt and water stress. *Plant Cell Environ.* **2002**, *25*, 239–250. 693
47. Heijden, M.G.A.V.D.; Bardgett, R.D.; Straalen, N.M.V. The unseen majority: Soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecol. Lett.* **2008**, *11*, 296–310. 694
695
48. Yoshihashi, T.; Huong, N.T.T.; Inatomi, H. Precursors of 2-acetyl-1 pyrroline, a potent flavor compound of an aromatic rice variety. *J. Agric. Food Chem.* **2002**, *50*(7), 2001–4, doi: 10.1021/jf011268s. 696
697
49. Luo, H.; Zhang, T.; Zheng, A.; He, L.; Lai, R.; Liu, J.; Xing, P.; Tang, X. Exogenous proline induces regulation in 2-acetyl-1-pyrroline (2-AP) biosynthesis and quality characters in fragrant rice (*Oryza sativa* L.). *Sci. Rep.* **2020**, *2020*(10), 13971, doi: 10.1038/s41598-020-70984-1. 698
699
700
50. Sansenya, S.; Hua, Y.; Chumanee S.; Sricheewin, C. The combination effect of gamma irradiation and salt concentration on 2-acetyl-1-pyrroline content, proline content and growth of Thai fragrant rice (KDML 105). *Orient. J. Chem.* **2019**, *35*(3), 938–946, doi: 10.13005/ojc/350304. 701
702
703
51. Poonlaphdecha, J.; Maraval, I.; Roques, S.; Audebert, A.; Boulanger, R.; Bry, X.; Gunata, Z. Effect of timing and duration of salt treatment during growth of a fragrant rice variety on yield and 2-acetyl-1-pyrroline, proline, and GABA levels. *J. Agric Food Chem.* **2012**, *60*, 3824–3830, doi: 10.1021/jf205130y. 704
705
706

52. Kongpun, A.; Jaisiri, P.; Rerkasem, B.; Prom-u-thai, C. Impact of soil salinity on grain yield and aromatic compound in Thai Hom Mali rice cv. Khao Dawk Mali 105. *Agr. Nat. Resour.* **2020**, *54*, 74–78, doi: 10.34044/j.anres.2020.54.1.10. 707
708
53. Kongpun, A.; Pusadee, T.; Jaksomsak, P.; Chinachanta, K.; Tuiwong, P.; Chan-in, P.; Konsaeng, S.; Pathom-aree, W.; Utasee, S.; Wangkaew, B.; Prom-u-thai, C. Abiotic and Biotic Factors Controlling Aromatic Flavor Along the Value Chain of Fragrant Rice: A Review. *Rice Sci.* **2023**, *In press*. 709
710
711
54. Flowers, T.J. Improving crop salt tolerance. *J. Exp. Bot.* **2004**, *55*, 307–319, doi: 10.1093/jxb/erh003. 712
55. Hamdia, M.B.E.; Shaddad, M.A.K.; Doaa, M.M. Mechanisms of salt tolerance and interactive effects of *Azospirillum brasilense* inoculation on maize cultivars grown under salt stress conditions. *Plant Growth Regul.* **2004**, *44*(2), 165–74, doi: 10.1023/B:GROW.0000049414.03099.9b. 713
714
715
56. Armada, E.; Roldán, A.; Azcon, R. Differential activity of autochthonous bacteria in controlling drought stress in native *Lavandula* and *Salvia* plants species under drought conditions in natural arid soil. *Microb Ecol* **2014**, *67*, 410–420, doi: 10.1007/s00248-013-0326-9. 716
717
718
57. Garcíadeblás, B.; Senn, M.E.; Bañuelos M.A.; Rodríguez-Navarro, A. Sodium transport and HKT transporters: The rice model. *The Plant J.* **2003**, *34*, 788–801, doi: 10.1046/j.1365-313x.2003.01764.x. 719
720
58. Gao, J.P.; Chao, D.Y.; Lin, H.X.; Understanding abiotic stress tolerance mechanisms: Recent studies on stress response in rice. *J. Integr. Plant Biol.* **2007**, *49*(6), 742–750, doi: 10.1111/j.1744-7909.2007.00495.x. 721
722
59. Folli-Pereira, M.D.S.; Meira-Haddad, L.S.A.; Kasuya, M.C.M. Plant-microorganism interactions: Effects on the tolerance of plants to biotic and abiotic stresses. In *Crop improvement: New Approaches and Modern Techniques*. Springer: Berlin, Germany **2013**, pp. 209–239. 723
724
725
60. Vogelsang, K.M.; Reynolds, H.L.; Bever, J.D. Mycorrhizal fungal identity and richness determine the diversity and productivity of a tallgrass prairie system. *New Phytol.* **2006**, *172*, 554–562. 726
727
61. Dodd, I.C.; Perez-Alfocea, F. Microbial amelioration of crop salinity stress. *J. Exp. Bot.* **2012**, *63*, 3415–3428. 728
62. Ash, C. Fungi help trees hunt for food. *Science* **2016**, *353*, 661–661. 729
63. Pieterse, C.M.; Van der Does, D.; Zamioudis, C.; Leon-Reyes, A.; Van Wees, S.C. Hormonal modulation of plant immunity. *Annu. Rev. Cell Dev. Biol.* **2012**, *28*, 489–521, doi: 10.1146/annurev-cellbio-092910-154055. 730
731
64. Qu, Q.; Zhang, Z.; Peijnenburg, W.; Liu, W.; Lu, T.; Hu, B.; Chen, J.; Chen, J.; Lin Z.; Qian, H. Rhizosphere microbiome assembly and its impact on plant growth. *J. Agric. Food Chem.* **2020**, *68*(18), 5024–5038, doi: 10.1021/acs.jafc.0c00073. 732
733
65. Spaepen, S.; Vanderleyden, J. Auxin and plant-microbe interactions. *Cold Spring Harb. Perspect. Biol.* **2011**, *3*(4), a001438, doi: 10.1101/cshperspect. a001438. 734
735
66. Jofre, E.; Fischer, S.; Rivarola, V.; Balegno, H.; Mori, G. Saline stress affects the attachment of *Azospirillum brasilense* Cd to maize and wheat roots. *Can. J. Microbiol.* **1998**, *44*(5), 416–422, doi: 10.1139/w98-024. 736
737
67. Chinachanta, K.; Herrmann, L.; Lesueur, D.; Jongkaewwattana, S.; Santasup, C.; Shutsrirung, A. Influences of farming practices on soil properties and the 2-acetyl-1-pyrroline content of Khao Dawk Mali 105 rice grains. *Appl. Environ. Soil Sci.* **2020**, *2020*, 8818922, doi: 10.1155/2020/8818922. 738
739
740
68. Iwai, R.; Uchida, S.; Yamaguchi, S.; Sonoda, F.; Tsunoda, K.; Nagata, H.; Nagata, D.; Koga, A.; Goto, M.; Maki, T.-a.; et al. Effects of Seed Bio-Priming by Purple Non-Sulfur Bacteria (PNSB) on the Root Development of Rice. *Microorganisms* **2022**, *10*, 2197. <https://doi.org/10.3390/microorganisms10112197>. 741
742
743
69. Epstein, E. *Mineral Nutrition of Plants: Principles and Perspectives*. Wiley, New York, 1972. 744
70. Cataldo, D.A.; Schrader, L.E.; Youngs, V.L. Analysis by digestion and colorimetric assay of total nitrogen in plant tissues high in nitrate. *Crop Sci.* **1974**, *14*, 854–856, doi: 10.2135/cropsci1974.0011183X001400060024x. 745
746
71. Fageria, N.K. Critical P, K, Ca, and Mg contents in the tops of rice and peanut plants. *Plant and Soil* **1976**, *45*, (2), 421–431. 747
72. Jackson ML. *Soil chemical analysis: advanced course*. 2nd ed. Madison: Parallel Press, University of Wisconsin-Madison Libraries; 2005. 748
749
73. Brand-Williams, W.; Cuvelier, M.E.; Berset, C. Use of a free radical method to evaluate antioxidant activity. *LWT-Food Sci Technol* **1995**, *28*, 25–30, doi: 10.1016/S0023-6438(95)80008-5. 750
751
74. Steel, R.G.D.; Torrie, J.H.; Dickey, D.A. *Principles and Procedures of Statistics; Abiometrical Approach*, 3rd ed.; McGraw-Hill, Boston. 1997. 752
753
75. R Development Core Team, R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria, 2011. 754
755
756