



Combined use of Sentinel-2 images and Sentinel-1-derived moisture maps for soil organic carbon content mapping in croplands, South-western France

Diego Urbina Salazar, Emmanuelle Vaudour, Nicolas Baghdadi, Eric Ceschia, Dominique Arrouays

► To cite this version:

Diego Urbina Salazar, Emmanuelle Vaudour, Nicolas Baghdadi, Eric Ceschia, Dominique Arrouays. Combined use of Sentinel-2 images and Sentinel-1-derived moisture maps for soil organic carbon content mapping in croplands, South-western France. EGU 2021, Apr 2021, Vienna (AUSTRIA), Austria. , 2021, 10.5194/egusphere-egu21-8836 . hal-04220921

HAL Id: hal-04220921

<https://hal.inrae.fr/hal-04220921>

Submitted on 28 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

COMBINED USE OF SENTINEL-2 IMAGES AND SENTINEL-1/2 DERIVED MOISTURE MAPS FOR SOIL ORGANIC CARBON CONTENT MAPPING IN CROPLANDS, SOUTH-WESTERN FRANCE


Study area / objective

Material & methods


Results & discussion

S2 prediction performance using surface soil moisture (SM) from (SMP) as a covariate (in bold characters, models using SM as a covariate)

General flowchart



Soil moisture products (SMP)



Imaging Date S2	SMP	NS	MD	R ² _{cv}	RMSE _{cv} (g kg ⁻¹)	RPD _{cv}	NC	Soil Moisture (mv Vol%)			
								Min	Me	$\bar{x}$	Max
6 April 2017	7 April 2017	59	No	0.08	1.69	1.05	1	6.8	17.8	17.38	24.6
	19 May 2017	50	Yes	0.3	1.48	1.2	3	9	20.6	20.4	28
16 May 2017	25 May 2017	58	No	0.47	2.06	1.39	4				
	26 May 2017	58	Yes	0.5	2.01	1.43	3				
22 March 2018	21 March 2018	55	No	0.46	1.33	1.38	4				
	21 April 2018	47	Yes	0.6	1.13	1.57	5	5.4	13.2	13.02	18.8
11 May 2018	20 May 2018	73	No	0.1	2.42	1.07	2	22	26.4	26.24	31.4
	21 May 2018	73	Yes	0.1	2.42	1.06	2				
	21 May 2018	73	No	0.09	3.58	1.06	2				
	21 May 2018	73	Yes	0.03	3.68	1.03	2	11.2	19.8	19.88	26.4
	21 May 2018	73	No	0.44	3.06	1.35	3				
	21 May 2018	73	Yes	0.48	2.95	1.4	3	21.4	28.4	28.21	33.4
	21 May 2018	73	No	0.48	2.91	1.4	7				
	21 May 2018	73	Yes	0.47	2.91	1.38	8	20.8	22.6	23.66	29

Study Aim: Predict topsoil organic carbon (SOC) content over croplands, addressing : (i) the influence of the sentinel image date , and (ii) the use of soil moisture products derived from the S1 & S2 satellites in the spectral models.

Sentinel-2 (S2A) images and soil moisture products were acquired from the Theia platform (<https://www.theia-land.fr/>) and Sentinel-1 images from the European Space Agency's Copernicus platform.

Bare soil reflectance spectra were extracted then centered and PLSR with leave-one-out cross-validation was used on the bands to construct prediction models for SOC content.

*Moisture data in drier periods improved performance
*S1 data in the models did not improve performance
*Higher residues were associated with loamy textures in 2018

Relationship between texture classes and SOC prediction residuals



S2 prediction performance using raw S1

Imaging Date S1	NS	P	R ² _{cv}	RMSE _{cv} (g kg ⁻¹)	RPD _{cv}	NC
7 April 2017	59	VH	0.13	1.64	1.08	3
		VV	0.09	1.68	1.06	1
19 May 2017	50	VH	0.6	2.08	1.38	5
		VV	0.44	2.12	1.35	5
25 May 2017	58	VH	0.45	1.31	1.36	5
		VV	0.45	1.31	1.36	5
21 March 2018	55	VH	0.09	2.44	1.05	2
		VV	0.3	2.08	1.24	2
19 April 2018	47	VH	0.05	3.64	1.04	2
		VV	0.08	3.58	1.06	2
13 May 2018	58	VH	0.46	3.01	1.37	2
		VV	0.46	2.9	1.37	2
20 May 2018	73	VH	0.46	2.97	1.37	9
		VV	0.45	3	1.36	9

Conclusion

*The date of acquisition of the S2 images and other disturbing factors affect SOC prediction
*Perspectives to study the effect of Soil moisture (SM) on more dates and under different conditions (Possibility of determining optimal thresholds above which the SM can be included in prediction models).