



**HAL**  
open science

## **NIVA UC1b Agro-environmental indicators**

Eric Ceschia, Clélia Sirami, Christian Bockstaller, David Sheeren, Ludovic Arnaud, Emmanuel de Laroche, Arthur Favreau, Guillaume Marchand, Dominique Laurent

► **To cite this version:**

Eric Ceschia, Clélia Sirami, Christian Bockstaller, David Sheeren, Ludovic Arnaud, et al.. NIVA UC1b Agro-environmental indicators. Stakeholder Forum, European Environmental Bureau, Dec 2020, online (Brussels), Belgium. hal-04221939

**HAL Id: hal-04221939**

**<https://hal.inrae.fr/hal-04221939v1>**

Submitted on 28 Sep 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# UC1b Agro-environmental indicators

Stackholder Forum, 10/12/2020

**INRAE** : Christian Bockstaller, Eric Ceschia, David Sheeren, Clélia Sirami,

**IGN** : Arthur Favreau, Dominique Laurent, Guillaume Marchand,

**ASP** : Ludovic Arnaud, Emmanuel De Laroche



This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No. 842009

# 1. NIVA's objectives

## **NIVA objectives**

1. Harnessing innovations to simplify the governance;
2. Reducing socio-economic and administrative burden to farmers;
3. Reducing the gap between IACS data use and potential broader uses

**=> including high ambitions to improve environmental conditions and mitigate climate change**

# 1. UC1b OBJECTIVES

## **UC1b fundamental objectives**

- Agricultural practices have a strong impact on environment
- This impact should be measured to orient farming practices (farmers, advisers, providers, market) and to support environmental policies. (Decision making -> implementation -> monitoring)

**⇒ Need for agro-environmental indicators that could be available to farmers, agricultural advisers, policy makers, ONG...**

**⇒ Develop indicators, produce them on a large scale, test them with stakeholders in France (APCA, French Biodiversity Agency) and other EU countries (DK, NL, SP so far)**

# List of Agro-Environmental Indicators

➤ We proposed 11 A.E. indicators addressing 3 CAP objectives and 5 categories of environmental issues related to:

- **Climate mitigation**: C budget, reduction of N fertilisers
- **Water quality**: nitrate leaching, pesticides, herbicides, fungicides
- **Biodiversity**: biodiv. conservation, biological control, pollination
- **Soils**: quality, erosion
- **Landscapes**: aesthetic value



➤ For each type of indicator we propose between 1 and 3 methods of calculations: TIERS 1 to 3. All of them are evidence-based, published and several were adapted from the H2020 DiverImpacts project (scoring systems from 0 to 1),

➤ Three were considered as a priority by the DG Agri, DG Climate and the Ministries of Agriculture (red boxes).

# Carbon budget Indicators

➤ Are calculated for each cropping year (at 10m/plot level), but can be summed over several years (crop rotation),

➤ 3 TIERS:

– TIER 1 (a proxy) and TIER 2 (C budget) are based on empirical approaches and can be applied to most crops species except rice,


– TIER 3 is based on the SAFY-CO2 crop model assimilating LAI derived from Sentinel 2 data ➔ allows other indicators to be calculated (biomass, yield, CO<sub>2</sub> fluxes...) but only for 4 crops species (wheat, sunflower, maize and soon rapeseed) + cover crops at this stage.

➤ A similar conceptual approach:

$$\text{C budget} = \text{Net CO}_2 \text{ flux} - \text{C harvested} + \text{Org. manure}$$

ecosystem convention

➤ Approaches have been discussed with JF Soussana (vice CEO of INRAe, member of IPCC, coordinator of CIRCASA)




# Carbon budget Indicators

➤ Empirical approaches: plot level/annual → most crop species


$$\text{C budget} = \text{Net CO}_2 \text{ flux} + \text{C harvested} - \text{Org. fertil.}$$

TIER 1



Ceschia et al. (2010)

• Reflects the effect of soil coverage (depending on the crop species and the presence of cover crops)




• The longer soil coverage, the better (same conclusion as the French 4/1000 expertise; directly estimated from Sentinel 2 data (NDVI dynamics → Copernicus Phenology service) Pellerin et al., 2019).

# Carbon budget Indicators

➤ Empirical approaches: plot level/annual → most crop species

TIER 1 Operational tool in test phase


Ain department, France



CO2 Flux Calculator

1. Enter csv file :  
Select the File :  Browse ...
2. Enter threshold (Between 0.1 and 0.9) :  
Threshold :  Submit
3. Enter data range (Optional) :  
 Analyse the data over a defined period of time  
Period start :    
Period end :   Submit
4. Process :  
CSV :   
Threshold :   
PROCESS

EXIT





# Carbon budget Indicators


➤ Empirical approaches: plot level/annual → most crop species

TIER 2

$$\text{C budget} = \text{Net CO}_2 \text{ flux} + \text{C harvested} - \text{Org. fertil.}$$

TIER 1

Farmer's data (FMIS)




*Ceschia et al. (2010)*

• What do we need to know from the farmers ?

- **C harvested:**
  - yield (grain t/ha)
  - eventually the amount of straw/cover crop exported (t/ha),
- Are organic amendments applied ? If yes:
  - type of amendment,
  - amount (t/ha).

# Carbon budget Indicators


➤ TIER 3, modelling approach: SAFY-CO2



# Carbon budget Indicators

TIER 3, modelling approach: SAFY-CO2


- This modelling approach was developed in the perspective of the Sentinel data,
- Need very few field data → suited for large scale applications at plot level, but only for a few crop species (wheat, maize, sunflower, rapeseed) + cover crops → to be applied in combination with TIER 2,
- Accounts for the « true effect » of crops/cover crop/regrowth/weeds development on the C budget (only approach that allows it),



- Analysis of large scale transposability during NIVA (research tool → operational MRV tool) → potential use for agricultural C market/Low C label...

# Nitrate leaching indicators

➤ **N biogeochemical** flows exceed planetary boundaries with a major role of agriculture (Campbell et al. 2017) -> N leaching




➤ **Main drivers** to consider (Beaudoin et al., 2005)

## Climate, soil




## Crop rotation



## N management

*N surplus*      *N timing*



➤ **Which indicator?**

➤ Limits of “classical” nitrogen surplus indicator (Bockstaller et al. 2015)

➤ **New approach from DiverImpacts** based on literature (e.g. Beaudoin et al. 2005)

# Nitrate leaching indicators

➤ Are calculated for **each couple previous/current cropping year** at plot level, but can be summed at rotation scale

➤ **2 TIERS:**

➤ TIER 1:

➤ Scoring method (between 0 and 1)

➤ TIER 2:

➤ Weighting factor (between 0 and 1)

## Crop rotation

Previous crop

Catch crop

Crop

Presence of CC  
Development intensity  
F factor

Type of catch  
Crop (FMIS)

Data  
needed



IACS data



Sentinel data 1 & 2



IACS data

Mineralisation

crop residue  
soil after  
previous crop (X2)

N uptake

Crop  
Catch crop

## Soil climate




External data  
Weather-soil  
map

# Biodiversity indicator



- **Method principle:**

- ✓ There is a good correlation between landscape structure + practices and biodiversity
- ✓ Biodiversity is correlated with pollination, biological control, cultural services
- ✓ FarmLand project on multidiversity (synthetic index plants, arthropods, birds) in 8 regions and 5 countries



# Biodiversity indicator



- **Calculation levels:**

Indicator will be assessed at the **landscape level**

**+ farm level** = potential contribution of the farm to landscape heterogeneity and biodiversity levels

- **Calculation period:**

- A cultural year

- Mid-October (year n-1) to mid-October (year n)

- **Variables:**

## Semi-natural habitats



*Including Agro-ecological infrastructures*

## Grassland



*Including fallows*

## Crop diversity



*Including catch crops*


## Field size



## Practices (organic/conventional)



*Impact on*



# Biodiversity indicator

- **Input data**

- LPIS: field size + crops + grassland and fallow land + organic
- Topographical data: semi-natural features

➤ **TIER 1:** proportion of SNH

## Prop of Semi-Natural Habitats



*Agro-ecological infrastructures*  
*Crops + cover crops*  
*Artificial surfaces (buildings...)*

➤ **TIER 2:** proportion + type of SNH  
(Advanced TIER 2: + farming intensity )

## Prop+Type of SNH



*Woods, hedges, grasslands, ponds...*  
*Crops + cover crops*  
*Artificial surfaces (buildings...)*

## Farming intensity



*Impact on*



*Weeds*



*Natural enemies*  
*Pollinators etc...*



# Biodiversity indicators

➤ Preserve landscape and biodiversity (Obj. 3 of new CAP) to promote pollinators services (Hass et al. 2018), biological control (Rush et al. 2010), crop production (Dainese et al. 2019), cultural services (Assandri et al. 2018)



➤ Indicator will be assessed at the landscape level

+ **farm level** = contribution of the farm to landscape and biodiversity levels in a given context

➤ **Main drivers** to consider (Holland et al. 2017, Sirami et al. 2019):

**Farming intensity**

**Type of crop and rotation**

**Field size**

**Crop diversity**

**Semi-natural habitats (SNH)**



Impact on



Including fallows



Including Agro-ecological infrastructures



Weeds




Natural enemies  
Pollinators etc...

# Biodiversity indicators

## ➤ 2 TIERS:

➤ **TIER 1:** proportion of semi natural habitat (SNH)


**Prop of SNH**



*Agro-ecological infrastructures*  
*Crops + cover crops*  
*Artificial surfaces (buildings...)*

➤ **TIER 2:** proportion of SNH + diversity of SNH

**Prop/type of SNH**




*Woods, hedges, grasslands, ponds...*  
*Crops + cover crops*  
*Artificial surfaces (buildings...)*


## Optional

➤ **TIER 3:** proportion of SNH + diversity of SNH + farming intensity

**Prop/type of SNH**



**Farming intensity**



# Conclusions

- We are working on 3 indicators that could be implemented operationally at plot/landscape levels all over Europe and then aggregated at relevant level,
- They address 3 objectives of the CAP and 3 categories of environmental issues/ecosystem services,
- TIER 1 approaches could easily be implemented everywhere thanks to the IACS data + the Sentinel data → use of the new Copernicus services (i.e. Phenology, Cropland),
- TIERS 2 and 3 are more complex to implement (require FMIS and/or external data) but they offer higher levels of accuracy/reliability,
- The 3 indicators will be implemented on test areas (France, Spain, Denmark...) within the frame of the UC1b of the NIVA project.

# THANKS for Your attention !



This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement no. 842009