



HAL
open science

Discrete simulations of granular materials with a Level Set shape description

Jérôme Duriez, Cedric Galusinski

► **To cite this version:**

Jérôme Duriez, Cedric Galusinski. Discrete simulations of granular materials with a Level Set shape description. Particles 2021, Oct 2021, Hamburg, Germany. hal-04222019

HAL Id: hal-04222019

<https://hal.inrae.fr/hal-04222019>

Submitted on 28 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Discrete simulations of granular materials with a Level Set shape description

Jérôme Duriez^{*,†} and Cédric Galusinski[‡]

[†] INRAE, Aix Marseille Univ, RECOVER, Aix-en-Provence, France

* e-mail: jerome.duriez@inrae.fr, web page: <https://www6.paca.inrae.fr/recover/Membres-du-laboratoire/Pages-personnelles/Jerome-DURIEZ>

[‡] IMATH, Université de Toulon, CS 60584, 83041 Toulon Cedex 9, France

ABSTRACT

Numerical modelling of granular materials classically adopt Discrete Element Methods (DEM) that provide a direct description of their discrete nature, together with all mechanical traits related with the latter. The Level Set-Discrete Element Method (LS-DEM) recently pushed DEM even closer to generic shapes, possibly mimicking real particles of a granular material [1]. LS-DEM relies onto a discrete form of the signed distance function to a particle's surface, indirectly describing the latter as the zero level set of the former, as well as onto boundary nodes being exactly located on a grain surface for the purpose of contact treatment.

The precision of LS-DEM is logically impacted by the resolution of the particle-attached grids and by the number and locations of boundary nodes. Computational costs then increase with the required precision and these are investigated by comparing LS-DEM with respect to DEM in the ideal case of spherical particles where a reference solution can be obtained, for a LS-DEM implementation based on the YADE code [2]. On the one hand, memory (e.g. RAM) cost increases, in line with the grid resolution mostly, from the order of megabytes to the order of gigabytes for a 3D REV. On the other hand, time costs are directly impacted by the loop over boundary nodes during contact treatment, turning the duration of a sequential triaxial simulation from hours into days [3]. This logical surge, by two or three order of magnitudes, of computational costs can nevertheless be alleviated by a simple OpenMP parallelization [3] and, possibly, algorithmic modifications [4].

Coming along increased computational costs, the versatility of the method is finally evidenced looking at the mechanical behaviour of superquadric particles, whose LS-DEM simulation is shown to be lighter than an alternative non-spherical description based on polyhedra.

REFERENCES

- [1] R. Kawamoto, E. Andò, G. Viggiani and J. E. Andrade, Level set discrete element method for three-dimensional computations with triaxial case study. *J. of the Mechanics and Physics of Solids*, Vol. 91, p. 1–13, 2016.
- [2] V. Šmilauer et al., Yade Documentation 2nd ed., *The Yade Project*, doi:10.5281/zenodo.34073 (<http://yade-dem.org/doc/>), 2015.
- [3] J. Duriez and S. Bonelli, Precision and computational costs of Level Set-Discrete Element Method (LS-DEM) with respect to DEM. *Computers and Geotechnics*, doi:10.1016/j.compgeo.2021.104033, 2021.
- [4] J. Duriez and C. Galusinski, Level set representation on octree for granular material with arbitrary grain shape. In David Šimurda and Tomáš Bodnár, editors, *Proceedings Topical Problems of Fluid Mechanics 2020*, p. 64–71, Prague, 2020.